

Geomorphology

Endogenic & Exogenic Forces

Introduction

Geomorphology is the study of the earth's surface features and how they are shaped by various forces. These forces can be divided into two main categories: endogenic (internal) and exogenic (external). Together, these forces create the physical and chemical changes that shape the earth's landscape, and are referred to as geomorphic processes.

- Endogenic geomorphic processes, such as diastrophism and volcanism, occur within the earth's interior. Diastrophism encompasses the deformation of the earth's crust caused by tectonic forces, while volcanism includes the formation of volcanic landforms due to the eruption of molten materials.
- On the other hand, exogenic geomorphic processes take place on the earth's surface and include weathering, mass wasting, erosion, and deposition. Weathering is the breakdown of rocks and minerals by chemical and physical processes. Mass wasting refers to the movement of soil, rock, and other materials downslope under the influence of gravity. Erosion is the removal of earth materials by agents such as wind, water, ice, and gravity, while deposition is the accumulation of these materials in new locations.
- Geomorphic agents are the mobile mediums, like running water, moving ice masses, wind, waves, and currents, that facilitate the removal, transportation, and deposition of earth materials. These agents play a significant role in shaping the earth's surface through the various geomorphic processes.

Endogenic Forces


- The forces or movements within the Earth's crust are generated through the interaction of matter and temperature. These Earth movements can be broadly classified into two categories: diastrophism and sudden

movements. The primary driving force behind these endogenic geomorphic processes is the energy that originates from within the Earth.

- This internal energy is primarily produced by radioactivity, rotational and tidal friction, as well as the primordial heat that has been present since the Earth's formation. Due to geothermal gradients and heat flow, this energy causes diastrophism (slow and gradual deformation of the Earth's crust) and volcanism (volcanic activity) within the lithosphere.

Diastrophism

- Diastrophism is the general term applied to **slow bending, folding, warping, and fracturing**.
 - **Warp**==make or become bent or twisted out of shape, make abnormal; distort.
- All processes that move, elevate or build up portions of the earth's crust come under diastrophism.
- **They include:**
 - **orogenic processes** involving mountain building through severe folding and affecting long and narrow belts of the earth's crust; In the process of orogeny, the crust is severely deformed into folds.
 - **epeirogenic processes** involving uplift or warping of large parts of the earth's crust;
- Due to epeirogeny, there may be simple deformation. **Orogeny is a mountain-building process whereas epeirogeny is a continental building process.**
- Through the processes of orogeny, epeirogeny, earthquakes, and plate tectonics, there can be faulting and fracturing of the crust. All these processes cause **pressure, volume, and temperature (PVT) changes which in turn induce metamorphism of rocks.**



Diastrophism

Which of the following is NOT an example of an endogenic geomorphic process? **A. Diastrophism B. Volcanism C.**

Weathering **D. Earthquakes**

Epeirogenic or Continent Forming

Movements [Vertical Movements]

- Epeirogenic movement is a geological process that involves the gradual rising or sinking of large landmasses, such as continents, over extended periods of time. These movements exhibit long undulations or wavelengths and minimal folding in the earth's crust. Cratons, the stable and ancient central portions of continents, are particularly susceptible to epeirogeny.
- These epeirogenic movements are also referred to as radial movements since they act along the earth's radius. They can either cause the land to subside (move closer to the center of the earth) or uplift (move away from the center of the earth). The consequences of epeirogenic movements can often be observed in the geological features and landforms of affected areas.

Uplift

- Evidence of uplift can be seen in features such as raised beaches, elevated wave-cut terraces, sea caves, and fossiliferous beds found above sea level. In various locations along the coasts of Kathiawar, Nellore, and Thirunelveli, raised beaches can be found, with some elevated as much as 15 to 30 meters above the current sea level.
- There are also several instances where places that were once on the coast are now located inland. For example, Coringa, which is situated near the mouth of the Godavari River, Kaveripattinam in the Kaveri Delta, and Korkai on the Thirunelveli coast were all thriving seaports approximately 1,000 to 2,000 years ago. These areas have shifted inland due to the uplift process, demonstrating the changes that have taken place over time.

Subsidence

- Submerged forests, valleys, and buildings serve as evidence of land subsidence, which is the sinking of


the earth's surface. An example of this occurred in 1819, when a section of the Rann of Kachchh was submerged due to an earthquake.

- The discovery of peat and lignite beds beneath sea level in Tirunelveli and the Sunderbans also indicates subsidence. In addition, the Andaman and Nicobar Islands have been separated from the Arakan coast as a result of the submersion of the land in between them.
- On the eastern side of Bombay island, trees have been discovered embedded in the mud approximately 4 meters below the low watermark. A similar submerged forest has been observed on the Thirunelveli coast in Tamil Nadu.
- A significant portion of the Gulf of Mannar and Palk Strait is shallow, indicating that it has been submerged in relatively recent geological times. Furthermore, part of the ancient town of Mahabalipuram, located near Chennai (formerly Madras), is now submerged in the sea.

Orogenic or the Mountain-Forming

Movements [Horizontal Movements]

- Orogenic or the mountain-forming movements act **tangentially to the earth's surface**, as in plate tectonics.
- Tensions** produce **fissures** (since this type of force acts away from a point in two directions) and **compression** produces **folds** (because this type of force acts towards a point from two or more directions).
- In the landforms so produced, the structurally identifiable units are difficult to recognize.
- In general, diastrophic forces that have uplifted lands have predominated over forces that have lowered them.



Ocean-Continent Convergence

Sudden Movements

- Rapid changes in the Earth's surface, also known as sudden geomorphic movements, predominantly take

place at the boundaries of lithospheric plates, or tectonic plate margins. These areas are highly unstable due to the immense pressure generated by the pushing and pulling of magma within the Earth's mantle, which is driven by convection currents.

- As a result, these movements can cause significant alterations to the Earth's surface within a relatively short timeframe.

Earthquakes

- An earthquake occurs when the built-up stress in the Earth's interior rocks is released through weak zones on the planet's surface. This release of energy results in wave motions that cause vibrations, which can sometimes be devastating, on the Earth's surface. These movements can lead to various consequences, such as an uplift in coastal areas. For example, an earthquake in Chile in 1822 resulted in a one-meter rise in the coastal regions.
- Other effects of earthquakes may include alterations in land contours, changes in river courses, and the creation of tsunamis (seismic waves generated in the ocean by an earthquake, as they are known in Japan), which can lead to changes in shorelines. Additionally, earthquakes can trigger remarkable glacial surges (as observed in Alaska), landslides, soil creep, and mass wasting, among other phenomena.

Volcanoes


- Volcanism involves the movement of molten rock, known as magma, towards or onto the Earth's surface. It also includes the formation of various intrusive and extrusive volcanic forms. A volcano is created when

magma from the Earth's interior escapes through vents and fissures in the crust, accompanied by steam, gases (such as hydrogen sulfide, sulfur dioxide, hydrogen chloride, carbon dioxide), and pyroclastic material. The shape of a volcano can vary depending on the chemical composition and viscosity of the lava.

- Pyroclastic flow is a dense, rapidly moving mixture of solidified lava pieces, volcanic ash, and hot gases, which occurs during certain volcanic eruptions. This flow is extremely hot and can incinerate anything in its path, potentially moving at speeds of up to 200 meters per second.

Exogenic (Exogenetic) Processes

- External forces, also known as exogenic forces, originate from outside the Earth's interior and primarily occur within its atmosphere. These forces are responsible for wearing down the Earth's surface, which is why they are referred to as land wearing forces.
- Exogenic processes are the result of stress applied to Earth's materials due to various forces that are generated by the heat from the sun. Stress, defined as the force applied per unit area, is created by pushing or pulling on a solid object.
- The development of stresses in Earth's materials ultimately leads to weathering, erosion, and deposition. Two key climatic factors that influence these processes are temperature and precipitation, as they play a significant role in inducing stress on Earth's materials.



Denudation

- All the exogenic geomorphic processes are covered under a general term, denudation.

- The word '**denude**' means to **strip off or to uncover**.
- **Weathering, mass wasting/movements, erosion, and transportation are included in denudation.**
- Denudation mainly depends on rock type and its structure that includes folds, faults, orientation and inclination of beds, presence or absence of joints, bedding planes, hardness or softness of constituent minerals, chemical susceptibility of mineral constituents; permeability, or impermeability, etc.
- The effects of most of the exogenic geomorphic processes are small and slow but will, in the long run, affect the rocks severely due to continued fatigue.

Denudation: 4 Phases

1. Weathering
2. Erosion
3. Transportation
4. Deposition

What are the two key climatic factors that influence exogenic geomorphic processes?

- A. Temperature and wind
- B. Precipitation and humidity
- C. Temperature and precipitation
- D. Wind and humidity

Conclusion

Geomorphology is the scientific study of the Earth's surface features and the processes that shape them. Endogenic and exogenic forces play crucial roles in shaping the Earth's landscape through various geomorphic processes, such as diastrophism, volcanism, weathering, mass wasting, erosion, and deposition. Endogenic forces originate from within the Earth's interior and are responsible for slow and sudden movements, while exogenic forces are external processes that act on the Earth's surface and are largely influenced by temperature and precipitation. Together, these forces and the geomorphic agents contribute to the continuous transformation and evolution of the Earth's surface features through the complex process of denudation.

Frequently Asked Questions (FAQs) of

Endogenic & Exogenic Forces

What is the main difference between endogenic and exogenic geomorphic processes?

Endogenic geomorphic processes occur within the Earth's interior and are driven by internal energy sources, such as radioactivity and geothermal heat. These processes include diastrophism and volcanism. Exogenic geomorphic processes occur on the Earth's surface and are driven by external forces, such as the sun's heat and precipitation. These processes include weathering, mass wasting, erosion, and deposition.

What are the primary driving forces behind endogenic geomorphic processes?

The primary driving forces behind endogenic geomorphic processes are the energy that originates from within the Earth, such as radioactivity, rotational and tidal friction, and the primordial heat that has been present since the Earth's formation.

What are some examples of epeirogenic movements, and how do they affect the Earth's surface?

Epeirogenic movements involve the gradual rising or sinking of large landmasses, such as continents, over extended periods of time. Examples of epeirogenic movements include uplift, which can cause raised beaches and elevated wave-cut terraces, and subsidence, which can lead to submerged forests, valleys, and buildings.

How do sudden movements, like earthquakes and volcanoes, impact the Earth's surface?

Sudden movements can cause significant alterations to the Earth's surface within a relatively short timeframe. Earthquakes can result in changes to land contours, alterations in river courses, and the creation of tsunamis, while volcanic eruptions can produce various intrusive and extrusive volcanic forms, as well as pyroclastic flows that can incinerate anything in their path.

What factors influence the rate and intensity of exogenic geomorphic processes?

Two key climatic factors that influence exogenic geomorphic processes are temperature and precipitation, as they play a significant role in inducing stress on Earth's materials. Additionally, factors such as rock type, structure, mineral composition, and permeability can also affect the rate and intensity of these processes.

1. What are endogenic forces?

Ans. Endogenic forces are the internal forces that cause changes in the Earth's crust. These forces are generated from within the Earth's surface due to the movement and collision of tectonic plates, volcanic eruptions, and earthquakes.

2. What are exogenic processes?

Ans. Exogenic processes are the external forces that cause changes on the Earth's surface. These processes are caused by factors such as weathering, erosion, deposition, and transportation of materials by wind, water, and ice.

3. What is denudation?

Ans. Denudation is the process of wearing away the Earth's surface due to the action of exogenic forces. It includes weathering, erosion, transportation, and deposition of materials by wind, water, and ice.

4. How do endogenic forces and exogenic processes affect the Earth's surface?

Ans. Endogenic forces and exogenic processes work together to shape the Earth's surface over time. Endogenic forces cause the movement and collision of tectonic plates, which can result in the creation of mountains, valleys, and other landforms. Exogenic processes then work to wear away and reshape these landforms through processes such as erosion and deposition.

5. Why are endogenic and exogenic forces important?

Ans. Endogenic and exogenic forces are important because they play a crucial role in shaping the Earth's surface and creating the diverse environments that support life on our planet. These forces also have significant impacts on human societies, as they can cause natural disasters such as earthquakes, landslides, and floods. Understanding these forces and their effects is important for managing and mitigating the risks associated with these natural hazards.

Physical Condition of Interior of Earth

Introduction - Interior of Earth

In this Geomorphology article, we will delve deeper into understanding the Earth's interior. One might wonder why studying the Earth's interior is important and what benefits can be derived from understanding its basic structure. Here are some reasons:

- Understanding the Earth's interior helps us comprehend the changes that occur on and beneath the Earth's surface.
- It allows us to better understand geophysical phenomena such as volcanism and earthquakes.
- We can gain insights into the internal structure of various objects within the solar system.
- It enables us to understand the evolution and present composition of the Earth's atmosphere.

- It is crucial for future deep-sea mineral exploration and other resource management initiatives.

Now that we understand the importance of studying the Earth's interior, let's discuss the sources that provide knowledge about it and explore the composition of the Earth's interior.

Sources to study the earth's interior

The methods for acquiring knowledge about the Earth's interior can be grouped into two categories:

- **Direct sources**
 - **Mining and drilling:** These activities enable us to directly access and observe the Earth's interior,


allowing us to collect samples and gather information about its composition and structure.

- **Volcanic eruptions:** Eruptions bring materials from deep within the Earth to the surface, providing us with valuable information about the composition and behavior of the Earth's interior.
- **Indirect sources:**
 - **Temperature and pressure variations:** By studying how temperature and pressure change with depth, we can infer information about the Earth's interior, such as its composition, density, and layering.
 - **Meteorites:** Analyses of meteorites can provide information about the composition of the Earth's interior, as they are believed to have originated from the same materials that formed the Earth.
 - **Seismicity:** Seismic waves generated by earthquakes travel through the Earth and are affected by the materials they encounter. By analyzing these waves, we can gain insights into the Earth's interior structure, such as the location of various layers and their properties.

Which of the following discontinuities separates the Earth's crust from the mantle? **A. Conorod Discontinuity B. Mohorovicic Discontinuity (Moho) C. Repiti Discontinuity D. Gutenberg Discontinuity**


Structure of the earth's interior

The structure of the earth's interior is made up of several concentric layers. Structure of Interior of The Earth is divided into three layers- Crust, Mantle, Core



Crust

- The Earth's crust is its outermost solid layer, typically measuring between 30-50 km in thickness. This layer varies in thickness depending on whether it is beneath oceanic or continental regions. The oceanic crust is generally thinner, ranging from 5-30 km, while the continental crust is thicker, measuring between 50-70 km. In areas with major mountain systems, such as the Himalayan region, the continental crust can be as thick as 70-100 km.
- The crust makes up only 0.5-1.0% of the Earth's total volume. Its outer covering is composed of sedimentary material, primarily granitic rocks. Below this layer, there are crystalline, igneous, and metamorphic rocks, which are generally acidic in nature. The lower portion of the crust consists of basaltic and ultra-basic rocks.
- In terms of density, the outer crust has an average density of 2.8, while the lower crust has a density of around 3.0. The continental crust is primarily composed of lighter silicates made of silica and aluminum, also known as "sial." In contrast, the oceanic crust contains heavier silicates made of silica and magnesium, referred to as "sima."
- The boundary between the Earth's lower crust and the upper mantle is known as the Mohorovicic (Moho) discontinuity.



Discontinuities

Mantle

- The mantle refers to the layer of the Earth's interior that lies beyond the crust. It stretches from a depth of about 35 kilometers, known as Moho's discontinuity, down to approximately 2,900 kilometers. The upper section of the mantle, called the asthenosphere, is situated just below the crust. Together, the crust and the uppermost part of the mantle make up the lithosphere, which has a thickness varying between 10 and 200 kilometers.
- Below the asthenosphere lies the lower mantle, which is in a solid state. The mantle is denser than the crust,

with a density ranging from 3.3 to 5.5. It comprises 83 percent of the Earth's total volume and accounts for 68 percent of its total mass. The mantle is primarily composed of silicate minerals that are rich in iron and magnesium.

Asthenosphere

- The upper portion of the mantle is called the asthenosphere.
- It is considered to be extending up to 400 km.
- It is the main source of magma that finds its way to the surface during volcanic eruptions. It has a density higher than the crust.

Core

- The core-mantle boundary is located at the depth of 2900 km.
- **Core lies between 2900 km and 6400 km** below the earth's surface.
- The **outer core is liquid while the inner core is solid**.
- The density of the core is higher than the mantle and varies from 5.5 to 13.6 g/cm³
- The **core is made up of heavy material mostly constituted by nickel and iron [nife]**.
- Volume and mass of core are 16% and 32% of the total volume and mass of the earth respectively.

Earth's Layers- Seismic

Discontinuities

- **Conorod Discontinuity**– between upper and lower crust
- **Mohorovicic Discontinuity (Moho)** – separates the crust from the mantle, its average depth being about 35 km.
- **Repiti Discontinuity** – between the upper and lower mantle
- **Gutenberg Discontinuity** – lies between the mantle and the outer core. Below 2900 km from earth's surface.
- **Lehman Discontinuity**- between inner and outer core.

The Major Elements of the Earth's Crust:

Elements	By Weight (%)
1. Oxygen	46.6
2. Silicon	27.72
3. Aluminium	8.13
4. Iron	5.00
5. Calcium	3.63
6. Sodium	2.83
7. Potassium	2.59
8. Magnesium	2.09
9. Others	1.41

Which of the following elements is NOT a major component of the Earth's crust? **A.** Silicon **B.** Aluminum **C.** Iron **D.** Gold

Conclusion

Understanding the Earth's interior is crucial for various reasons, including predicting geophysical phenomena, studying the evolution of the atmosphere, and managing resources. The Earth's interior is composed of several concentric layers, including the crust, mantle, and core, each with distinct properties and compositions. Knowledge about the Earth's interior is derived from both direct and indirect sources, such as mining, volcanic eruptions, temperature and pressure variations, meteors, and seismicity. Studying these layers and their properties helps us gain valuable insights into the Earth's structure and behavior, contributing to our overall knowledge of the planet and the universe.

Why is studying the Earth's interior important?

Studying the Earth's interior is essential for understanding changes that occur on and beneath the Earth's surface, comprehending geophysical phenomena such as volcanism and earthquakes, gaining insights into the internal structure of objects within the solar system, understanding the evolution and composition of the Earth's atmosphere, and supporting future deep-sea mineral exploration and resource management initiatives.

What are the direct sources for studying the Earth's interior?

Direct sources for studying the Earth's interior include mining and drilling activities, which allow us to directly access and observe the Earth's interior, and volcanic eruptions, which bring materials from deep within the Earth to the surface.

What are the indirect sources for studying the Earth's interior?

Indirect sources for studying the Earth's interior include studying temperature and pressure variations with depth, analyzing

meteorites, and examining seismic waves generated by earthquakes.

What are the three main layers of the Earth's interior?

The three main layers of the Earth's interior are the crust, the mantle, and the core.

The major seismic discontinuities in the Earth's interior include the Conrad Discontinuity (between upper and lower crust), the Mohorovicic Discontinuity (Moho, between the crust and mantle), the Repiti Discontinuity (between upper and lower mantle), the Gutenberg Discontinuity (between the mantle and outer core), and the Lehman Discontinuity (between the inner and outer core).

What are the major seismic discontinuities in the Earth's interior?

1. What are the sources to study the earth's interior?



Ans. The sources to study the earth's interior include seismic waves, which are generated by earthquakes and can provide valuable information about the structure of the earth. Other sources include geophysical surveys, such as gravity and magnetic surveys, which help in mapping the variations in density and magnetic properties within the earth.

2. What is the structure of the earth's interior?



Ans. The earth's interior is divided into several layers. The outermost layer is the crust, which is relatively thin and made up of solid rock. Beneath the crust is the mantle, which is much thicker and consists of solid rock that can flow slowly under pressure. The core is located at the center of the earth and is divided into the outer core, which is liquid, and the inner core, which is solid.

3. What are seismic discontinuities in the earth's layers?



Ans. Seismic discontinuities are boundaries within the earth's layers where seismic waves change their velocity or direction of travel. These discontinuities provide important clues about the composition and density of the materials in the earth's interior. Examples of seismic discontinuities include the Mohorovičić discontinuity (Moho), which separates the crust from the mantle, and the core-mantle boundary, which separates the mantle from the core.

4. How do seismic waves help in studying the earth's interior?



Ans. Seismic waves generated by earthquakes travel through the earth and can be detected and recorded by seismographs. By analyzing the properties of seismic waves, such as their speed, direction, and how they are refracted or reflected at different layers, scientists can infer the composition, density, and temperature variations within the earth's interior. This helps in understanding the structure and dynamics of the earth's interior.

5. What are some frequently asked questions about the physical condition of the earth's interior?



Ans. Some frequently asked questions about the physical condition of the earth's interior include: - What is the temperature of the earth's core? - How do scientists determine the thickness of the earth's crust? - What causes the movement of the tectonic plates? - Is the earth's mantle completely solid? - Can seismic waves travel through the earth's core?

Origin & Evolution of Earth's Crust

Introduction

- The Earth's crust is the thinnest and most fundamental layer of our planet, serving as home to all living organisms that have ever existed. Despite being a relatively simple structure, the crust is highly dynamic and forms a key part of the Earth's various layers.
- Characterized by its distinct chemical composition, the crust can be divided into two main types: oceanic and continental. These differences arise due to plate tectonics, which involve the movement of plates above the asthenosphere. This movement drives lithospheric processes that result in natural phenomena such as earthquakes and ridges. As one of the Earth's five chemical layers, the crust showcases the unique chemical properties found at each level.
- In conjunction with the upper mantle, the Earth's crust plays a crucial role in the ongoing creation and destruction of the surface where all living beings thrive. This section will explore various aspects of the crust, delving into its origin, structure, and composition, as well as shedding light on its continuous evolution up to the present day.

solidification of materials brought to Earth by asteroids and other celestial bodies. While this theory could potentially explain the composition of the oceanic crust, which is primarily composed of basalt, it does not account for the fact that many impact events on Earth occurred after the oceanic crust had already formed. Additionally, the amount of basalt produced by impact events is considered too small to have formed the crust.

- The terrestrial model is currently the most widely accepted explanation for the formation of the Earth's crust. This model posits that the crust originated from the Earth's internal processes. After the late accretion of the Earth, the heat retained by the planet caused the upper mantle to melt, creating a magma ocean that covered the Earth's surface. As the Earth cooled, the magma ocean crystallized, forming a widespread crust. Alternatively, the melted upper mantle could have risen to the surface and solidified, forming the crust.
- The terrestrial model is considered the most likely explanation for the formation of the Earth's crust, as it can account for the uniform composition of the crust as well as its layered structure. The cooling of the magma ocean over time would have resulted in the formation of the different layers observed in the Earth's crust today.

Origin of Earth's Crust

- The Earth's crust is believed to have formed approximately 4.5 billion years ago, after the late stages of planetary accretion. There are three main theories that have been proposed to explain the formation of the crust: the inhomogeneous accretion model, the impact model, and the terrestrial model.
- The inhomogeneous accretion model suggests that the Earth's crust was formed during the accretion of the planet, with lighter and volatile elements forming a thin layer on the primitive planet, which eventually became the crust. However, this model has been criticized for not accounting for the presence of non-volatile elements, such as uranium and thorium, in the Earth's crust.
- The impact model proposes that the Earth's crust was formed through the melting and subsequent

What is the most widely accepted explanation for the formation of the Earth's crust?

- A. Inhomogeneous Accretion Model
- B. Impact Model
- C. Terrestrial Model
- D. Nebular Hypothesis Theory

Nebular Hypothesis Theory

- The Nebular Hypothesis Theory was initially proposed by German philosopher Immanuel Kant and later revised by

mathematician Laplace in 1776. According to the theory, the Sun was encircled by a solar nebula, primarily composed of hydrogen and helium, along with dust particles. The friction and collision of these particles led to the creation of a disk-shaped cloud. Gradually, through the process of accretion, planets were formed from the material surrounding the young Sun.

- However, the Nebular Hypothesis Theory faced several criticisms. For instance, the theory fails to explain the origin of the nebula itself. It also does not account for the sources of heat and motion within the nebula. Furthermore, the hypothesis suggests that since the nebula was gaseous, the planets should also be in a gaseous state, which is not the case as planets are solid. The transition from a gaseous nebula to solid planets is not adequately explained by the theory.
- In addition, the Nebular Hypothesis Theory predicts that all planets and their satellites should rotate in the same direction as the rotation of the nebula. However, this is not observed in reality, as Venus and Uranus rotate in the opposite direction compared to other planets in our solar system.

Planetesimal Hypothesis

The Planetesimal Hypothesis, proposed by Chamberlain and Moulton in 1900, offers an explanation for the origin of Earth and the formation of planets. According to this theory, before the formation of planets, there was only a sun, which was cold at that time. Another sun passed close to our sun, and due to their gravitational attraction, a cigar-shaped bulge of material was pulled from our sun. This bulge eventually separated from the solar surface and formed smaller bodies called planetesimals. These planetesimals then came together and combined to form larger celestial bodies, which we now know as planets.

However, the Planetesimal Hypothesis has been criticized for several reasons:

- The theory was unable to explain why there are only nine planets in our solar system (now recognized as eight, with Pluto being reclassified as a dwarf planet).
- It failed to account for the various sizes of planets in our solar system.
- The distance between the two suns was immense, and it is unlikely that the gravitational pull between them would have been strong enough to extract such significant amounts of material from our sun.

The Big Bang Theory

The Big Bang Theory is a modern scientific explanation that seeks to address various questions about the origin and evolution of the universe, the solar system, the Earth's crust, and the atmosphere. This theory, also known as the expanding universe hypothesis, has significantly advanced our understanding of the cosmos.

- According to the Big Bang Theory, all the matter that makes up the universe was initially concentrated in a single point called the singularity. This point had an incredibly small volume, infinite temperature, and infinite density. Around 13.7 billion years ago, the singularity exploded in a massive event known as the Big Bang. Within the first three minutes after the explosion, the initial atoms began to form as energy was converted into matter. Approximately 300,000 years after the Big Bang, the universe became transparent.
- The formation of stars, as explained by the Big Bang Theory, began about 5 billion years ago. In the early universe, the distribution of matter and energy was uneven. This uneven distribution led to differences in density, which in turn created gravitational forces that caused matter to come together. The universe is composed of a vast number of galaxies, each containing a large number of stars.

A galaxy is a massive collection of stars that formed from nebulae, which are clouds of hydrogen and helium gas. The formation of planets takes place in several stages:

- **Stage 1:** Within the nebula, stars begin as lumps of gas. Gravitational forces within these lumps lead to the formation of a core, surrounded by a huge rotating disc of gas and dust.
- **Stage 2:** The matter around the core starts to condense into smaller, rounded objects called planetesimals. These planetesimals collide with one another, forming larger bodies in the process.
- **Stage 3:** The smaller terrestrial bodies, or planetesimals, continue to combine and form larger planetary bodies.

In summary, the Big Bang Theory is a crucial modern theory that delves into the origin and evolution of the universe, the solar system, and the Earth. This theory provides a comprehensive explanation for the formation of stars, galaxies, and planets, greatly expanding our knowledge of the cosmos.

Which theory provides a comprehensive explanation for the formation of stars, galaxies, and planets in the context of the origin and evolution of the universe?

A. Nebular Hypothesis Theory B. Planetesimal Hypothesis C. Big Bang Theory D. Inhomogeneous Accretion Model

Evolution of earth

- The evolution of Earth can be traced back to when it was initially a barren, rocky, and hot object with a thin atmosphere composed of hydrogen and helium. Over time, Earth developed a layered structure, which is not uniform, due to the gradual increase in density and temperature within the planet. This caused materials to separate based on their density, with heavier elements like iron sinking towards the center and lighter ones moving towards the surface. The formation of the Moon further heated the Earth, leading to the development of three major layers: the crust, mantle, and core.
- The atmosphere and hydrosphere of Earth evolved through three main stages. In the first stage, primordial gases like helium and hydrogen were lost as they were blown away by solar winds. In the second stage, the hot interior of Earth contributed to the development of the atmosphere. As the planet cooled, gases and water vapor were released from its interior into the atmosphere, leading to a composition consisting mainly of water vapor, nitrogen, carbon dioxide, methane, ammonia, and very little free oxygen. This process is known as degassing.
- In the third stage, living organisms played a significant role in modifying the atmosphere's composition. Through the process of photosynthesis, oxygen began to accumulate in the atmosphere. Meanwhile, carbon dioxide levels decreased as it dissolved in rainwater, leading to more condensation and rainfall.
- The origin of life on Earth is believed to be a result of chemical reactions that transformed inorganic materials into living organisms. Around 3.8 billion years ago, life began to evolve, starting with simple organisms like blue-green algae.

Conclusion

The Earth's crust, atmosphere, and hydrosphere have undergone significant transformations since their formation billions of years ago. The terrestrial model, the Big Bang Theory, and the evolution of Earth's atmosphere and hydrosphere provide insights into the processes that have

shaped our planet and its dynamic crust. From the formation of stars, galaxies, and planets to the emergence of life, these theories have greatly expanded our understanding of the cosmos, the origins of Earth, and the complex interactions between its various layers and components.

What are the two main types of Earth's crust?

The Earth's crust can be divided into two main types: oceanic and continental. These differences arise due to plate tectonics, which involve the movement of plates above the asthenosphere.

What is the most widely accepted explanation for the formation of the Earth's crust?

The terrestrial model is currently the most widely accepted explanation for the formation of the Earth's crust. This model posits that the crust originated from the Earth's internal processes, involving the melting and solidification of the upper mantle.

What are the main criticisms of the Nebular Hypothesis Theory?

The Nebular Hypothesis Theory faces several criticisms, including its inability to explain the origin of the nebula itself, the sources of heat and motion within the nebula, and the transition from a gaseous nebula to solid planets. It also does not account for the opposite rotation of Venus and Uranus compared to other planets in our solar system.

How does the Big Bang Theory contribute to our understanding of the formation of the Earth and its crust?

The Big Bang Theory provides a comprehensive explanation for the formation of stars, galaxies, and planets, greatly expanding our knowledge of the cosmos. This theory delves into the origin and evolution of the universe, the solar system, and the Earth, shedding light on the processes that led to the formation of the Earth's crust.

How did life on Earth influence the evolution of the atmosphere?

Living organisms, particularly through the process of photosynthesis, played a significant role in modifying the atmosphere's composition. Oxygen began to accumulate in the atmosphere, while carbon dioxide levels decreased as it dissolved in rainwater. This led to more condensation and rainfall, further shaping the Earth's atmosphere and hydrosphere.

1. What is the origin of Earth's crust?

Ans. The Earth's crust is believed to have originated from the cooling and solidification of molten rock called magma. This process occurred about 4.6 billion years ago during the formation of the Earth.

2. What is the Nebular Hypothesis theory?

Ans. The Nebular Hypothesis theory suggests that the formation of the solar system, including the Earth's crust, occurred from a giant rotating cloud of gas and dust known as the solar nebula. This cloud collapsed under its own gravitational force, causing it to flatten into a spinning disk. The central region of this disk eventually formed the Sun, while the outer regions formed the planets, including the Earth.

3. How does the Big Bang Theory relate to the origin of Earth's crust?

Ans. The Big Bang Theory explains the origin of the universe itself, not specifically the Earth's crust. According to this theory, the universe began as a singularity, an infinitely hot and dense point, approximately 13.8 billion years ago. It then expanded rapidly and continues to expand to this day. The formation of the Earth's crust occurred much later, through processes such as accretion and differentiation.

4. What is accretion in the context of the origin of Earth's crust?

Ans. Accretion refers to the process by which small particles, such as dust and debris, come together to form larger objects. In the context of the Earth's crust, accretion involved the collision and sticking together of these particles, forming planetesimals. Over time, these planetesimals collided and merged to form the proto-Earth, which eventually differentiated into its core, mantle, and crust.

5. How does the origin and evolution of Earth's crust relate to the UPSC exam?

Ans. The origin and evolution of Earth's crust is a topic of importance in the UPSC exam, particularly in the Geology and Geography sections. Understanding the formation and composition of the Earth's crust is crucial for comprehending various geological processes, plate tectonics, and the distribution of natural resources. It is important for candidates to have a solid understanding of these concepts to perform well in the exam.

Fundamentals of Geomagnetism

Introduction


Geomagnetism is a relatively recent field of study in geomorphology that has significantly expanded our understanding of geological processes and the Earth's interior. This area of study is attributed to William Gilbert and focuses on the dynamics of the Earth's magnetic field, which originates in the outer core.

- The Earth's magnetic field primarily acts as a geo-axial dipole, with its north and south magnetic poles located near the geographic poles. Throughout the Earth's geological history, its magnetic field has shifted and reversed, typically at a slow rate of around 9 km per year. However, since the beginning of the 21st century, this speed has accelerated to 50 km per year.
- The Earth functions as a dipole magnet, with its geomagnetic south pole near the Earth's geographic

north and vice versa. The magnetosphere, the region where the Earth's magnetic field can be detected, extends around 60,000 km above the Earth's surface. This magnetic field reaches up to 3,200 km away from the Earth's surface, helping to deflect solar wind and protect the planet from potential catastrophes.

- The Earth's magnetic field serves as a shield against solar flares and harmful gamma rays from the sun. Interestingly, there have been periods in the Earth's geological past when its magnetic field disappeared, and these periods often coincide with major extinction events on the planet.

In summary, geomagnetism is a vital area of study that investigates the Earth's magnetic field and its impacts on the planet's geological processes and protection from cosmic threats. The field has contributed greatly to our understanding of the Earth's interior and the dynamics of its magnetic field, which is crucial for the planet's safety and preservation.



Earth's Magnetic Field


Magnetic Poles & Geomagnetic Poles

Magnetic Poles and Geomagnetic Poles are two terms used to describe different aspects of Earth's magnetic field.

- A magnet's North pole is the pole that is attracted to Earth's North Magnetic Pole when allowed to rotate freely. Since opposite poles attract each other, Earth's North Magnetic Pole is actually the south pole of its own magnetic field. Earth's magnetic field, which resembles a simple bar magnet's north-south field, is typically closely aligned with its rotational axis.
- This means that the magnetic poles are usually quite close to the geographic poles (the poles through which Earth's axis passes), which is why compasses work effectively.


- However, the magnetic field's dipole component reverses every few thousand years, causing the locations of the north and south magnetic poles to switch. Currently, Earth's magnetic North Pole is rapidly moving from the Canadian Arctic towards Russia.

The Earth's Magnetic Field



Earth's Magnetic Field


- On the other hand, Geomagnetic Poles refer to the points where Earth's surface intersects with the axis of a hypothetical bar magnet placed at Earth's center.
- There is one such pole in each hemisphere, known as the "geomagnetic north pole" and the "geomagnetic south pole."
- The geomagnetic dipole is currently tilted at an angle of about 11 degrees to Earth's rotational axis. The magnetic poles, however, are the points where magnetic needles become vertical.
- The difference in positions between the magnetic poles and geomagnetic poles is due to Earth's uneven and complex magnetic field distribution.



North Pole

- Geomagnetic poles are antipodal points where the axis of a best-fitting dipole intersects Earth's surface.
- This theoretical dipole is akin to a powerful bar magnet at Earth's center and provides the most accurate model for describing the magnetic field observed on Earth's

surface. In contrast, Earth's actual magnetic poles are not antipodal, meaning the line connecting them does not pass through Earth's center.



What is the difference between Magnetic Poles and Geomagnetic Poles?

- A.** Magnetic Poles are fixed points, while Geomagnetic Poles change position over time.
- B.** Magnetic Poles are antipodal points, while Geomagnetic Poles are not.
- C.** Geomagnetic Poles are points where a hypothetical bar magnet intersects Earth's surface, while Magnetic Poles are points where magnetic needles become vertical.
- D.** Geomagnetic Poles are related to Earth's rotation, while Magnetic Poles are related to Earth's magnetic field.

Geomagnetic Reversal

A geomagnetic reversal refers to a change in the Earth's magnetic field where the positions of magnetic north and magnetic south switch places. This phenomenon has been studied through paleomagnetism, which examines the magnetism in rocks formed under the influence of Earth's magnetic field at the time of their creation. Over the past 20 million years, it has been observed that the magnetic north and south poles have flipped approximately every 200,000 to 300,000 years.

- However, the reversal process is not as regular as the sun's magnetic field, which reverses every 11 years. On Earth, the time between magnetic reversals can range from as short as 10,000 years to as long as 25 million years, and the reversal process itself can take anywhere from a few hundred to a few thousand years. Throughout the reversal, the magnetic poles emerge at unusual latitudes.
- The Earth's magnetic field has experienced periods of normal polarity, where the predominant direction is the same as it is today, and reverse polarity, where the direction is opposite. The North and South Magnetic Poles shift due to changes in the Earth's magnetic field, and their locations are not fixed. Currently, the North Magnetic Pole is located in northern Canada and is rapidly drifting towards Siberia, while the South Magnetic Pole is positioned off the coast of Antarctica.
- Scientists estimate that the North Magnetic Pole moves about 10 kilometers per year, but recently, its speed has increased to around 50 kilometers per year, potentially reaching Siberia within a few decades. Since the Earth's magnetic field is not perfectly symmetrical, the North and South Magnetic Poles do not have antipodal positions, meaning a straight line drawn between them does not pass through the Earth's center.
- The Earth's North and South Magnetic Poles are also referred to as Magnetic Dip Poles due to the vertical "dip" of the magnetic field lines at these points. For example, if a magnetic compass needle is freely suspended at the magnetic poles, it will point straight down at the North Magnetic Pole (the south pole of Earth's magnetic field) and straight up at the South Magnetic Pole (the north pole of Earth's magnetic field).

Sunspot Theory

- The Sunspot Theory suggests that the Sun, being a massive magnet, can have a significant impact on Earth's magnetic field during periods of increased sunspot activity. When the Sun's magnetic field is waxing or becoming stronger, the solar wind can penetrate Earth's magnetic shield and interact with the planet's magnetic field, provided that Earth's magnetism is in a waning or weakening phase. In this way, Earth can potentially acquire the Sun's magnetic polarity.
- Furthermore, collisions with meteorites or comets can cause a reversal of Earth's magnetic poles. When these impacts occur, the Earth's magnetic field can become

temporarily disrupted and lose its magnetism. As the magnetism re-establishes, the magnetic poles may reverse. This process suggests that external factors, such as solar activity and celestial collisions, can play a significant role in the behavior of Earth's magnetic field.

Magnetic Declination

- Magnetic Declination, also known as the horizontal component, refers to the angle on the horizontal plane between magnetic North and geographic (true) North. By convention, the declination is considered positive when magnetic North is east of true North, and negative when it is to the west. Currently, the magnetic North Pole is slightly west of the geographic North Pole, while the magnetic South Pole is slightly east of the geographic South Pole. This is due to the molten, iron-rich core of the Earth, which spins like a dynamo and causes a gradual westward drift of the geomagnetic field, resulting in ever-changing magnetic declination throughout Earth's geological history. Magnetic declination can be measured using a magnetic compass.
- Magnetic Inclination, also known as Dip Angle or Magnetic Dip, is the vertical component of geomagnetism. It is defined as the angular inclination between a freely suspended magnetic needle and the Earth's horizontal surface. The magnetic dip occurs due to a magnet's tendency to align itself with the line of force. As latitude increases, the dip angle also increases, reaching 90 degrees at the magnetic poles and 0 degrees at the magnetic equator. The inclination can be measured using a dip needle.
- The Magnetic Equator is the line or locus of points where the dip angle is zero, meaning the magnetic force is perfectly horizontal at these locations.


Theories about origin of earth's magnetism

There are several theories about the origin of Earth's magnetism:

1. **Gilbert Theory:** Sir Gilbert first discovered Earth's magnetism in the 1600s. He believed that a bar magnet inside the Earth's interior was responsible for the magnetism. However, this theory is not widely accepted, as it cannot explain how the bar magnet's magnetism

could sustain itself in the high temperatures and molten rocks of the Earth's interior. It also cannot explain the periodic reversal of Earth's magnetic polarity without assuming the rotation of the magnet itself, which is improbable.

2. **Rock Magnetism:** Another theory suggests that Earth's magnetism is due to rock magnetism. While rock magnetism does exist, it is not considered the primary cause of Earth's geomagnetism, as randomly created magnetic rocks cannot explain the specific orientation of Earth's magnetic field. Additionally, the high temperatures of the Earth's interior should have destroyed any magnetism, as no magnetism can maintain itself in such extreme heat and molten fluid state.
3. **Dynamo Theory:** In 1939, Walter Alsec proposed the Dynamo theory, which states that Earth's magnetism may be the result of the unique structure of Earth's core. The core consists of an iron-rich solid inner core, which is magnetic, and an outer core that is hot and fluid, with free electrons in a near plasma state. Due to Earth's rotation, the outer core and its free electrons also rotate around the inner core, creating electricity. This arrangement is called a solenoid. The solenoid generates a magnetic field, which in turn induces a strong electric field, further strengthening the magnetic field. This self-exciting dynamo mechanism is the most widely accepted explanation for Earth's magnetism. However, it is still unable to explain the fluctuation in the magnetic field and reversal of polarity. Even though Earth's rotation has been smooth and consistent throughout its geological history, the Earth's spin has never reversed.



What is the primary cause of Earth's magnetism according to the most widely accepted theory? **A.** A large bar magnet inside the Earth **B.** Rock magnetism **C.** Dynamo Theory **D.** The Sun's magnetic field

surface intersects with the axis of a hypothetical bar magnet placed at Earth's center. The magnetic poles are subject to change and movement, while the geomagnetic poles are more stable and theoretical in nature.

Conclusion

Geomagnetism is a crucial field of study that enhances our understanding of Earth's magnetic field and its role in geological processes and protection from cosmic threats. The Earth's magnetic field, which primarily functions as a geo-axial dipole, has experienced shifts and reversals throughout its history, impacting the planet's magnetic poles and geomagnetic poles. The widely accepted Dynamo Theory attributes the origin of Earth's magnetism to the unique structure and behavior of the planet's core. Studying Earth's magnetic field and its fluctuations can provide valuable insights into the dynamics of the planet's interior and help predict future changes that may affect the planet's safety and preservation.

What is geomagnetism, and why is it important?

Geomagnetism is the study of Earth's magnetic field and its impacts on the planet's geological processes and protection from cosmic threats. It is crucial for understanding the Earth's interior, the dynamics of its magnetic field, and the protection it provides from solar flares and harmful gamma rays.

What is the difference between Magnetic Poles and Geomagnetic Poles?

Magnetic Poles refer to the points on Earth's surface where the magnetic field lines are vertical, near the geographic poles. Geomagnetic Poles, on the other hand, are the points where Earth's

What is a geomagnetic reversal, and how often does it occur?

A geomagnetic reversal is a change in Earth's magnetic field where the positions of magnetic north and magnetic south switch places. Over the past 20 million years, the magnetic north and south poles have flipped approximately every 200,000 to 300,000 years. However, the reversal process is not regular, and the time between reversals can range from as short as 10,000 years to as long as 25 million years.

What is the Dynamo Theory, and how does it explain Earth's magnetism?

The Dynamo Theory suggests that Earth's magnetism is a result of the unique structure of Earth's core, which consists of an iron-rich solid inner core and a hot, fluid outer core with free electrons. Due to Earth's rotation, the outer core and its free electrons also rotate around the inner core, creating electricity and generating a magnetic field. This self-exciting dynamo mechanism is the most widely accepted explanation for Earth's magnetism.

What is magnetic declination, and how is it measured?

Magnetic declination is the angle on the horizontal plane between magnetic North and geographic (true) North. It is considered positive when magnetic North is east of true North and negative when it is to the west. Magnetic declination can be measured using a magnetic compass and is crucial for navigation, as it allows for the correction of compass readings to account for the difference between magnetic and true North.

1. What are magnetic poles and geomagnetic poles?

Ans. Magnetic poles are the regions on a magnet where the magnetic field lines converge or diverge. There are two types of magnetic poles: the north pole and the south pole. Geomagnetic poles, on the other hand, refer to the locations on the Earth's surface where the magnetic field lines are perfectly vertical.

2. What is geomagnetic reversal?

Ans. Geomagnetic reversal, also known as magnetic polarity reversal, is the process in which the Earth's magnetic field flips, resulting in the north and south magnetic poles switching places. This phenomenon has occurred numerous times throughout Earth's history and is recorded in rocks and sediments as magnetic stripes.

3. What are some theories about the origin of Earth's magnetism?



Ans. There are several theories proposed to explain the origin of Earth's magnetism. One theory suggests that it is generated by the movement of molten iron in the outer core of the Earth. Another theory proposes that it is a result of the magnetization of minerals in the Earth's crust. Additionally, some scientists believe that the Earth's magnetic field is influenced by the interaction between the solar wind and the Earth's magnetosphere.

4. How does geomagnetism impact our daily lives?



Ans. Geomagnetism has various impacts on our daily lives. It plays a crucial role in navigation, as compasses rely on the Earth's magnetic field to determine direction. Geomagnetic storms caused by solar activity can disrupt satellite communications, power grids, and GPS systems. Additionally, some animals, such as birds and turtles, use geomagnetism for navigation during migration.

5. How do scientists study geomagnetism?



Ans. Scientists study geomagnetism through various methods. They use magnetometers to measure the strength and direction of the Earth's magnetic field. By analyzing rocks and sediments, they can reconstruct the Earth's magnetic field history and study geomagnetic reversals. Satellites equipped with magnetometers also provide valuable data about the Earth's magnetic field and its variations.

Geosynclines

Introduction to Geosynclines

Geosynclines refer to expansive, elongated areas of shallow water bodies that are surrounded by more stable landmasses. These areas receive massive deposits of sediment from their surrounding regions. They are typically characterized as long, narrow ocean basins that have experienced continuous sediment deposition and subsidence over extended periods.

These sediment deposits within geosynclines eventually become uplifted and folded, leading to the formation of folded mountain ranges. As a result, it is often said that such mountains originate from geosynclines. Examples of geosynclines in geological history include the Tethys Sea in Asia, the Appalachian geosynclines in North America, and the Caledonian geosynclines in Great Britain.

Concept of Geosyncline Theory and Its Development

The concept of geosynclines was first introduced by James Hall and James Dwight Dana and later expanded upon by Émile Haug. Hall's research on the stratigraphy and structure of the northern Appalachians led him to discover that folded Paleozoic sediments in mountain ranges were shallow-water marine types, with a thickness of 12 kilometers. This thickness was ten to twenty times greater compared to unfolded rock strata of corresponding ages found in the interior lowlands to the west.

- Hall's findings suggested that the deposition of massive sequences of shale, sandstone, and limestone indicated that the underlying floor of older rocks had subsided by a similar amount. Mountain formation was preceded by extended periods of down-warping, during which sediment accumulation balanced the subsidence of the

crust. Dana (1873) termed such troughs as "geosynclinal troughs," which later became known as "geosynclines."

- H. Stille developed an important classification of geosynclines, dividing the earth's crust into two primary divisions: Cratons and Ortho-geosynclines. Cratons are further divided into hochkraton (stable continental crust) and tiefkraton (stable oceanic crust). Ortho-geosynclines are subdivided into miogeosynclines and eugeosynclines, with eugeosynclines characterized by intermittent volcanic activity during sedimentation and miogeosynclines having low volcanic activity.
- E. Haug defined geosynclines as deepwater regions with considerable length but relatively narrow width. He created paleogeographical maps of the world to demonstrate that present-day fold mountains originated from the massive geosynclines of the past. Haug identified five major landmasses from the Mesozoic Era and four geosynclines located between these rigid masses.
- According to Haug, the transgressional and regressional phases of seas directly impact the littoral margins of geosynclines. Evans further explained that geosynclines could be found between two landmasses, in front of a mountain or plateau, along continental margins, or in front of a river mouth.

Geosynclinal Orogen Theory of Kober

- The renowned German geologist Kober provided a comprehensive and systematic analysis of the Earth's surface features in his book, 'Der Bau der Erde.' His primary goal was to establish a connection between ancient, rigid landmasses or tablelands and more mobile areas or geosynclines, which he referred to as orogen.
- Kober aimed to explain not only the origins of mountains using his geosynclinal theory but also to delve into various aspects of mountain formation, such as their creation, geological history, evolution, and development. He viewed the old, rigid landmasses as the foundational building blocks for today's continents, asserting that modern continents have evolved from these rigid structures. He described the process of mountain building, or orogenesis, as the mechanism connecting rigid landmasses with geosynclines. In other words, mountains are created from geosynclines as a result of the influence of rigid landmasses.
- Kober's geosynclinal theory is founded on the idea that the Earth's contraction forces, caused by cooling, play a

significant role in its history. As J.A. Steers stated in 1932, "Kober is definitely a constructionist, with contraction providing the motive force for the compressive stress." This means that the contraction forces generated by the Earth's cooling lead to horizontal movements of rigid landmasses or forelands, which in turn compress, buckle, and fold sediments to form mountain ranges.

Base of the Geosynclinal Orogen Theory

Kober proposed that in the past, there were mobile water zones in locations where mountains currently exist. He referred to these mobile water zones as geosynclines or orogens, which are areas where mountain formation occurs. Kober also identified rigid land masses surrounding these geosynclines, which he called kratogens.

Some examples of these old rigid masses include the Canadian Shield, Baltic Shield or Russian Massif, Siberian Shield, Chinese Massif, Peninsular India, African Shield, Brazilian Mass, and the rigid masses of Australia and Antarctica. Kober also suggested that the mid-Pacific geosyncline separated northern and southern landmasses, which later sank to form the Pacific Ocean.

During the Mesozoic era, Kober identified eight distinct morphotectonic units based on the earth's surface features. These units include:

1. **Africa, along with parts of the Atlantic and Indian Oceans**
2. **The combined landmass of India and Australia**
3. **Eurasia**
4. **North Pacific continents**
5. **South Pacific continents**
6. **South America and Antarctica**

Kober recognized six major periods of mountain formation throughout Earth's history. Three of these periods occurred during the pre-Cambrian period, though little is known about them. The Palaeozoic era saw two significant mountain-building periods: the Caledonian orogenesis, which ended during the Silurian period, and the Variscan orogeny, which culminated in the Permo-Carboniferous period. The final (6th) mountain-building event, the Alpine orogeny, was completed during the Tertiary epoch.

Kober believed that mountains were formed from geosynclines. He described geosynclines as long, wide water areas characterized by sedimentation and subsidence, which are the locations where

mountain formation (or orogenesis) occurs. J.A. Steers (1932) stated that Kober's ideas combined the older geosynclinal hypothesis by Hall and Dana (later developed by Haug) with his own views on orogenesis.


What are the three stages or phases of the mountain-building process?

- A. Lithogenesis, Orogenesis, Gliptogenesis
- B. Sedimentation, Compression, Erosion
- C. Deposition, Folding, Uplift
- D. Subsidence, Convergence, Metamorphism

Phases of the Mountain Building

There are three different stages or phases of the mountain-building process.

- **Lithogenesis:** Lithogenesis is the initial stage of mountain formation, characterized by the creation, sedimentation, and subsidence of geosynclines. Geosynclines form as a result of the Earth's cooling process, which leads to contraction. The surrounding stable areas, or forelands, experience erosion, causing rocks and boulders to wear away and be deposited into the geosynclines. As more sediment accumulates, the geosynclines continue to subside, leading to an increased thickness of sediments.




The stage of lithogenesis: creation of geosyncline followed by sedimentation and subsidence.

Lithogenesis


- **Orogenesis:** Orogenesis is the stage in which the sediments in the geosynclines are compressed and

folded, forming mountain ranges. This occurs due to the convergence of forelands towards each other, which is caused by the Earth's contraction. The immense pressure generated by the moving forelands results in the contraction, squeezing, and folding of the sediments that have accumulated in the geosynclines.



The stage of orogenesis: squeezing and folding of geosynclinal sediments due to compressive forces; the whole of geosynclinal sediments are folded when the compressive forces coming from the sides of geosyncline is enormous and acute.

- Orogenesis
- **Gliptogenesis:** The final stage of mountain building, gliptogenesis, involves the gradual rise of mountain ranges and the ongoing erosion processes by natural agents, such as wind, water, and ice. This stage highlights the continuous evolution of the mountains as they are subjected to various geological processes.



Folding of marginal sediments into marginal ranges and formation of median mass when the compressive forces are moderate.

Gliptogenesis

Types of Geosynclines

There are seven types of geosynclines as follows: Geosynclines are elongated depressions in the Earth's crust that accumulate large amounts of sediments and volcanic rocks. These basins eventually undergo deformation, leading to the formation of fold-mountain chains. There are seven types of geosynclines, each with unique features and formation processes:

1. **Ortho-geosynclines:** These are elongated basins that fill up with a significant thickness of sediments before

undergoing deformation, resulting in the formation of a fold-mountain chain.


2. **Eugeosynclines:** These geosynclines contain large amounts of sediments and volcanic rocks and are typically situated away from shield areas or stable regions of the Earth's crust, known as Kratons.
3. **Miogeosynclines:** Forming adjacent to Kratons, these geosynclines have a thinner layer of sediments and lack volcanic rocks.
4. **Taphrogeosynclines:** These elongated depressions result from faulting in the Earth's crust and are also called 'graben' or 'rift-valleys.'
5. **Parageosynclines:** These geosynclines are located within the Kraton itself.
6. **Zeugo-geosynclines:** These are parageosynclines that have marginal uplifts, or areas of the crust that have risen relative to their surroundings.
7. **Auto-geosynclines:** Unlike zeugo-geosynclines, these are parageosynclines that do not have marginal uplifts.

Most mountain-building events, or orogenies, occur at the sites of geosynclines. The resulting mountains typically consist of sediments and volcanic rocks that have been deformed and metamorphosed to varying extents, depending on their position and depth within the orogenic belt.

The Modern Concept of Geosyncline

The concept of geosynclines has evolved significantly with the introduction of Plate Tectonic Theory. Active margins refer to continental margins situated along plate boundaries where subduction, collision, or transform fault motion occurs. Passive margins, on the other hand, are continental margins that move away from a spreading axis.

- An example of a passive margin can be found on the east coast of North America, where sediment deposits accumulate as the continent moves away from the spreading axis. As the lithosphere cools and becomes denser, the ocean floor deepens off the passive margin, and sediments continue to be deposited. This thick sediment accumulation along the border of a passive margin is called a geosyncline.
- Research conducted during the latter half of the 20th century has shown that a geosyncline is a thick, rapidly accumulating body parallel to the continent. The traditional idea of a geosyncline as an intra-cratonic trough bordered by mountains providing sediments is no longer considered accurate. Instead, sediment accumulation can occur on the continental shelf and slope or in a trough or trench.
- The term "geocline" is now used because the structure of a geosyncline is not a two-sided trough, but rather, it is more open towards the ocean. Geoclines of passive continental margins can be classified into two types: miogeoclines, which are wedges of shallow-water sediments of marine origin that make up the continental shelves; and eugeoclines, which are wedges of deep-sea sediment deposited at the base of the continental slope and located on oceanic crust. Both types of geoclines are formed by the accumulation of sediments as the lithosphere slowly subsides.
- In the Gulf of Mexico, miogeocline sediments can reach a thickness of 20 km at the outer edge of the continental shelf. Eugeocline sediments can be found on the oceanic crust above an oceanic volcano. The continuous accumulation of sediments in miogeoclines for about 200 million years has been enabled by the sinking of the crust due to sediment loading. Miogeocline areas are economically significant due to the presence of mineral oil.



The world distribution of young fold mountains, active volcanoes and earthquake zones.

Fold mountain locations

What is the primary difference between miogeoclines and eugeoclines?

- A.** Miogeoclines have a thinner layer of sediments and lack volcanic rocks, while eugeoclines contain large amounts of sediments and volcanic rocks.
- B.** Miogeoclines are formed in deep-sea trenches, while eugeoclines are formed on continental shelves.
- C.** Miogeoclines are situated away from shield areas, while eugeoclines are formed adjacent to Kratons.
- D.** Miogeoclines are found on oceanic crust, while eugeoclines are found on continental crust.

Conclusion

Geosynclines are elongated depressions in the Earth's crust that accumulate large amounts of sediments and volcanic rocks, eventually leading to the formation of fold-mountain chains. The concept of geosynclines has evolved over time, with the introduction of Plate Tectonic Theory further refining our understanding of their formation and classification. Geoclines, which include miogeoclines and eugeoclines, are now understood as thick sediment accumulations along passive continental margins that play a significant role in the geological history of the Earth and the formation of mountain ranges.

What is a geosyncline?

A geosyncline is an expansive, elongated area of shallow water bodies that receive massive deposits of sediment from their surrounding regions. They are characterized as long, narrow ocean basins that have experienced continuous sediment deposition and subsidence over extended periods.

How do geosynclines contribute to the formation of mountain ranges?

Sediment deposits within geosynclines eventually become uplifted and folded, leading to the formation of folded mountain ranges. As a result, it is often said that such mountains originate from geosynclines.

Who first introduced the concept of geosynclines?

The concept of geosynclines was first introduced by James Hall and James Dwight Dana, and later expanded upon by Émile Haug.

What are the three stages of the mountain-building process?

The three stages of the mountain-building process are lithogenesis (creation, sedimentation, and subsidence of geosynclines), orogenesis (compression and folding of sediments in geosynclines, forming mountain ranges), and gliptogenesis (gradual rise of mountain ranges and ongoing erosion processes).

How has the concept of geosynclines evolved with the introduction of Plate Tectonic Theory?

The concept of geosynclines has evolved significantly with the introduction of Plate Tectonic Theory, which has led to the understanding of active and passive continental margins. This has changed the view of geosynclines as intra-cratonic troughs bordered by mountains providing sediments to a more open structure, with thick sediment accumulation occurring on the continental shelf and slope or in a trough or trench.

1. What is a geosyncline?



A geosyncline is a trough-like depression in the Earth's crust that forms due to the accumulation of sedimentary rocks over millions of years. It is often associated with the formation of mountain ranges and is characterized by the presence of thick sedimentary deposits.

2. What is the geosyncline theory and how did it develop?



The geosyncline theory, also known as the geosynclinal theory, is an early geological concept that explains the formation of mountain ranges through the gradual accumulation and subsequent folding and uplifting of sedimentary rocks in geosynclines. It was first proposed by James Hall in the 19th century and further developed by geologists such as James Dana and Charles Schuchert.

3. What are the phases of mountain building in geosynclines?



The phases of mountain building in geosynclines typically involve three stages: the geosynclinal stage, the eugeosynclinal stage, and the miogeosynclinal stage. During the geosynclinal stage, sediments accumulate in the geosyncline. In the eugeosynclinal stage, intense folding, faulting, and volcanic activity occur, leading to the formation of a mountain range. Finally, in the miogeosynclinal stage, erosion and sedimentation processes dominate, causing the gradual subsidence of the mountain range.

4. What are the types of geosynclines?



There are two main types of geosynclines: oceanic geosynclines and continental geosynclines. Oceanic geosynclines occur in ocean basins and are associated with the formation of island arcs and volcanic mountain ranges. Continental geosynclines, on the other hand, occur on continental margins and are characterized by the accumulation of thick sedimentary sequences.

5. What is the modern concept of geosyncline?



The modern concept of geosyncline has evolved from the original geosyncline theory and is now understood in the context of plate tectonics. Geosynclines are seen as areas of crustal subsidence where sedimentary rocks accumulate, often in association with subduction zones or continental collisions. They are considered as dynamic features that play a role in the formation of mountain ranges and the overall evolution of the Earth's crust.

Continental Drift

Introduction - Continental Drift

Alfred Wegener, a multifaceted German scientist with expertise in meteorology, polar exploration, astronomy, and geology, is widely regarded as the father of the continental drift theory. In 1912, during a lecture, Wegener introduced his groundbreaking concept of 'continental drift,' which significantly changed our understanding of the Earth's geological processes.

Continental Drift Theory (Alfred Wegener)

The Continental Drift Theory was proposed by Alfred Wegener, who suggested that all of Earth's land was once combined into a single supercontinent, which he called "Pangaea" (meaning "all Earth"), surrounded by an enormous ocean named "Panthalassa" (meaning "all ocean"). According to Wegener's theory, this massive landmass began breaking apart approximately 200 million years ago, with the pieces gradually

moving to their current positions and continuing to shift even today.

Wegener became interested in this idea due to several intriguing questions he had:

1. How could tropical ferns have grown in locations such as London, Paris, Bonn, and even Greenland?
2. Why are coal belts found in the extremely cold tundra regions?
3. How can evidence of glaciers be found in tropical areas like Brazil, the Indian Peninsula, Australia, and the Congo Basin?

Considering these points, Wegener came up with two possible explanations:

- The climate zones may have shifted from one region to another, while the continents remained stationary.
- The climate zones stayed in place, but the continents moved.

Since climate zones are determined by factors such as the Earth's tilt and the sun's position, it seemed unlikely that they would shift. Therefore, Wegener concluded that the more plausible explanation was that the continents had changed their positions over time.

What were the two primary directions of continental drift, according to Alfred Wegener's theory? **A.** Northward and Eastward **B.** Southward and Westward **C.** Towards the Equator and Westward **D.** Towards the Equator and Eastward

The Direction of Drift


The theory of continental drift, proposed by Alfred Wegener, suggests that the continents have moved in two primary directions:

- **Towards the Equator:** Wegener attributed the equatorward drift to the Earth's rotation and equatorial bulge, as well as the gravitational pull of the moon and the sun. This drift pushed Africa and Eurasia closer together, raising the Tethys Sea deposits between them into the folded mountains of the Alps, Atlas, Tien Shan, Zagros, Hindukush, and the Himalayas. Additionally, the Indian and African peninsulas separated from Australia and Antarctica.

- **Towards the West:** The westward drift caused North America and South America to separate from Europe and Africa, resulting in the formation of the Atlantic Ocean.

Evidence supporting the Continental Drift Theory includes:

- **Jigsaw Fit:** The shapes of the continents, especially the coastlines of Africa and South America, fit together like pieces of a puzzle, indicating that they were once joined.



- **Geological Structure:** The geological structures on the continents' coastlines show remarkable similarities, suggesting they were once connected. For example, the Appalachian Mountains in North America align with the old Hercynian Mountains in Ireland, Wales, and central Europe.



Geological Structure

- **Permo-Carboniferous Glaciations:** Evidence of ancient glaciations is found in Brazil, Falkland Islands, South Africa, the Indian Peninsula, and Australia. This suggests that these landmasses were once joined, as it is difficult to explain these glaciations based on the current distribution of land and water.
- **Fossil Remains:** Similar fossil remains of terrestrial animals are found on both sides of the Atlantic, suggesting that the continents were once connected. It would have been impossible for these animals to swim across the Atlantic Ocean.
- **Paleoclimatic Evidence:** Coal deposits have been discovered in temperate and polar regions, even though coal is formed in tropical regions. This indicates that the continents have shifted over time.
- **Biological Evidence:** Some animal species, such as the lemming, display migratory patterns that suggest the continents were once joined. Lemmings migrate westward across North America and fall into the Atlantic Ocean, possibly following an ancient route that connected North America and Europe.

Criticism of Continental Drift Theory

The continental drift theory, while quite persuasive, also faced significant criticism and controversy, largely due to the speculative nature of some aspects of the theory and insufficient supporting evidence. The most contentious part of this theory was the proposed forces that caused the drift.

- Experts argued that if the gravitational forces of the moon or sun were strong enough to break up the landmass, it would have also stopped the Earth's rotation, making it stationary. Additionally, the rotational speed required to cause a drift in the landmass would have been so high that it would have flung the atmosphere and everything else in outer space away from Earth's gravitational pull.
- Another issue with the theory was the lack of knowledge about the Earth's history before the Carboniferous period. Furthermore, the proposed drift only considered northward and westward movements, without explaining why other directions were not involved.
- The idea of the sial (continental crust) floating over the sima (oceanic crust) was also criticized, as it was later discovered that the lithosphere actually floats on the asthenosphere. The explanation for the formation of

mountains like the Rockies and Andes due to friction with the sima was considered self-contradictory.

- Finally, the continental drift theory failed to explain the formation of oceanic ridges and island arcs. Overall, while the theory was groundbreaking at the time, it faced significant criticism, which later led to the development of the more comprehensive plate tectonics theory.

Which of the following is NOT a piece of evidence supporting the Continental Drift Theory? **A.** Jigsaw Fit of the continents **B.** Similar geological structures on coastlines **C.** Permo-Carboniferous Glaciations **D.** The Earth's stationary rotation

Conclusion

Alfred Wegener's continental drift theory significantly changed our understanding of the Earth's geological processes. The theory suggests that Earth's landmasses were once combined into a single supercontinent, Pangaea, which broke apart and moved to their current positions over time. While the theory provided an explanation for various geological, paleoclimatic, and biological evidence, it faced significant criticism due to its speculative nature and insufficient supporting evidence. These criticisms eventually led to the development of the more comprehensive plate tectonics theory.

What is the Continental Drift Theory and who proposed it?

The Continental Drift Theory was proposed by German scientist Alfred Wegener. It suggests that all of Earth's land was once combined into a single supercontinent called Pangaea, which later broke apart and moved to their current positions, continuing to shift even today.

What were the two main directions of continental drift according to Wegener's theory?

Wegener's theory suggests that the continents have moved in two primary directions: towards the Equator and towards the West. The equatorward drift contributed to the formation of various mountain ranges, while the westward drift led to the separation of the Americas from Europe and Africa, and the formation of the Atlantic Ocean.

What evidence supports the Continental Drift Theory?

Some key evidence supporting the Continental Drift Theory includes the jigsaw fit of continents' coastlines, similarities in geological structures, evidence of Permo-Carboniferous glaciations, fossil remains, paleoclimatic evidence, and biological evidence such as migratory patterns of certain animal species.

Why was the Continental Drift Theory criticized?

The theory faced significant criticism mainly due to the speculative nature of some aspects, insufficient supporting evidence, and the inability to explain certain geological phenomena. Critics argued that the proposed forces causing the drift were not plausible, and the theory could not account for the Earth's history before the

Carboniferous period or explain the formation of oceanic ridges and island arcs.

How does the Continental Drift Theory relate to the Plate Tectonics Theory?

The Continental Drift Theory was a precursor to the Plate Tectonics Theory. While the former focused on the movement of continents, the latter expanded upon this idea by considering the movement of Earth's lithospheric plates. The criticisms of the Continental Drift Theory led scientists to develop the more comprehensive Plate Tectonics Theory, which is now widely accepted as the explanation for Earth's geological processes.

1. What is continental drift and how does it explain the movement of continents?

Ans. Continental drift is a scientific theory that suggests that the Earth's continents were once joined together as a single landmass called Pangaea and have since drifted apart over millions of years. This theory explains the movement of continents by proposing that the Earth's lithosphere, which is composed of tectonic plates, moves slowly over the semi-fluid asthenosphere. This movement is driven by convection currents in the mantle.

2. What evidence supports the theory of continental drift?

Ans. There is various evidence that supports the theory of continental drift. One of the key pieces of evidence is the fit of the continents' coastlines, particularly the east coast of South America and the west coast of Africa. The similarity of rock types and geological structures across continents, such as the Appalachian Mountains in North America and the Caledonian Mountains in Europe, also supports the theory. Additionally, the distribution of fossils and the presence of similar plant and animal species on different continents provide further evidence for continental drift.

3. What is the significance of the direction of drift in continental drift theory?

Ans. The direction of drift in continental drift theory is significant as it helps explain the movement of continents and the formation of various geological features. The theory suggests that continents have moved away from each other, forming ocean basins in the process. The direction of drift provides insights into the history of Earth's landmasses, the formation of mountain ranges, and the distribution of plant and animal species across continents.

4. What are some criticisms of the continental drift theory?

Ans. The continental drift theory has faced criticism in the past, primarily due to the lack of a convincing mechanism to explain the movement of continents. Critics argued that the theory lacked a solid scientific foundation until the discovery of plate tectonics in the 1960s provided a mechanism for continental drift. Additionally, some geologists initially resisted the theory as it challenged the prevailing belief in static continents. However, with the accumulation of more evidence and the development of plate tectonics, the continental drift theory gained widespread acceptance in the scientific community.

5. How does the theory of continental drift contribute to our understanding of Earth's geological history?

Ans. The theory of continental drift has revolutionized our understanding of Earth's geological history. It has helped explain the formation of various geological features, such as mountain ranges and ocean basins. By studying the distribution of fossils and plant and animal species across continents, scientists have been able to reconstruct past environments and track the movement of landmasses. This theory has also provided insights into the formation of supercontinents, the opening and closing of ocean basins, and the evolution of Earth's climate over millions of years.

Isostasy

Define Isostasy

Isostasy, derived from the Greek words 'iso-stasios' meaning 'equal standing,' is a term first proposed by American geologist Clarence Dutton in 1889. It refers to the state of balance that exists between the Earth's large upstanding features, such as mountain ranges and plateaus. The theory behind isostasy suggests that the Earth's lighter surface materials (sial) float on top of the denser magma (sima) found in the Earth's interior.

- As we explore deeper into the Earth, we find various concentric layers, with the densest material forming the core and the lightest material making up the Earth's surface. Each layer and surface feature rests on one another, maintaining an isostatic balance. For instance, the core has an average density of 13.5 gm/cm³, the mantle's density ranges from 3.3 to 5.7 gm/cm³, and the continental crust's density is around 2.7 gm/cm³.
- The concept of isostasy is particularly useful for explaining the 'glacial adjustment' observed in Scandinavian countries following the Pleistocene great ice age. The raised beaches of Finland, for example, demonstrate that an uplift of about 250 meters has occurred over the last 8,000 years due to isostatic adjustment.


Development of the Isostasy Concept

The concept of isostasy was developed by geologists to explain the gravitational attraction of massive mountainous regions. The idea was first introduced when Pierre Bouguer, during his Andes expedition from 1735-45, discovered that the huge volcanic peak of Chimborazo did not attract the plumb line as expected. He concluded that the Andes' gravitational attraction was much smaller than anticipated based on the mass of the mountains.

Similarly, during the geodetic survey of the Indo-Gangetic plain in 1859, discrepancies were noted in determining latitudes under the supervision of Sir George Everest, then Surveyor General of India. The latitudinal difference between Kalianpur and Kaliana, which are 370 miles apart, was calculated using both direct triangulation and astronomical methods. Kaliana was just 96 km away from the Himalayas.

- The results of the two methods differed by 5.23 seconds:
- Results obtained through triangulation: 5° 23' 42.294"
- Results obtained through astronomical method: 5° 23' 37.058"
- Difference: 5.236"

This discrepancy was attributed to the decreased gravitational attraction of the Himalayas, which caused the plumb bob used in the astronomical latitude determination to be deflected. Various theories have been proposed to explain the gravitational attraction, deflection, and isostatic balance among different landforms.




Which geologist first proposed the term "isostasy"? **A.** Pierre Bouguer **B.** Sir George Everest **C.** Clarence Dutton **D.** Archdeacon Pratt

Concept of Sir George Airy

- The concept of Sir George Airy revolves around the idea that the Earth's crust is not hollow, but rather, the weight of the mountains is balanced by lighter materials below. According to Airy, the crust, made of lighter materials called "sial," floats on a denser substratum known as "sima." This means that massive mountain ranges like the Himalayas are floating on denser magma, similar to how a boat floats on water.
- Airy's concept is based on the principle of flotation, which states that the ratio of the part of an object floating above the water (freeboard) to the part submerged (draught) is 1 to 9. Applying this principle to the Earth's crust, it can

be assumed that for every 1 part of the crust above the substratum, there are 9 parts submerged. For example, considering the Himalayas' height of 8,848 meters, there must be a downward projection of lighter material beneath the mountain reaching a depth of around 80,000 meters.

- However, another scientist named Joly suggested a different freeboard-to-draught ratio of 1 to 8, which would result in a depth of around 70,784 meters for the Himalayas. Despite the difference in ratios, both theories agree that a significant amount of lighter material must be submerged beneath the mountains to maintain their balance.



Concept of Sir George


- According to Airy, the density of the Earth's crust remains the same regardless of the height of the landmass. This means that the continents are made of rocks with uniform density, but their thickness varies from place to place. To demonstrate this concept, Airy used iron pieces of different lengths, which submerged in mercury to varying depths depending on their lengths. This experiment can also be replicated using wooden pieces submerged in water.
- While Airy's concept is widely respected among the scientific community, it has some flaws. For instance, it is difficult to accept that the Himalayas have a downward projection of lighter material reaching depths of around 80,000 or 70,784 meters, as this would likely melt due to the high temperatures at such depths. Nevertheless, Airy's concept of isostasy remains an important contribution to the understanding of the Earth's crust and its underlying structure.

Concept of Archdeacon Pratt

- The concept of Archdeacon Pratt's isostasy was derived from his study of the difference in gravitational deflection during the geodetic survey of Kaliana and Kalianpur. He calculated the gravitational force of the Himalayas by taking the average density of the mountain range as 2.75 and found that the difference should have been 15.885 seconds. This led him to the conclusion that the


horizontal plane between two sites must be askew, affecting the vertical direction as well.

- Pratt further studied the rocks and their densities in the Himalayas and neighboring plains, discovering an inverse relationship between the height of the landforms and their density. In other words, the density of mountains is less than that of plateaus, which in turn is less than the density of plains and the ocean floor. According to Pratt, there is a level of compensation above which density varies among different land columns, but there is no change in density below this level. Within one column, density does not change, but it does vary between columns above the compensation level.



Line of Compensation


- Line of Compensation
- Pratt's central theme on isostasy can be summarized as "uniform depth with varying density." He believed that along the line of compensation, equal surface areas must underlie equal mass. For example, if there are two columns (A and B) with equal surface areas but different heights, the density of column A should be less than that of column B in order to achieve equal weight along the line of compensation. This concept suggests that larger columns have lower density, while smaller columns have higher density.
- While Pratt's concept of isostasy does not directly align with the law of floatation or the idea of root formation, there are indirect implications of these principles in his work. Bowie has noted that a close examination of Pratt's concept reveals glimpses of both floatation and root formation.
- In comparing the views of Airy and Pratt on isostasy, Bowie observed that "the fundamental difference between Airy's and Pratt's views is that the former postulated a uniform density with varying thickness and the latter a uniform depth with varying density."



Pratt's Concept

Concept of Hayford and Bowie

- The concepts of isostasy proposed by Hayford and Bowie are quite similar to those of Pratt. According to Hayford and Bowie, there is a plane of compensation where the crustal parts are entirely balanced. The densities of these crustal parts vary with the elevations of columns above this compensation plane.
- In this concept, the density of mountains is lower than that of the ocean floor, meaning that the crust is made of lighter material under mountains than under the ocean floor. Additionally, there is a zone beneath the compensation plane where density remains uniform in a lateral direction.
- Hayford and Bowie propose that there is an inverse relationship between the height of crustal columns and their respective densities (as assumed by Pratt) above the compensation plane. The compensation plane, or level of compensation, is thought to be located at a depth of about 100 km. Columns with rocks of lower density stand higher than columns with rocks of higher density.
- To illustrate this concept, consider four imaginary columns (interior plain, plateau, coastal plain, and offshore region) reaching the level of compensation. Although their heights differ, they are balanced by their varying densities. The assumption is that the differing volume of matter in each column is compensated by its density so that they all exert equal downward pressure at the level of compensation and balance each other.
- This statement may be understood with the help of fig below:



- This concept can be visualized using a diagram showing different columns of equal cross-sections made from various metals and ores with varying densities floating in a basin of mercury. All of these columns reach the same line (level of compensation) and exert equal weight along that line.

- Bowie compared the views of Airy and Pratt on isostasy and found considerable similarity between them. However, he also noted that they were not identical. Bowie recognized elements of Airy's concept of root formation and the law of flotation in Pratt's views, albeit indirectly.
- However, the idea that crustal parts (various reliefs) are in the form of vertical columns, as proposed by Hayford and Bowie, is not entirely accurate because crustal features are typically found in horizontal layers.

Concept of Jolly

- Jolly proposed his ideas on isostasy in 1925, challenging Hayford and Bowie's belief that there was a level of compensation at a depth of about 100 km beneath the Earth's surface. Jolly argued that the high temperatures at this depth would lead to complete liquefaction, making it impossible for such a level of compensation to exist.
- Additionally, Jolly disagreed with Hayford and Bowie's idea that the density above the level of compensation varied, while remaining uniform below it. He contended that this condition would be unstable, as geological events could easily disturb it and disrupt the level of compensation.
- Instead, Jolly suggested that there was a zone of compensation, rather than a single level. He believed that below a shell of uniform density, there was a 10-mile (16 km) thick layer where the density varied. This concept implied that the compensation process was not a linear phenomenon, but rather a zonal one.
- Jolly's ideas seemed to be more in line with the law of flotation, although he did not explicitly mention it. His concept was more similar to Airy's ideas than those of Hayford and Bowie. In Jolly's model, areas of low density within the 10-mile layer corresponded to the downward projections of the lighter continental crust, while areas of high density represented the spaces filled with heavier underlying material.

Pratt	Airy
All blocks have equal depth/thickness with varying density	All blocks have equal density with varying depth/thickness
Based on law of compensation	Based on law of flotation
He believed in contraction of earth in the process of origin of earth	He believed in upstanding position was result of expansion of earth interior after getting heated
His concept was based on local observation	His concept was also based on local observation
Pratt model is used for mid oceanic ridges	Airy model is used for continental topography especially mountain ridges
Pratt has not used word root	Height is proportional to root, higher the elevation thicker the block and root.

What is the central theme of Archdeacon Pratt's concept of isostasy? **A.** Uniform density with varying thickness **B.** Uniform depth with varying density **C.** Varying depth with uniform density **D.** Varying density with uniform thickness

Conclusion

The concept of isostasy has evolved through the contributions of various geologists, including Airy, Pratt, Hayford, Bowie, and Joly. These theories attempt to explain the balance between the Earth's large upstanding features and the underlying denser materials. Although their approaches differ, they all agree on the existence of a compensation mechanism that maintains this balance. Isostasy is an important concept in understanding the Earth's crust and its underlying structure, and has been instrumental in explaining phenomena such as glacial adjustment and gravitational deflection.

What is the main concept behind isostasy?

Isostasy refers to the state of balance that exists between the Earth's large upstanding features, such as mountain ranges and plateaus. It suggests that the Earth's lighter surface materials (sial) float on top of the denser magma (sima) found in the Earth's interior.

Who first proposed the concept of isostasy?

The concept of isostasy was first proposed by American geologist Clarence Dutton in 1889.

What are the key differences between the isostasy concepts proposed by Sir George Airy and Archdeacon Pratt?

Sir George Airy's concept of isostasy is based on the idea that the Earth's crust has a uniform density and varies in thickness, with lighter materials beneath mountain ranges balancing their weight. Archdeacon Pratt's concept, on the other hand, suggests that the crust maintains a uniform depth but varies in density, with larger columns having lower density and smaller columns having higher density.

What is the level or zone of compensation in the context of isostasy?

The level or zone of compensation refers to a specific depth or area within the Earth's crust where the crustal parts are entirely balanced. According to Hayford and Bowie, this level of compensation is located at a depth of about 100 km, while Joly suggests that it is a zone rather than a single level.

How does the concept of isostasy help explain the uplift of Scandinavian countries following the last ice age?

Isostasy can help explain the "glacial adjustment" observed in Scandinavian countries following the Pleistocene great ice age. The raised beaches of Finland, for example, demonstrate that an uplift of about 250 meters has occurred over the last 8,000 years due to isostatic adjustment.

1. What is the concept of isostasy?

Ans. Isostasy is a geological concept that refers to the equilibrium or balance between the Earth's lithosphere and asthenosphere. It explains the vertical movement of the Earth's crust in response to the redistribution of mass on the Earth's surface.

2. How does isostasy work?

Ans. Isostasy works based on the principle of gravitational equilibrium. When there is an increase in the load on the Earth's crust, such as the deposition of sediments or the melting of glaciers, the crust sinks to maintain balance. Conversely, when there is a decrease in the load, the crust rises to regain equilibrium.

3. What are the factors that influence isostatic adjustments?

Ans. Isostatic adjustments are influenced by several factors, including the thickness and density of the crust and underlying mantle, the presence of large bodies of water, the accumulation or erosion of sediment, and the tectonic activity in the region.

4. How does isostasy contribute to the formation of mountains and valleys?

Ans. Isostasy plays a significant role in the formation of mountains and valleys. When tectonic forces push up mountain ranges, the crust beneath them undergoes isostatic adjustment and sinks deeper into the mantle. Similarly, when erosion wears down mountains, the crust rebounds and uplifts, resulting in the formation of valleys.

5. What are the practical applications of isostasy in geology?

Ans. Isostasy has several practical applications in geology. It helps in understanding the regional variations in elevation, predicting the subsidence or uplift of coastal areas due to processes like sea level change or sediment deposition, and interpreting the isostatic rebound following the melting of glaciers. It is also used to analyze the isostatic response to tectonic events and to estimate the thickness of the Earth's crust in different regions.

Plate Tectonics

Tuzo Wilson from the **University of Toronto** was the first to use the term plate tectonics, while **W.J Morgan** from **Princeton University** published the **Plate Tectonics theory** for the first time in **1962**.

Plate Tectonics Theory


The Plate Tectonics Theory is a comprehensive framework that seeks to explain the dynamic processes and features of Earth's crust and the forces within it. This theory was developed in the 1960s, largely due to advancements in seafloor mapping. It is based on two main hypotheses: Arthur Holmes' convection current hypothesis and the concept of seafloor spreading, as proposed by Hess. The Plate Tectonics Theory is an improvement over Wegener's Continental Drift Theory and is regarded as the most sophisticated and comprehensive explanation for the movement of continents and the expansion of seafloors.

The main principles of the Plate Tectonics Theory can be summarized as follows:

1. Earth's interior can be classified based on mechanical rigidity into three main layers: the lithosphere, asthenosphere, and mesosphere. This classification rejects the earlier concepts of SIAL and SIMA.
2. The lithosphere is composed of Earth's crust and the upper part of the mantle. It is about 100 kilometers thick and has a lower density than the material beneath it, allowing it to "float."

3. The asthenosphere is a plastic, semi-fluid layer located below the lithosphere. The plates of the lithosphere essentially float on the asthenosphere, allowing for their movement.

In summary, the Plate Tectonics Theory provides a comprehensive explanation for the movement of Earth's crust and the forces that shape it, based on the interactions between the lithosphere and asthenosphere. This theory has significantly advanced our understanding of Earth's geological processes and the formation of its diverse features.



Interior of the Earth

According to Plate tectonics theory

The theory of plate tectonics suggests that the Earth's lithosphere, which includes the crust and upper mantle, is

fragmented into several pieces that float on a ductile layer known as the asthenosphere. These fragments, or tectonic plates, move due to convection currents generated in the upper mantle. The plates slide horizontally over the asthenosphere as rigid units.

- The thickness of the lithosphere varies, with oceanic regions ranging from 5-100 km thick and continental areas reaching approximately 200 km thick. Oceanic plates primarily consist of Simatic crust and are relatively thin, while continental plates are made up of Sialic material and are thicker in comparison.
- Tectonic plates can be classified into minor and major plates, as well as continental plates (e.g., Arabian plate) and oceanic plates (e.g., Pacific plate). Some plates are a combination of both continental and oceanic material, such as the Indo-Australian plate.
- The movement of these plates, driven by convection currents in the mantle, is responsible for the formation of various landforms and is the primary cause of all Earth movements. The boundaries between plates are areas of significant geological activity, including seafloor spreading, volcanic eruptions, crustal deformation, mountain building, and continental drift.
- The term "tectonics" originates from the Greek word "tektonikos," meaning building or construction. It refers to the deformation of the Earth's crust as a result of internal forces.

Which type of convergent boundary results in the formation of the Himalayas? **A.** Oceanic-continental convergence **B.** Oceanic-oceanic convergence **C.** Continental-continental convergence **D.** Transform fault boundary

Lithospheric Plates

A plate, in the context of Earth's structure, refers to a large section of the lithosphere, which includes the crust and the rigid upper mantle. These plates float on the more fluid-like asthenosphere below and move independently from one another. There are generally two types of plates: continental plates and oceanic plates. According to the geologist Xavier Le Pichon, the Earth's surface is made up of seven major plates and nine minor plates.

Major Tectonic Plates

The major tectonic plates of the Earth include:

- **Antarctica Plate:** This plate encompasses Antarctica and the surrounding oceanic plate. It is characterized by divergent boundaries, where the plate is moving away from its neighbors.
- **North American Plate:** This plate is shifting westwards at a velocity of 4-5 cm per year. It is a combination of both oceanic and continental crust.
- **South American Plate:** Similar to the North American Plate, the South American Plate is also moving westwards, but at a slightly slower rate of 3-4 cm per year. It is also half oceanic and half continental in composition.
- **Pacific Plate:** This plate is purely oceanic and is moving northwest at a rate of 2-3 cm per year.
- **India-Australia-New Zealand Plate:** This plate is a combination of the Indian Plate, the Australian Plate, and the New Zealand Plate, all of which are moving in different directions and at varying speeds.
- **Africa Plate:** This plate includes the continent of Africa as well as the eastern Atlantic Ocean floor. It is a combination of both oceanic and continental crust.
- **Eurasia Plate:** This plate is primarily continental and is shifting eastwards at a velocity of 2-3 cm per year. It includes the landmass of Europe and Asia as well as the adjacent oceanic plate.




Plate tectonics in the World

Minor Tectonic Plates

- **Arabian Plate:** This plate mainly consists of the Saudi Arabian landmass.
- **Bismarck Plate:** Divided into the North Bismarck Plate and South Bismarck Plate.
- **Caribbean Plate:** A tectonic plate encompassing the Caribbean Sea and surrounding regions.
- **Carolina Plate:** Located in the eastern hemisphere, this plate straddles the Equator north of New Guinea.

- **Cocos Plate:** A tectonic plate beneath the Pacific Ocean near Central America.
- **Juan de Fuca Plate:** Situated between the Pacific and North American plates, this plate is off the coast of the Pacific Northwest.
- **Nazca Plate:** A tectonic plate located in the eastern Pacific Ocean, off the west coast of South America.
- **Philippine Plate:** Positioned between the Asiatic and Pacific plates, this plate is beneath the Philippine Sea.
- **Persian Plate:** A tectonic plate that includes parts of Iran and surrounding regions.
- **Anatolian Plate:** Also known as the Turkish Plate, it comprises most of the Anatolia (Asia Minor) peninsula, including the country of Turkey.
- **China Plate:** A tectonic plate that covers the majority of mainland China and its surrounding regions.
- **Fiji Plate:** Situated between the Pacific Plate and the Indo-Australia Plate, this plate encompasses the Fiji island group.






Plate Boundaries Major and Minor



Persian Plate



Carolina Plate

Most of the plates include both continental and oceanic crusts. The area of the plates is fairly large in comparison to their depth and thickness. It has also been established that the depth of the plates is even less under the oceanic crust.

Three types of motion are possible between the plates:

1. Separation or divergent or constructive plate margins
2. Closing together or convergent or destructive plate margins
3. Transform or conservative plate margin

Type of Margin	Divergent	Convergent	Transform
Motion	Spreading	Subduction	Lateral sliding
Effect	Constructive (oceanic lithosphere created)	Destructive (oceanic lithosphere destroyed)	Conservative (lithosphere neither created or destroyed)
Topography	Ridge/Rift	Trench	No major effect
Volcanic activity?	Yes	Yes	No

Diagram (a) shows a divergent boundary with a ridge. Diagram (b) shows a convergent boundary with a trench and volcanoes. Diagram (c) shows a transform boundary with earthquakes within the crust. Labels include Ridge, Trench, Earthquakes, Lithosphere, and Asthenosphere.


Convergent And Divergent Plate Boundaries

Divergent Boundaries

The boundary between two plates that are moving apart or rifting. Rifting causes Seafloor Spreading.

Features of Divergent Boundaries –

- Mid-ocean ridges
- rift valleys
- fissure volcanoes



Different stages of development of Divergent Boundary

Continental Rift Valleys

Continental rift valleys can form when divergent boundaries occur within a continent. A prime example of this is the Great East African Rift Valley, which stretches from Ethiopia to Mozambique. Similarly, the Red Sea is a result of continental spreading, where the separation has been significant enough to create a "proto-ocean."



Great African Rift Valley

Convergent Boundaries

Convergent boundaries occur when tectonic plates collide, resulting in the destruction or compression of the Earth's crust. These boundaries are responsible for creating some of the most remarkable landforms on Earth, such as major mountain ranges, volcanoes, and oceanic trenches.

There are three types of convergent boundaries:

1. **Oceanic-continental convergence**
2. **Oceanic-oceanic convergence**
3. **Continental-continental convergence**

1. Oceanic-Continental Convergence: In this type of convergence, the denser oceanic lithosphere, which consists of basaltic crust, collides with the lighter continental lithosphere. As a result, the oceanic plate is forced beneath the continental plate in a process known as subduction. The subducting slab pulls the rest of the plate along with it, which is believed to be the primary driver of plate movement.

2. Oceanic-Oceanic Convergence: When two oceanic plates collide, subduction also occurs. As one oceanic plate is forced beneath the other, an oceanic trench forms, accompanied by shallow and deep-focus earthquakes and volcanic activity. Over time, a volcanic island arc develops, such as the Aleutian

Islands and the Mariana Islands. Eventually, these arcs may mature into more complex island arc systems, like Japan and the islands of Sumatra and Java in Indonesia.

3. Continental-Continental Convergence: In cases where two continental plates converge, subduction does not occur because the continental crust is too buoyant. Instead, massive mountain ranges are formed, such as the Alps. The most striking example of continental collision today is the formation of the Himalayas.

In summary, convergent boundaries are zones where tectonic plates collide, resulting in the formation of various landforms such as mountain ranges, volcanoes, and oceanic trenches. There are three types of convergent boundaries: oceanic-continental, oceanic-oceanic, and continental-continental, each with their unique characteristics and resulting geological formations.




Alps

Transform Fault Boundaries

- A transform fault, also known as a transform boundary or strike-slip boundary, is a type of fault that occurs along plate boundaries where the motion is primarily horizontal. Transform plate boundaries are areas where two tectonic plates slide past each other, resulting in deformation of the existing landforms rather than the creation or destruction of new landforms. The fracture zone that constitutes a transform plate boundary is referred to as a transform fault.
- In the ocean, these faults typically form perpendicular to mid-oceanic ridges, acting as planes of separation. On continents, well-known examples of transform faults include the North Anatolian Fault and the San Andreas Fault, which lies alarmingly close to Silicon Valley along the western coast of the United States. These faults demonstrate the characteristics of a transcurrent edge, where tectonic plates slide past one another horizontally.



San Andreas Fault California



North Anatolian Fault

Evidence supporting plate tectonics includes:

- **Deep ocean drilling:** The Glomar Challenger was a research vessel that drilled into the ocean floor, providing valuable information about the Earth's crust and plate tectonics.
- **JOIDES (Joint Oceanographic Institutions for Deep Earth Sampling):** This program collects and analyzes samples from the ocean floor to study the Earth's structure and history, further supporting the theory of plate tectonics.
- **Hot spots:** These are regions of intense volcanic activity that occur far from plate boundaries. Their existence suggests that the Earth's mantle is dynamic and that heat from the core drives the movement of tectonic plates.

- **Paleomagnetism:** This is the study of the Earth's magnetic field in the past. It provides evidence for the movement of continents over time and supports the theory of plate tectonics.
- **Magnetic reversal and seafloor spreading:** The pattern of magnetic reversals found in rocks on the ocean floor provides strong evidence for the process of seafloor spreading, which is a key component of plate tectonics.

The significance of plate tectonics is vast, as it influences various aspects of the Earth's geology:

- **Formation of major landforms:** Most major landforms, such as mountain ranges and ocean trenches, are formed as a result of plate tectonic processes.
- **Creation of new minerals:** Magmatic eruptions that occur during plate tectonic activity can bring new minerals to the Earth's surface.
- **Economic value:** Minerals like copper and uranium, which are valuable resources, are often found near plate boundaries.
- **Predicting future landmasses:** By studying the current movement of tectonic plates, scientists can predict how the Earth's landmasses will evolve in the future. For example, if current trends continue, North and South America may eventually separate, a piece of land may break off from the east coast of Africa, and Australia may move closer to Asia.

What is the primary driver of plate movement in the context of oceanic-continental convergence? **A.** Subduction of the oceanic plate beneath the continental plate **B.** Formation of rift valleys **C.** Sliding of plates along transform faults **D.** Mid-ocean ridges

Conclusion

The Plate Tectonics Theory is a groundbreaking framework that explains the dynamic processes and features of Earth's crust and the forces within it. Developed in the 1960s, this theory has advanced our understanding of Earth's geological processes and the formation of diverse features. By studying the interactions between the lithosphere and asthenosphere, as well as the movement of tectonic plates, scientists can explain the creation of major landforms, predict future landmasses, and uncover valuable resources. The evidence supporting plate tectonics, such as deep ocean drilling, hot spots, and

paleomagnetism, has significantly contributed to our understanding of Earth's complex geology.

What is the main difference between Wegener's Continental Drift Theory and the Plate Tectonics Theory?

Wegener's Continental Drift Theory proposed that continents were once connected and gradually drifted apart, but it lacked a mechanism to explain this movement. Plate Tectonics Theory, on the other hand, provides a comprehensive explanation of the movement of the Earth's crust by describing the interactions between the lithosphere and asthenosphere, including the process of seafloor spreading and the role of convection currents in the mantle.

What are the three types of plate boundaries and their associated geological features?

The three types of plate boundaries are divergent, convergent, and transform. Divergent boundaries are characterized by plates moving apart, leading to features such as mid-ocean ridges and rift valleys. Convergent boundaries involve plates colliding, resulting in the formation of mountain ranges, volcanoes, and oceanic trenches. Transform boundaries occur when plates slide past each other, causing deformation of landforms rather than creating or destroying them.

How do convection currents in the mantle contribute to plate tectonics?

Convection currents are generated in the mantle due to the heat from the Earth's core. These currents cause the semi-fluid asthenosphere to rise, spread out, and then sink back down, creating a circular motion. This movement of the asthenosphere drives the movement of the lithospheric plates that float on top of it, facilitating the processes of seafloor spreading, subduction, and plate movement.

What role do subduction and seafloor spreading play in the Plate Tectonics Theory?

Subduction is the process by which one tectonic plate is forced beneath another, typically occurring at convergent boundaries. This process recycles the crust material back into the mantle and drives the movement of the plates. Seafloor spreading, on the other hand, occurs at divergent boundaries where new crust is formed as molten material rises from the mantle and solidifies. This process results in the expansion of the seafloor and the movement of continents.

How does the study of paleomagnetism support the Plate Tectonics Theory?

Paleomagnetism involves the study of the Earth's magnetic field in the past, as recorded in rocks and minerals. This field of study has provided evidence for the movement of continents over time, showing that the Earth's magnetic poles have changed positions relative to the continents. Additionally, the pattern of magnetic reversals found in rocks on the ocean floor provides strong evidence for seafloor spreading, a key component of plate tectonics.

1. What is plate tectonics theory?



Ans. Plate tectonics theory is a scientific explanation for the movement of the Earth's lithospheric plates. It states that the Earth's outer shell is divided into several large and small plates that float on the semi-fluid asthenosphere beneath them. These plates interact with each other at their boundaries, leading to various geological phenomena such as earthquakes, volcanic activity, and the formation of mountain ranges.

2. What are lithospheric plates?



Ans. Lithospheric plates are large and rigid pieces of the Earth's outer shell called the lithosphere. The lithosphere is composed of the crust and uppermost part of the mantle. These plates float on the semi-fluid asthenosphere beneath them and are constantly moving and interacting with each other. There are several major plates, such as the Pacific Plate, North American Plate, Eurasian Plate, and many smaller plates.

3. What are convergent boundaries?

Ans. Convergent boundaries occur where two lithospheric plates collide or move towards each other. There are three types of convergent boundaries: oceanic-oceanic, oceanic-continental, and continental-continental. At oceanic-oceanic convergent boundaries, one oceanic plate is forced beneath the other, forming a subduction zone and often leading to the formation of volcanic arcs. At oceanic-continental convergent boundaries, the denser oceanic plate subducts beneath the less dense continental plate, causing volcanic activity and the formation of mountain ranges. At continental-continental convergent boundaries, both plates are of similar density and collide, resulting in the formation of large mountain ranges.

4. What are transform fault boundaries?

Ans. Transform fault boundaries occur where two lithospheric plates slide past each other horizontally. These boundaries are characterized by intense seismic activity and are responsible for creating strike-slip faults. The most famous transform fault boundary is the San Andreas Fault in California, where the Pacific Plate and North American Plate slide past each other. Transform boundaries do not lead to the creation or destruction of lithosphere but are significant sources of earthquakes.

5. How does plate tectonics theory explain geological phenomena?

Ans. Plate tectonics theory explains various geological phenomena by describing how lithospheric plates interact at their boundaries. For example, earthquakes occur when plates slide past each other at transform boundaries or when subduction occurs at convergent boundaries. Volcanic activity is often associated with convergent boundaries where subduction of oceanic plates leads to the formation of magma. The formation of mountain ranges is also explained by plate tectonics theory, as it involves the collision of continental plates at convergent boundaries. Overall, plate tectonics theory provides a comprehensive framework for understanding the dynamic nature of the Earth's surface and its geological processes.

Recent Views on Mountain Building

Introduction

Orogeny refers to the crucial geological process responsible for the formation and differentiation of the Earth's crust. In 1890, American geologist G.K. Gilbert coined the term 'orogeny' to describe the process of mountain building, specifically focusing on the formation of fold mountain belts like the Alps and the Rockies. As defined by the Oxford Dictionary of Geography, 'orogeny' refers to the tectonic movements of the Earth that involve the folding of sediments, faulting, and metamorphism.

- There are two primary types of orogeny: the cordilleran type, which involves the severe deformation of geosynclinal deposits, and the type formed by continental collision, in which oceanic crust and sediment are trapped between two masses of continental crust. The term 'orogeny' is not applicable to submerged mountains found in seas, as these underwater mountains are formed through processes that are significantly different from those responsible for creating mountain systems like the Alps and the Rockies.

- Two modern theories attempt to explain mountain building: the Geosynclinal theory and the Plate Tectonic theory. Both theories offer different perspectives on the geological processes involved in the formation of mountains and the Earth's crust.

Which of the following types of geosynclines is characterized by a high amount of volcanic products during their formation? **A.**

Monogeosynclines **B.** Polygeosynclines **C.** Eugeosynclines **D.** Miogeosynclines

Geosynclines

Geosynclines are significant geological features that have played a crucial role in the formation and development of the Earth's continents and ocean basins. These shallow, long, and narrow depressions of water are characterized by gradual sedimentation and subsidence. They are believed to have originated from mobile water zones surrounded by stable, rigid landmasses, which are the ancient nuclei of present-day continents.

- Geosynclines have not remained constant throughout geological history; their location, shape, dimensions, and extent have undergone significant changes due to various earth movements and geological processes. These depressions are typically bordered by two rigid masses called forelands, forming the edges of the geosynclinal regions.
- The concept of geosynclines was first introduced by James Hall and Dana, but it was further developed and expanded upon by Haug. J.A. Steers (1932) acknowledged their contributions, stating, "While the theory of geosynclines is due to Haug, the concept of the idea belongs to Hall and Dana." Geosynclinal theory gained widespread acceptance as it shed light on the unique features of folded mountains and the processes involved in their formation.

In summary, geosynclines are long, narrow, and shallow water depressions that have played a crucial role in the Earth's geological history. Their continuous sedimentation and subsidence have contributed to the formation of mountains and other landforms. The concept of geosynclines has evolved over time, thanks to the work of various geologists and geographers who have further developed and refined the theory.

Concept of Hall and Dana

The concept of Hall and Dana revolves around the study of folded mountains and their relationship with geosynclines. Dana examined the sediments found in the rocks of these mountains and theorized that they originated from marine sources. He proposed that these sediments were deposited in elongated, narrow, and shallow seas, which he referred to as geosynclines.

- Dana was the first to define geosynclines as long, narrow, and shallow sinking beds of seas. Hall further developed this concept, providing substantial evidence to demonstrate a connection between geosynclines and folded mountains. He argued that the rocks found in folded mountains were initially deposited in shallow seas, which eventually formed geosynclines.
- According to Hall, the continuous process of sedimentation causes the beds of geosynclines to subside. However, despite this subsidence, the depth of water in these geosynclines remains constant. It is important to note that geosynclines are characterized by their length, which is significantly greater than their width. This concept of Hall and Dana provides a foundation for understanding the geological processes that contribute to the formation of folded mountains and their relationship with geosynclines.





Fig. 11.2 : Sinking beds of geosynclines due to sedimentation and subsidence.

Concept of E. Haug

- The concept of geosynclines, credited to Hall and Dana, was further developed by E. Haug. He defined geosynclines as long, deep bodies of water that were much longer than they were wide. Haug created paleogeographical maps of the world, illustrating the long and narrow oceanic tracts that were later folded into mountain ranges.
- Haug proposed that the locations of present-day mountains were once occupied by geosynclines or oceanic tracts. These geosynclines existed as mobile

water zones between rigid masses. He identified five major rigid masses during the Mesozoic era: the North Atlantic Mass, the Sino-Siberian Mass, the Africa-Brazil Mass, the Australia-India-Madagascar Mass, and the Pacific Mass. He also located four geosynclines between these ancient rigid masses: the Rockies geosyncline, the Ural geosyncline, the Tethys geosyncline, and the Circum-Pacific geosyncline.

- According to Haug, systematic sedimentation occurred in the geosynclines. The littoral margins of the geosynclines were impacted by transgressional and regressive phases of the seas. The marginal areas of the geosynclines contained shallow water where larger sediments were deposited, while finer sediments were deposited in the central parts of the geosynclines. These sediments were compressed and folded into mountain ranges due to the compressive forces coming from the margins of the geosynclines.
- Haug also noted that not all geosynclines necessarily went through the complete cycle of sedimentation, subsidence, compression, and folding of sediments. In some cases, mountains were not formed from the geosynclines despite continuous sedimentation over long geological time periods.
- While Haug's contributions to the development of the geosyncline concept are commendable, his theory has some limitations and confusing aspects. For instance, his paleogeographic map of the Mesozoic era depicted an unrealistic extent of rigid masses (land areas) compared to the geosynclines (oceanic areas).



Concept of J.W. Evans

J.W. Evans posits that geosynclines are highly diverse in their forms and locations, which makes it challenging to provide a clear definition for their shape and position. Geosynclines are characterized by their beds undergoing gradual subsidence due to sedimentation, and their forms and shapes evolve in

response to changing environmental conditions. As a result, a geosyncline can be either narrow or wide and take on various shapes.

There are several possible scenarios for the placement of geosynclines:

- They can be situated between two land masses, such as the Tethys geosyncline that was located between Laurasia and Gondwanaland.
- They can be found in front of a mountain or plateau, like the long trench that formed after the Himalayas' emergence, which was later filled with sediments to create the Indo-Gangetic Plains.
- They can be positioned along the edges of continents.
- They can be present in front of a river mouth, among other locations.

Evans emphasizes that all geosynclines, despite their diverse forms, shapes, and locations, share the common features of sedimentation and subsidence. Over time, geosynclines that have experienced prolonged periods of sedimentation eventually become compressed and folded, giving rise to mountain ranges.

Views of Schuchert

Schuchert's classification of geosynclines focused on their characteristics, such as size, location, and evolutionary history. He divided them into three categories:

- **Monogeosynclines:** These are exceptionally long, narrow, and shallow water tracts, as described by Hall and Dana. Monogeosynclines are situated either within a continent or along its borders and are characterized by continuous subsidence due to gradual sedimentation and the resulting load. They are called "mono" because they undergo only one cycle of sedimentation and mountain building. The Appalachian geosyncline in the United States is considered the best example of monogeosynclines. It was present during the Pre-Cambrian period and bordered by the highland mass known as Appalachia in the east. The Appalachian geosyncline experienced folding from the Ordovician to Permian periods.
- **Polygeosynclines:** These are long and wide water bodies, broader than monogeosynclines. Polygeosynclines have relatively longer lifespans and more complex evolutionary histories. They are thought to have experienced more than one phase of orogenesis,

leading to the formation of one or more parallel geanticlines arising from their floors during the squeezing process. Polygeosynclines originated in positions similar to those of monogeosynclines. The Rocky and Ural geosynclines are representative examples of polygeosynclines.

- **Mesogeosynclines:** These are very long, narrow, and mobile ocean basins that are surrounded by continents on all sides. Mesogeosynclines are characterized by great abyssal depths and long, complex geological histories. They undergo several geosynclinal phases, such as sedimentation, subsidence, and folding. Mesogeosynclines are similar to the geosynclines described by Haug. The Tethys geosyncline is a typical example of this type, and the Mediterranean Sea is its remnant. The Tethys geosyncline was folded into the Alpine mountains of Europe and the Himalayas of Asia. The unfolded remaining portion of the Tethys geosyncline became the Mediterranean Sea, an example of Kober's median mass.

Concept of Arthur Holmes

A. Holmes, a renowned geologist, has made significant contributions to the understanding of geosynclines, including their main characteristics, causes of origin, and the processes involved in sedimentation, subsidence, and orogenesis. He believed that sedimentation leads to subsidence, but this alone cannot account for the greater thickness of sediments in geosynclines. Instead, earth movements can cause substantial subsidence in geosynclinal beds. Holmes also noted that the subsidence of geosynclinal beds occurs gradually, not suddenly.

For instance, the deposition of sediments up to 12,160 meters (40,000 feet) thick in the Appalachian geosyncline could have occurred over a 300-million-year span from the Cambrian period to the early Permian period, at a rate of one foot of sedimentation every 7,500 years.

Holmes identified four major types of geosynclines and described their origins separately:

- **Geosynclines formed by magma migration:** Holmes posited that the earth's crust consists of three layers, and the migration of magma from the intermediate layer to neighboring areas can cause the collapse and subsidence of the outer layer, forming a geosyncline. The Coral Sea is a present-day example of this process.
- **Geosynclines formed by metamorphism:** Holmes argued that converging convective currents cause compression-induced metamorphism in the lower layer of the crust, increasing the density of rocks and causing subsidence, which leads to the formation of a geosyncline. The Caribbean Sea, western Mediterranean Sea, and Banda Sea are examples of this type of geosyncline. However, this concept has been criticized because compression caused by convergent convective currents might not result in metamorphism, but rather in the melting of rocks due to high temperatures.
- **Geosynclines formed by compression:** Some geosynclines result from compression and subsequent subsidence of the outer crustal layer caused by convergent convective currents. The Persian Gulf and the Indo-Gangetic trough are considered typical examples of this type of geosyncline.
- **Geosynclines formed by thinning of the sialic layer:** Holmes suggested two possibilities when a column of rising convective currents diverges in opposite directions after reaching the lower crustal layer: (a) the sialic layer stretches apart due to tensile forces exerted by diverging convective currents, causing thinning and resulting in the creation of a geosyncline, such as the former Tethys geosyncline; or (b) the continental mass may separate due to enormous tensile forces generated by the divergent convective currents, as is thought to have occurred in the formation of the former Ural geosyncline.

View of Others

Dustar has categorized geosynclines into three types based on the structure of mountain ranges:

- Inter-continental geosynclines are located between two continental or land masses. This type is similar to Schuchert's mesogeosyncline classification. The Ural geosyncline is a representative example of this category.
- Circum-continental geosynclines are typically found along the edges of continents. This type corresponds to Schuchert's monogeosyncline classification.
- Circum-oceanic geosynclines are generally situated in the marginal areas where continents meet oceanic margins. Stille refers to these as marginal geosynclines, while others call them special or unique geosynclines. More extensive geosynclines are referred to as orthogeosynclines by Stille.

Stille has also classified geosynclines based on the presence of volcanic activity during their formation:

- Eugeosynclines have a relatively high amount of volcanic products, indicating a high level of igneous activity (the Greek prefix "eu" means high status).
- Miogeosynclines contain a lower amount of volcanic products, suggesting a lower level of igneous activity (the prefix "mio" means low).

In summary, Duxter's classification of geosynclines involves three types based on their location in relation to continents and oceans, while Stille's classification is based on the level of volcanic activity during the formation of geosynclines.

What are the three main stages of the geosynclinal history? **A.**


Lithogenesis, orogenesis, and gliptogenesis **B.** Sedimentation, compression, and denudation **C.** Formation, subsidence, and folding **D.** Erosion, deposition, and mountain building

Stages of Geosynclines

The geosynclinal history is divided into three main stages, including:

Lithogenesis (the stage of geosyncline formation, sedimentation, and subsidence of geosynclinal beds):


- This stage is characterized by the formation, sedimentation, and subsidence of geosynclines. These geosynclines form due to the contraction caused by the Earth's cooling process. The adjacent stable areas or kratogens, which surround the geosynclines, are subjected to the forces of erosion.
- As a result, rocks and boulders from the kratogens are continuously worn away and deposited onto the beds of the geosynclines. This leads to the subsidence of the geosynclines, as the weight of the deposited sediments causes them to sink. The combined processes of sediment deposition and the resulting subsidence contribute to the ongoing accumulation of sediments, which increases the overall thickness of the geosynclinal beds.



The stage of lithogenesis: creation of geosyncline followed by sedimentation and subsidence.

Orogenesis (the stage of squeezing and folding of geosynclinal sediments into mountain ranges:


- In this stage the geosynclinal sediments are squeezed and folded into mountain ranges. There is a convergence of forelands towards each other due to the force of the contraction of the earth. The enormous compressive forces produced by these moving forelands produce contraction, squeezing and folding of sediments deposited on the geosynclinal bed.
- The parallel mountain ranges found on both sides of the geosyncline have been termed by Kober as rand ketten meaning marginal ranges. Kober viewed the folding of geosynclinal sediments to be dependent upon the intensity of the compressive forces. Compressive forces of normal and moderate intensity produce marginal ranges on two sides of the geosyncline leaving the middle part unaffected.
- The unfolded middle part is termed as zwischengebirge (between mountains) or median mass. He viewed the Tethys geosyncline as bordered by the European foreland in the north and by the African foreland in the south.
- The sedimentary deposits of the Tethys geosyncline had undergone massive compression due to the converging movement of the European landmass (foreland) and the African foreland, leading to the formation of the Alpine mountain system.



The stage of orogenesis: squeezing and folding of geosynclinal sediments due to compressive forces; the whole of geosynclinal sediments are folded when the compressive forces coming from the sides of geosyncline is enormous and acute.

Gliptogenesis (the stage of gradual rise of mountains, and their denudation and consequent lowering of their heights):

- This phase of mountain- building is characterized by a gradual ascent of mountain ranges and the ongoing denudation processes by natural agents.



Folding of marginal sediments into marginal ranges and formation of median mass when the compressive forces are moderate.

Conclusion

Orogeny is a crucial geological process responsible for the formation and differentiation of the Earth's crust, specifically the creation of fold mountain belts. Geosynclines, long and narrow depressions of water, play a significant role in the Earth's geological history, including the formation of mountains and other landforms. Various geologists and geographers have contributed to our understanding of geosynclines, their relationship with mountain building, and their classifications. The study of geosynclines and orogeny helps us understand the complex processes that have shaped the Earth's crust and contributed to the formation of the diverse landscapes we see today.

What is orogeny, and what are the two primary types of orogeny?

Orogeny refers to the geological process responsible for the formation and differentiation of the Earth's crust, specifically focusing on the formation of fold mountain belts. The two primary types of orogeny are the cordilleran type, which involves the severe deformation of geosynclinal deposits, and the type formed by continental collision, in which oceanic crust and sediment are trapped between two masses of continental crust.

What are geosynclines, and how do they contribute to the formation of mountains?

Geosynclines are long, narrow, and shallow water depressions that have played a crucial role in the Earth's geological history. Their continuous sedimentation and subsidence have contributed to the formation of mountains and other landforms. The concept of geosynclines has evolved over time, thanks to the work of various geologists and geographers who have further developed and refined the theory.

What is the difference between monogeosynclines, polygeosynclines, and mesogeosynclines?

Monogeosynclines are long, narrow, and shallow water tracts that undergo only one cycle of sedimentation and mountain building. Polygeosynclines are long and wide water bodies that have relatively longer lifespans and more complex evolutionary histories, experiencing more than one phase of orogenesis. Mesogeosynclines are very long, narrow, and mobile ocean basins that are surrounded by continents on all sides, characterized by great abyssal depths and long, complex geological histories.

How do the concepts of Hall and Dana, E. Haug, J.W. Evans, and Arthur Holmes contribute to our understanding of geosynclines?

The concepts of Hall and Dana were the first to define geosynclines and establish a connection between them and folded mountains. E. Haug further developed the concept, identifying major geosynclines and their relationships with rigid landmasses. J.W. Evans emphasized the diversity of geosynclines in terms of their forms and locations. Arthur Holmes contributed by explaining the possible origins and causes of geosynclines, as well as the processes involved in sedimentation, subsidence, and orogenesis.

What are the three main stages of geosynclinal history?

The three main stages of geosynclinal history are lithogenesis (the stage of geosyncline formation, sedimentation, and subsidence of geosynclinal beds), orogeneis (the stage of squeezing and folding of geosynclinal sediments into mountain ranges), and gliptogenesis (the stage of the gradual rise of mountains and their denudation and consequent lowering of their heights).

5. How do recent views on mountain building differ from the views of Schuchert?

Ans. Recent views on mountain building differ from Schuchert's views in terms of the mechanism and driving forces behind mountain formation. While Schuchert attributed mountain building to the accumulation of sediments in geosynclines, recent views emphasize the role of plate tectonics and the collision of tectonic plates. Plate tectonics provide a more comprehensive understanding of mountain building processes, incorporating factors such as subduction, crustal movement, and the interaction of different lithospheric plates.

3. What are the stages of geosynclines?

Ans. Geosynclines go through several stages of development. The first stage is the initial subsidence, where the trough starts to form. This is followed by the accumulation of sediments, which leads to the second stage known as the sedimentation stage. The third stage is the tectonic or orogenic stage, where the sediments undergo deformation and folding, resulting in the formation of mountain ranges. The final stage is the uplift and erosion stage, where the mountains are gradually worn down by weathering and erosion processes.

4. What are the recent views on mountain building?

Ans. Recent views on mountain building emphasize the role of plate tectonics in the formation of mountains. It is now understood that the collision of tectonic plates, along with subduction and the movement of Earth's crust, plays a significant role in the uplift and formation of mountain ranges. This view is supported by geological evidence and the observation of ongoing mountain building processes in various parts of the world.

1. What are geosynclines?

Ans. Geosynclines are elongated, subsiding troughs in the Earth's crust where sedimentary deposits accumulate. These troughs are often associated with the formation of mountain ranges.

2. What are the views of Schuchert regarding geosynclines?

Ans. Charles Schuchert, an American paleontologist, proposed the concept of geosynclines in the early 20th century. He believed that geosynclines were the primary sites of sedimentation and the source of materials that eventually formed mountain ranges.

Volcanicity

A **volcano** is a vent or opening through which molten lava, ash, gases, and other materials from the Earth's interior are released. The majority of volcanoes are found at convergent and divergent plate boundaries, while some are located within plate interiors and connected to hot spots.


Types of Volcanic Eruptions

- **Icelandic eruptions** are characterized by fissures that allow basaltic lava to flow out in large quantities. This

type of eruption is generally non-violent and results in the formation of plateaus. Examples include the Columbia River Basalt Group in the United States and the Deccan Plateau in India.

- **Hawaiian eruptions** can involve fissures, craters, or calderas, and typically produce small lava domes from which lava and gases are released. These eruptions are characterized by fire fountains and relatively low amounts of gas emissions.

- **Strombolian eruptions** create stratovolcanoes, where the lava flows out and solidifies in layers. These eruptions involve rhythmic to continuous explosions caused by the intermittent release of gas. Additionally, they can eject clots of lava, which result in the formation of bombs and scoria.
- **Vulcanian eruptions** also form stratovolcanoes, but their associated lava is more viscous. The eruptions can be more violent due to the formation of a solid crust over the vent, leading to a buildup of gas pressure. These eruptions typically involve the ejection of bombs, pumice, and ash, followed by lava flows from the top or flank of the volcano.
- **Vesuvian eruptions** are extremely violent and involve the explosive release of gas-charged magma from a stratovolcano vent. These eruptions usually occur after a period of relative calm and involve the ejection of lava in an explosive spray, accompanied by gas clouds that can reach great heights and deposit tephra.
- **Plinian eruptions** are a more intense form of Vesuvian eruptions, characterized by the rapid upward movement of gas clouds that can extend for several miles. These clouds are typically narrow at the base and expand outward at higher elevations, but they generally contain low amounts of tephra.
- **Pelean eruptions** result from high-viscosity lava and delayed explosiveness. The conduit of a stratovolcano can become blocked by a dome or plug, with gas escaping through lateral openings or by the destruction or uplift of the plug.
- **Katmaian eruptions** are a variant of Pelean eruptions, characterized by the massive release of fluidized ash flows and widespread explosive tephra. These eruptions commonly produce ignimbrites and can also lead to the formation of hot springs and fumaroles.



© 2011 Encyclopædia Britannica, Inc.

Classification of Volcanoes

On the basis of material erupted:

1. **Basalt cone:** basalt cones are very rare. They are low rather than high because of the fluidity of basaltic lava.
 - Ex- Rangitoto (New Zealand) and Skjaldbreit (Iceland) are the most suitable examples.
2. **Basalt dome:** basalt domes are flat domes formed by fluid basalt lava. They are less in height and broad.
 - Hawaiian volcanoes are the best example.
3. **Ash and cinder cones:** these are high volcanoes formed by high viscous lava. Ash and cinder cones are built where eruptions are the explosive type with predominance of pyroclastic material. The growth of an ash or cinder cone begins around a crater.
4. **Composite or Strato cone:** they are formed by alternate sheets of lava and pyroclastic material. It is formed by alternate periods of explosive and quiet eruptions. Most of the largest volcanoes of the world fall under this category.
 - **Mt Fujiyama of Japan, Vesuvian of Italy, Cotapaxi, and Chimborazo of Ecuador** are good examples.

What are the two types of lava that can be produced during volcanic eruptions? **A.** Basaltic lava and Acidic lava **B.** Shield lava and Cinder lava **C.** Fluid lava and Explosive lava **D.** Hot lava and Cold lava

On the basis of periodicity

1. **Active volcanoes:** volcanoes that constantly eject lava, gases, ash, cinder, etc. are known as active volcanoes. There are about 600 active volcanoes in the world, most of them being around the Pacific "ring of fire".
 - The Stromboli volcano erupts so much fire that it has been termed as the lighthouse of the Mediterranean.
2. **Dormant volcano:** A volcano that has not erupted for a long time but still has the probability of eruption is called a dormant volcano. **Mt Kilimanjaro is one such volcano.**
3. **Extinct volcano:** a volcano that functioned in a long geological past but is no longer active is called an extinct volcano. They are only the remnants of the volcanoes. In

many of the cases crater, lakes are formed by filling of water into craters.


- **Aconcagua of Andes** is a typical example of an extinct volcano.

Volcanic Mountains

These mountains are made of volcanoes (made of material ejected from fissures in the earth's crust)

The material includes:

- Molten lava
- Volcanic bombs
- Cinders
- Ashes
- Dust and liquid mud



These materials fell around the vent in successive layers, building up a characteristic volcanic cone.

These mountains are often called as **mountains of accumulation**.

Example:

- Mt. Fuji (Japan)
- **Mt. Merapi (Sumatra)**
- Mt. Mayon (Philippines)
- Mt. Agung (Bali)
- Mt. Cotopaxi (Ecuador)

Plateaux

Elevated upland with extensive leveled surfaces and usually descend steeply to the surrounding lowland.

- Also known as **Tablelands**
- Most highlands are subject to the erosional process and so, plateaux are no exception.
- According to the mode of formation and physical appearance, the plateau is of 3 types – **Tectonic, Volcanic, and Dissected**.

Tectonic Plateaux

- As the name suggests they are formed because of Movements inside the earth which sometimes caused uplifting of leveled land.
- They have uniform altitude and considerable size.
- Example – Deccan plateau (India)
 - Meseta of central Iberia (tilted tectonic plateau)
 - Harz of Germany (faulted)

Intermontane Plateaux

- Plateaus enclosed by fold mountains (Intermontane means situated between mountains)
- Example – Tibet plateau between Himalayas and Kunlun
- The intermontane plateaus are some of the most extensive and highest plateaux in the world.

Volcanic Plateaux

- Similar to the volcanic mountains, the volcanic plateau formed from the solidification successive layers deposited by basaltic lava which comes from inside the earth.
- Also known as lava plateau
- Example-
 - Antrim plateau of Northern Ireland
 - The northwestern part of Deccan plateau
 - Columbia – Snake plateau

Dissected Plateaux

- As the name dissected suggests, they are as a result of weathering and erosion by running water, ice and winds
- Weathering and erosion and gradually worn down and made surfaces irregular
- In Humid highland, stream and sometimes glaciation cut deep narrow valleys in the plateau which are dissected plateaux
- **Example – Scottish Highlands**
- In drier countries, vertical corrosion by rivers and abrasion by winds will dissect the plateau into steep-sided tabular masses termed mesas and buttes, intersected by deep canyons. This is a common feature of arid and semi-arid areas, e.g. in the south-western USA
- Mineral-rich – Mot plateau are mineral-rich
- Example-

- **African plateau** – Gold, Diamond, copper, manganese, and chromium
- **Brazilian plateau** – Iron and Manganese
- **Deccan plateau** – Manganese, Coal, and Iron
- **Western Australia plateau** – Gold and Iron

Types of Plains

A plain can be –

- Level or undulation
- Rises up to 100 feet above sea level

These are the best land in a country and intensively cultivated and populated. Economic importance of a plain increases if a river also passes through it.

Example:

- Indo – Gangetic plains
- Mississippi plain
- Yangtze plain

Some most extensive temperate plains are grasslands like Russian steppes, North American Prairies and Argentina Pampas.

On the basis of their formation 3 types – **Structural, depositional and erosional** plains.

Structural Plains

- Structurally depressed areas of the world
- Most extensive natural lowlands
- Because they are formed from the horizontally bedded rocks they are relatively undisturbed by the crustal movements of the earth.
- Example –
 - Great plains of the Russian platform
 - Great plains of USA
 - Central lowlands of Australia

Depositional Plains

- The deposition name itself suggests that these plains are formed by the deposition of materials brought by various agents of transportation.
- Rise towards adjacent highlands
- Their fertility and economic development depend on types of sediments that are deposited by transportation.
- Active erosion (by the river) in the upper course results in the large quantities of alluvium brought down to the lower

course and deposited to form extensive alluvial plains, flood plains, and deltaic plains.

- These are most productive in agriculture.
- Example – Nile delta of Egypt (Rice and cotton) & Ganges delta (Rice and Jute).


Landforms Associated with Volcanic Activities

Volcanic activity affects Earth's landforms. Solid, liquid or gaseous materials may find their way to the surface from some deep-seated reservoir beneath. For example – Geyser and springs, volcanic eruption, sometimes gas sources are found. Molten magma is a mobile rock. It mostly enters weak portions of the crust.


The resultant landforms depend on many factors like –

- Strength and fluidity of the magma
- Types of cracks, faults, and joints penetrated by magma
- The manner in which magma escapes to the surface.


Intrusive landforms – Magma while thrusting its way up to the surface may cool and solidify within the crust as plutonic rocks.



Extrusive landforms – Magmas that reach the surface and solidify, form extrusive landforms.



Rocks formed by either plutonic or volcanic activity are called igneous rocks.




Landforms of Igneous Intrusions

Sill-

- Sill: Intrusion of molten magma horizontally along the bedding planes of sedimentary rocks.
- Denudation of overlying sedimentary strata in sill exposes the intrusion which looks like bold escarpment (a steep slope or cliff, such as one which marks the edge of a range of hills)
- Example – Great Whin sill of NE England

Dyke-

- Dyke – intrusion of molten magma vertically
- Denudation results – Upstanding walls or shallow trenches
- Example-
 - Cleveland Dyke of Yorkshire, England
 - Isles of Mull and Arran in Scotland
 - Quartzite dyke, North of Kuala Lumpur




Liths – igneous intrusions on a larger scale

Various types of liths – Laccolith, Lopolith, Phacolith, Batholith. All of them took place differently in the Earth's crust and solidified within the upper layers of the crust.


Laccolith

- Large blister or igneous mound with a dome-shaped upper surface and a level base fed by a pipe-like conduit from below.
- Shape similarity – Dome
- Example – Henry mountains, Utah, USA




Lopolith

- A large igneous intrusion which is lenticular in shape with a depressed central region.
- Shape similarity – Saucer
- Example – Bushveld lopolith of Transvaal, South Africa



Phacolith

- Igneous rocks occupying the crest of an anticline or the bottom of a syncline and fed by a conduit which is below it.
- Shape similarity – Lens
- Example – Corndon hill in Shropshire, England



Batholith

- A very large igneous intrusion extending to an unknown depth in the earth's crust.
- Shape similarity – irregular shape
- Example-
 - Wicklow Mountains of Ireland
 - Uplands of Brittany, France
 - Main Range of West Malaysia
- The orientation of Batholith –
 - It is believed that large masses of magma that were going up came into contact with country rock and metamorphosed them.
 - These metamorphosed rocks with solidified magma gave rise to extensive Batholiths.

Origin of Volcanoes

- We have read those volcanic activities are associated with crustal disturbances and mostly takes place where the crust has a weak portion because of faulting or folding.
- As we go deep inside crust temperature increases (1°F increase with 65 feet) and so we can expect the interior of the earth in the semi-molten state which could consist of solid, liquid, and gaseous material, collectively called as magma.
- Gases like – carbon dioxide, sulphurated hydrogen, and small proportions of Nitrogen chlorine, and other volatile substances charged magma heavily.
- Gases and vapor increase the mobility and explosiveness of Lava.

Types of lava

- Basaltic lava
- Acidic lava

All volcanoes pass through Active, Dormant, and Extinct stages but we are not sure about when they will be extinct. Example – Mt. Vesuvius and Mt. Krakatau. Both known as extinct but erupted violently.

Extrusive Landforms

- Nature and composition of Lava and ejected materials which reach the surface of the earth.
- As we have already read that basic lava is very fluid and it flows for long distances resulting in extensive lava plains.
- Example – our Deccan plateau has formed in a similar way & Snake basin, USA.
- Basalt plateau example – NW Deccan and also found in Iceland.

Volcanic Cones

- Highly fluid lava (i.e. Basic lava) forms lava domes or Shield volcanoes (with gently rising slopes and Broad, flattened tops)
- Example – Hawain volcano (best-developed lava domes)
- Mauna Loa
- Kilauea (steep-walled caldera with lava pit of Halemaumau)

Cinder Cone

- Less fluid lavas (i.e. acid lava) forms – Ash and cinder cones (with large central craters and steep slopes)

- They are small volcanoes (not greater than 1000 feet height)
- Example – Mt. Nuovo (near Naples) & Mt. Paricutin (Mexico)
- They form – Lava tongue, lava dammed lakes, lava bridges, lava tunnels

What comes out from the vent of the Volcano and what they do?

- Fine particles (Volcanic dust) are shot high in the sky. They can travel worldwide before coming to earth.
- They came down in the form of “Black Snow” and can bury houses or people.
- Pyroclasts – coarser fragmental rocks which include lapilli (rock fragments ejected from a volcano), scoria (basaltic lava ejected as fragments from a volcano, typically with a frothy texture), pumice and volcanic bombs.

Composite Cones

- Most volcanoes have this kind of cone.
- Built by – Several eruptions of lava, ashes and other volcanic materials (come out through conduit from the main conduit from magma chamber/reservoir) which forms a lot of layers and increases the height.
- From the main conduit subsidiary pipes or dykes reach the surface as a feeder to parasitic cones (like water pipelines in our house from a main big pipeline). From these subsidiary pipes, lava comes out to the sides of the cone.
- **Example –**
 - Mt. Etna (Sicily)
 - Mt. Stromboli (Lighthouse of Mediterranean)
 - Mt. Vesuvius
 - Mt. Fuji
 - Mt. Popocatepetl
 - Mt. Chimborazo

Caldera Cone

Crater formation – during an eruption the material from the top of the cone is blown off or collapses into vent widening the orifice into a large crater.

- Greatly enlarged depressions are called Calderas

- These are the result of violent eruptions accompanied by the subsidence of much of the volcano into the magma beneath.
- Water may collect in the crater or the caldera-forming crater or Caldera lakes e.g. Lake Toba in Sumatra.

Some Volcanic Eruptions

Mt. Vesuvius

- A somma-stratovolcano located on the Gulf of Naples in Campania, Italy
- Vesuvius has erupted many times since and is the only volcano on the European mainland to have erupted within the last hundred years.
- Today, it is regarded as one of the most dangerous volcanoes in the world because of the population of 3,000,000 people living nearby.
- Vesuvius was formed as a result of the collision of two tectonic plates, the African and the Eurasian. The former was subducted beneath the latter, deeper into the earth.

Mt. Krakatau

- A volcanic island situated in the Sunda Strait between the islands of Java and Sumatra in the Indonesian province of Lampung.
- The most notable eruptions of Krakatoa culminated in a series of massive explosions over August 26–27, 1883, which were among the most violent volcanic events in recorded history.
- The 1883 eruption ejected approximately 25 km³ (6 cubic miles) of rock (OMG)
- After remaining dormant for almost half a century, an eruption in 1927 pushed up a cinder cone from the submarine floor.
- This new volcanic island was named Anak Krakatau, meaning 'the child of Mt. Krakatau'.

Mt. Pelee

- A semi-active volcano at the northern end of Martinique, an island in the Lesser Antilles island arc of the Caribbean.
- Its volcanic cone is composed of layers of volcanic ash and hardened lava.
- The volcano is currently in a dormant state.
- Erupted white-hot lava and super-heated steam.

Kilauea-Hawaii

- Kilauea is Hawaii's most active volcano and parts of it have been continuously erupting since January 3, 1983. But the last major volcanic eruption at Kilauea was in 2014 and lasted for several months.

Mount Etna-Italy


- Europe's biggest active volcano, Mount Etna-erupted recently.

The Distribution of Volcanoes in The World

- Volcanoes are mostly located in intensely folded or faulted regions
- Active volcanoes – 500
- Dormant and extinct volcanoes – 1000s
- Occurrence – Coastal mountain ranges, off-shore islands and in the midst of oceans, but there are few in the interiors of continents.

Pacific Region

- Greatest concentration – circum pacific region (Pacific ring of fire) – Includes 2/3rd world's volcanoes.
- The western pacific – chain starts from the Aleutian Islands, Japan extending in the south up to the Philippines, Indonesia (Java and Sumatra particularly), pacific islands of Solomon, New Hebrides, Tease and North Island, New Zealand.
- Eastern pacific – Starting from Andes to Central America (particularly Guatemala, Costa Rica and Nicaragua), Mexico and right up to Alaska.



Atlantic

- Few volcanoes
- Many dormant or extinct
- Example – Madeira, Ascension, St. Helena, Cape Verde Island. and Canary Island
 - Active – Iceland and Azores

Mediterranean

- In alpine – folds – Vesuvius, Etna, Stromboli, and the Aegean Islands

Asia

- Mt. Ararat (recall with “are rat” or “Airavat” (Indra’s elephant name))
- Mt. Elbrus (elbows or eyebrows)
- It’s interesting and surprising that the Himalayas have no active volcanoes.

Africa

- Mt. Kilimanjaro and Mt. Kenya (east African rift valley – extinct)
- Mt. Cameroon – an only active volcano of west Africa.
- Madagascar – some volcano cones but not active volcanoes
- West Indian islands – Mt. Pelee and St. Vincent (violent explosion in recent times)
- Lesser Antilles – made of volcanic islands
- Interior volcanoes in continents – rare.

Most Recent Volcanic Eruptions

- Stromboli – Italy
- Dukono – Indonesia
- Ibu – Indonesia
- Sangeang Api – Indonesia
- Krakatau – Indonesia
- Merapi – Indonesia
- Kerinci – Indonesia
- Karangetang – Indonesia
- Barren Island – Andaman and Nicobar, India
- Sinabung – Indonesia
- Semeru volcano – Indonesia’s East Java province

Which type of plateau is formed due to the solidification of successive layers deposited by basaltic lava? **A.** Tectonic Plateau **B.** Intermontane Plateau **C.** Volcanic Plateau **D.** Dissected Plateau

Geysers and Hot Springs

Geysers

- Fountains of hot water and superheated steam.

- It can sprout up to the height of 150 feet from under the earth.

How do geysers work?


- Water near the volcanic or thermal region is heated beyond the boiling point.
- The jet of water emitted with the explosion and it is triggered or cause any gases which seep out from heated rocks

Where they are found?

- Mostly in 3 areas –
 - Iceland
 - New Zealand – Rotorua district of North Island
 - USA – Yellowstone Park (old faithful – world’s best-known geyser)

Hot Spring

- It can be found anywhere on the Earth where water is sunk deep enough beneath the surface to be heated by interior forces.
- Rises without any explosion (in geyser explosion is there)
- Contains – dissolved minerals
- Uses – dissolved minerals, heat house, swimming pools, and domestic purposes.
- Locations – Honduras, Canada, Chile, Hungary, Iceland, Israel, Japan, New Zealand, Fiji, and the United State



© 2011 Encyclopædia Britannica, Inc.

Semeru Volcano

- Semeru – also known as “The Great Mountain” – is the highest volcano in Java and one of the most active.
- It previously erupted in December 2019.
- Indonesia, with the maximum number of active volcanoes in the world, is prone to seismic upheaval due to its location on the Pacific's Ring of Fire.
- Semeru volcano is also part of the Island arcs formed by the subduction of the Indo-Australian plate below Sunda Plate (part of Eurasian Plate). The trench formed here is called the Sunda trench whose major section is the Java Trench.

Mt Merapi

- Merapi (Mountain of Fire) is the most active of Indonesia's 130 active volcanoes.
- It rises to 2,911 meters and has steep slopes with dense vegetation on its lower sides.
- It is located near the center of the island of Java and Indonesia's cultural capital, Yogyakarta.
- Mt Merapi's last significant eruption was in 2010. At that time, more than 300 people were killed and some 2,80,000 residents were forced to evacuate the surrounding areas.



Conclusion

Volcanic eruptions and their resulting landforms significantly impact the Earth's surface. Various types of eruptions, such as Icelandic, Hawaiian, Strombolian, Vulcanian, Vesuvian, Plinian, Pelean, and Katmaian, create diverse landforms like plateaus, volcanic mountains, and plains. The composition of lava and the manner in which it escapes the Earth's crust determines the formation of intrusive and extrusive landforms. Furthermore, volcanic activities are associated with crustal disturbances, which mostly take place in the weak portions of the crust due to faulting or folding. As such, understanding volcanic eruptions,

their types, and the landforms they create is crucial for comprehending the Earth's geological processes and landscape evolution.

What are the different types of volcanic eruptions?

There are several types of volcanic eruptions, including Icelandic, Hawaiian, Strombolian, Vulcanian, Vesuvian, Plinian, Pelean, and Katmaian eruptions. These eruptions differ in the level of violence, the type of lava produced, and the resulting landforms created.

How are volcanoes classified based on the material erupted and periodicity?

Volcanoes are classified into basalt cones, basalt domes, ash and cinder cones, and composite or strato cones based on the material erupted. They can also be classified as active, dormant, or extinct based on their periodicity.

What are some examples of volcanic mountains and plateaux?

Examples of volcanic mountains include Mt. Fuji (Japan), Mt. Merapi (Sumatra), Mt. Mayon (Philippines), Mt. Agung (Bali), and Mt. Cotopaxi (Ecuador). Examples of volcanic plateaux include the Antrim plateau of Northern Ireland, the northwestern part of the Deccan plateau, and the Columbia-Snake plateau.

What are the different types of plains and their formation?

Plains can be classified into structural, depositional, and erosional plains based on their formation. Structural plains are formed due to the Earth's crustal movements, while depositional plains are created by the deposition of materials brought by various agents of transportation. Erosional plains are formed as a result of weathering and erosion by running water, ice, and winds.

What are the different types of landforms associated with volcanic activities?

Volcanic activities can result in both intrusive and extrusive landforms. Intrusive landforms include sills, dykes, and various types of liths (laccoliths, lopoliths, phacoliths, and batholiths). Extrusive landforms include volcanic cones, lava domes, ash and cinder cones, and composite cones.

5. What are igneous intrusions and their associated landforms?

Ans. Igneous intrusions refer to the underground movement of magma, which solidifies and forms igneous rocks. These intrusions can create various landforms, such as batholiths, dikes, and sills. Batholiths are large bodies of intrusive igneous rocks that form deep beneath the Earth's surface. They are often exposed through erosion and form mountain ranges. Dikes are vertical or near-vertical sheets of magma that cut across existing rock layers. Sills are horizontal or near-horizontal intrusions that form between existing rock layers. They can create prominent landscape features, such as cliffs or steps.

1. What are the different types of volcanic eruptions?

Ans. There are several types of volcanic eruptions, including effusive eruptions, explosive eruptions, phreatomagmatic eruptions, and pyroclastic eruptions. Effusive eruptions occur when magma reaches the surface and flows out gently, forming lava flows. Explosive eruptions involve the sudden release of trapped gases in the magma, resulting in a violent explosion. Phreatomagmatic eruptions occur when magma interacts with water, causing steam-driven explosions. Pyroclastic eruptions involve the ejection of hot ash, rocks, and gas into the air, creating pyroclastic flows.

2. How are volcanoes classified?

Ans. Volcanoes are classified based on their shape, size, and eruption style. The two main types of volcanoes are shield volcanoes and stratovolcanoes. Shield volcanoes have broad, gently sloping sides and are formed by repeated eruptions of fluid lava flows. Stratovolcanoes, on the other hand, are tall and steep with layers of lava, ash, and volcanic rocks. They are formed by alternating eruptions of explosive and effusive lava.

3. What are volcanic mountains?

Ans. Volcanic mountains, also known as volcanic cones or volcanic peaks, are mountains that are primarily formed by volcanic activity. They are created when magma rises to the surface and erupts, depositing layers of lava, ash, and other volcanic materials. Over time, repeated eruptions build up the mountain, forming a distinct cone shape. Examples of volcanic mountains include Mount Vesuvius in Italy and Mount Fuji in Japan.

4. What are the landforms associated with volcanic activities?

Ans. Volcanic activities can create various landforms, including calderas, volcanic craters, lava domes, volcanic islands, and volcanic plateaus. Calderas are large, bowl-shaped depressions that form when the roof of a volcano's magma chamber collapses after a large eruption. Volcanic craters are circular depressions at the summit of a volcano, often formed during explosive eruptions. Lava domes are mounds of thick, viscous lava that accumulate around a volcanic vent. Volcanic islands are formed when volcanic eruptions occur underwater, creating new land masses. Volcanic plateaus are extensive, flat areas of land formed by repeated volcanic eruptions over long periods.

Earthquakes and Tsunamis

Earthquake

- An earthquake refers to the shaking or trembling of the Earth's surface, which occurs due to a sudden release of energy within the Earth's lithosphere. This energy takes the form of seismic waves, which travel through the surface layers of the Earth. Earthquakes can be caused by various factors, such as faulting, folding, plate movements, volcanic eruptions, and human-made factors like dams and reservoirs.
- Among all natural disasters, earthquakes are the most unpredictable and have the potential to cause significant destruction. Earthquakes can range in severity from minor tremors, which are caused by gentle waves of vibration within the Earth's crust and occur frequently, to major earthquakes that result from movement along faults. These major earthquakes can be particularly devastating, especially in densely populated areas.

Earthquake-Related Terminology

- **Earthquake intensity:** The measure of the severity of shaking produced by an earthquake at a given location.
- **Earthquake magnitude:** The measure of the energy released by an earthquake.
- **Richter Scale:** A scale developed by Charles F. Richter in 1935 to quantify the magnitude of an earthquake.
- **Mercalli Scale:** A seismic scale used to measure the intensity of an earthquake by observing its effects on people and structures.
- **Fault:** A fracture or zone of fractures in the Earth's crust where rocks on either side have moved relative to each other.
- **Focus (or hypocenter):** The point within the Earth where an earthquake begins.
- **Epicenter:** The point on the Earth's surface directly above the focus.

- **Seismic wave:** Waves of energy caused by the sudden breaking of rock within the Earth, which travel through the Earth and are recorded on seismographs.
- **Seismograph:** An instrument used to detect and record seismic waves.

Focus and Epicenter

The focus, or hypocenter, is the starting point of an earthquake within the Earth. The epicenter is the point on the Earth's surface directly above the focus. The intensity of the earthquake is highest at the epicenter and decreases as one moves away from it.

Richter Scale

The Richter magnitude scale measures the energy released by an earthquake. The scale ranges from 0 to 9, with each whole number increase representing a tenfold increase in shaking amplitude and a 31.6 times increase in energy released.

Mercalli Scale

The Mercalli intensity scale measures the effects of an earthquake on people, structures, and the Earth's surface. The scale ranges from 1 to 12, with higher numbers representing more severe effects.

Seismic Waves


Seismic waves are energy waves that result from the sudden breaking of rock within the Earth. There are two main types of waves: body waves and surface waves.

Body Waves

- **Primary waves (P-waves):** These are the fastest seismic waves, capable of moving through gases, solid rock, and fluids. P-waves cause rocks to move back and forth, similar to the movement of sound waves through the air.
- **Secondary waves (S-waves):** These waves are slower than P-waves and can only travel through solid rock. S-waves cause rocks to move up and down or side-to-side and arrive at the surface after P-waves.

Surface Waves

- **Love waves (L-waves):** Named after British mathematician A.E.H. Love, these waves are the fastest surface waves, moving the ground from side-to-side.
- **Rayleigh waves:** These waves cause the ground to move in an elliptical motion, with both vertical and horizontal components.



Rayleigh Waves

- The other kind of **surface wave is the Rayleigh wave**, named after Lord Rayleigh.
- A Rayleigh **wave rolls along the ground just like a wave rolls across a lake or an ocean**.
- Because it rolls, it moves the ground **up and down, and side-to-side** in the same direction that the wave is moving.
- **Most of the shaking felt from an earthquake is due to the Rayleigh wave, which can be much larger than the other waves.**

Classification of Earthquake

1. Classification of Earthquakes Based on Causative Factors:

Natural Earthquakes:

- **Volcanic:** Caused by volcanic activity
- **Tectonic:** Caused by movement of tectonic plates
- **Isostatic:** Caused by adjustments of the Earth's crust due to changes in weight distribution
- **Plutonic:** Caused by the movement of magma beneath the Earth's surface
- **Artificial Earthquakes:** Human-induced, such as those caused by mining or reservoir construction

2. Classification of Earthquakes Based on Depth of Focus:

- **Shallow focus (0-50 km):** Earthquakes with a focal depth of less than 50 km
- **Intermediate focus (50-250 km):** Earthquakes with a focal depth between 50 and 250 km
- **Deep focus (250-700 km):** Earthquakes with a focal depth greater than 250 km

3. Classification of Earthquakes Based on Human

Casualties:

- **Moderate:** Earthquakes resulting in fewer than 50,000 deaths
- **Highly hazardous:** Earthquakes resulting in 51,000 to 100,000 deaths
- **Most hazardous:** Earthquakes resulting in more than 100,000 deaths

Global Distribution of Earthquakes: Earthquake distribution around the world is closely related to the distribution of volcanoes.

The areas with the highest seismic activity are found in the Circum-Pacific regions, with the most frequent earthquakes occurring along the "Pacific Ring of Fire." Approximately 70% of all earthquakes take place in this belt.

Another 20% of earthquakes occur in the Mediterranean-Himalayan belt, which includes Asia Minor, the Himalayas, and parts of northwestern China.

The remaining 10% of earthquakes occur within the interiors of tectonic plates and at spreading ridge centers.

Earthquake Causes

Earthquakes are primarily caused by an imbalance or dis-equilibrium in the Earth's crust. This imbalance can be attributed to various natural and man-made factors.

1. Natural Causes:

- Volcanic eruptions
- Faulting and folding
- Upwarping and downwarping
- Expansion and contraction of gases within the Earth
- Movement of tectonic plates
- Landslides

2. Anthropogenic (Man-made) Causes:

- Deep underground mining
- Blasting of rocks with dynamite for construction purposes
- Deep underground tunnels
- Nuclear explosions
- Reservoir-Induced Seismicity (e.g., the 1967 Koyna Reservoir earthquake)
- Hydrostatic pressure from man-made water bodies like reservoirs and lakes


Plate tectonics offer the most logical explanation for the occurrence of earthquakes and volcanoes. Earthquakes usually occur along three types of plate boundaries: convergent, divergent, and transform.

- In India, earthquakes are more frequent in areas where the Indian Plate converges with the Eurasian Plate,

such as the Himalayan region. Although the peninsular part of India is generally considered a stable block, occasional earthquakes are felt along the margins of minor plates, like the 1967 Koyna earthquake and the 1993 Latur earthquake.

- Indian seismology experts have classified the country into four seismic zones: Zone II, Zone III, Zone IV, and Zone V. The highest and high-risk categories, Zone V and Zone IV, include the entire Himalayan region, the northeastern states, Western and Northern Punjab, Haryana, Uttar Pradesh, Delhi, and parts of Gujarat. The remaining parts of the northern plains and western coastal areas fall under the moderate-risk zone, while a large portion of the peninsular region is in the low-risk zone.

Seismic zones in India



Consequences of Earthquake

Impact of Earthquakes

Earthquakes can cause significant damage to human life and property due to the deformation of the ground surface resulting from the vertical and horizontal movement of the Earth's crust. For instance, the 2015 Nepal earthquake, which had a magnitude of 7.8 and a depth of 8.2 km, led to massive casualties and destruction due to poorly designed buildings and unscientifically constructed structures in urban areas. The city of Kathmandu experienced significant damage, with a death toll of around 8,000 people and an economic loss of 10 billion USD.

- **Landslides and Avalanches:** In mountainous regions, earthquake tremors can lead to slope instability and failure, resulting in landslides as debris falls down the slope. Additionally, large masses of ice may become dislodged from snow-covered peaks due to earthquakes, causing avalanches. For example, the 2015 Nepal

earthquake triggered several avalanches around Mount Everest, while the 2011 Sikkim earthquake led to landslides and significant damage to life and property, particularly the Singik and Upper Teesta hydropower projects.

- **Floods:** Earthquakes can cause severe disruption to dams and reservoirs, potentially leading to flash floods. Landslides and avalanches can also block river courses, contributing to flooding. The 1950 Assam earthquake, for instance, created a barrier in the Dihang River due to the accumulation of massive debris, causing flash floods in the upstream section.
- **Tsunamis:** Tsunamis are waves generated by the disruption of ocean basins and the displacement of large volumes of water. Seismic waves from an earthquake can displace the seafloor and create high sea waves, such as tsunamis. The 2004 Indian Ocean tsunami, which occurred on December 26th, was caused by an earthquake off the coast of Sumatra. The event was triggered by the subduction of the Indian plate under the Burmese plate, killing around 240,000 people in countries surrounding the Indian Ocean.
- **Fukushima Nuclear Accident:** The 2011 Tohoku earthquake in Japan, which had a magnitude of 9, resulted in tsunami waves reaching heights of 10 meters due to an undersea earthquake. These waves destroyed the emergency generators that cooled the reactors at the Fukushima Daiichi nuclear plant, leading to nuclear meltdown and radioactive fallout. The incident became a global concern as the radiation spread worldwide.


Earthquake Management

Earthquake management involves organizing and managing resources and responsibilities to address all humanitarian aspects of emergencies, with the goal of minimizing the negative effects of such hazards. Earthquake management encompasses everything from pre-earthquake risk reduction measures to post-earthquake recovery efforts.

- **Risk Recognition:** The first step in earthquake management is identifying areas that are more susceptible to earthquakes. This helps in prioritizing resources and planning for potential disasters.
- **Earthquake Monitoring and Early Warning Systems:** Accurately predicting the occurrence of an earthquake in a specific region remains a challenge for seismologists. However, the focus has shifted towards developing earthquake forecasting systems to mitigate

the impact of potential disasters. For example, Japan has implemented an earthquake early warning system that utilizes electronic signals, which travel faster than earthquake waves, to provide advance notice of an impending quake.

- **Structural Solutions:** Historical data reveals that over 95% of lives lost in earthquakes are due to the collapse of non-earthquake resistant buildings. Constructing earthquake-resistant buildings is more expensive compared to regular structures, which poses a challenge for countries with limited financial resources, such as India. A possible solution to this issue is seismic strengthening, which involves prioritizing structures for retrofitting and ensuring they meet safety standards. To implement this approach effectively, it is crucial to develop an earthquake hazard map that categorizes different zones based on their vulnerability to earthquakes.




Tsunami

A tsunami is a series of large ocean waves caused by seismic activity, such as earthquakes, that displace large volumes of water. The term "tsunami" comes from the Japanese words "tsu," meaning harbor, and "nami," meaning wave. These waves can cause massive destruction in coastal areas where underwater earthquakes occur.

- For a tsunami to occur, two conditions must be met: there must be an earthquake that transfers energy, and there must be a vertical displacement of water. This typically happens near oceanic trenches, where tectonic plates are being subducted. Tsunamis are not formed in areas where there is no sudden vertical movement, such as the Mid-Atlantic ridge. Seamounts breaking can also trigger tsunamis by causing a vertical displacement of water.
- When a tsunami is generated, its height-to-length ratio is very small, which allows it to pass unnoticed beneath ships at sea. As the wave approaches the shore, its height increases rapidly due to the rebound from the

shallow surface. The wave's period remains constant, but its velocity decreases and its height increases. In confined coastal waters, tsunamis can reach heights of over 30 meters.

- Tsunamis travel at speeds of 100-150 km/h in deep water, which can increase to 650-900 km/h. As they enter shallower waters near the coast, their velocity decreases and their height increases due to the "shoaling" effect. This causes a tsunami that was barely noticeable in deep water to grow several feet or more in height as it approaches the shore.
- The Pacific Ocean has the highest frequency of tsunamis, and since 1948, an International Tsunami Warning Network has been in place to alert coastal residents of potential dangers.
- Tsunamis can be caused by various factors, including earthquakes, landslides, undersea volcanoes, meteor and asteroid impacts, and human activities such as nuclear explosions.
- Since scientists cannot predict when earthquakes will occur, they cannot accurately predict when a tsunami will be generated. However, historical records and numerical models can help determine where tsunamis are most likely to occur. The most effective way to mitigate the impact of a tsunami is through an early warning system.
- A tsunami early warning system (TWS) detects tsunamis in advance and issues warnings to protect people's lives. It consists of two equally important components: a network of sensors that detect tsunamis and a communication infrastructure that issues timely alerts for evacuation of coastal areas. While it is not possible to predict the exact timing of a tsunami, early warning systems can help save lives and minimize damage by providing advanced notice to those in affected areas.



- Numerous early warning systems are in place around the world to alert citizens of potential natural disasters. These systems use various methods to communicate warnings, such as text messages, radio and television broadcasts, sirens, mosque loudspeakers, and police vehicles with loudspeakers.
- India joined the International Tsunami Warning System following the devastating tsunami in December 2004. Established in 2007, the Indian Tsunami Early Warning Centre (ITEWC) is part of the Indian National Centre for Ocean Information Sciences in Hyderabad, an autonomous organization under the Ministry of Earth Sciences. The ITEWC is equipped with the Deep Ocean Assessment and Reporting of Tsunamis (DART) system and is considered one of the best in the world.
- The ITEWC monitors seismic activity through a network of seventeen broadband seismic stations designed to detect tsunamigenic earthquakes and provide timely warnings to vulnerable populations. Additionally, the center collects data from global networks to identify earthquakes with a magnitude of at least 6.5. Since its inception in 2007, the ITEWC has monitored 339 such earthquakes.
- Furthermore, the ITEWC serves as one of the Regional Tsunami Advisory Service Providers (RTSP) for the Indian Ocean region, alongside Australia and Indonesia.

Conclusion

Earthquakes and tsunamis are natural disasters that occur due to the sudden release of energy within the Earth's lithosphere, leading to destructive seismic waves. Various natural and human-induced factors contribute to the occurrence of earthquakes, which can be categorized based on their causes,

depth, and the extent of human casualties. On the other hand, tsunamis result from events such as earthquakes, landslides, and undersea volcanic eruptions, among others. These disasters can cause significant damage to human life, property, and the environment. To minimize the impacts of earthquakes and tsunamis, it is crucial to establish early warning systems. Organizations like the International Tsunami Warning System and the Indian Tsunami Early Warning Centre play a vital role in detecting tsunamis and providing timely alerts. These systems allow for the evacuation of coastal areas, reducing the potential loss of life and property during such disasters.

What is an earthquake, and what causes it?

An earthquake is the shaking of the Earth's surface, resulting from the sudden release of energy in the Earth's lithosphere that creates seismic waves. It can be caused by factors such as faulting, folding, plate movement, volcanic eruptions, and anthropogenic factors like dams and reservoirs.

What is the difference between the Richter Scale and the Mercalli Scale?

The Richter Scale measures the magnitude of energy released by an earthquake, with values ranging from 0 to 9. The Mercalli Scale, on the other hand, measures the intensity of an earthquake and its effects, with values ranging from 1 to 12.

What are the different types of seismic waves?

Seismic waves can be classified into two main types: body waves and surface waves. Body waves include Primary waves (P-waves) and Secondary waves (S-waves). Surface waves include Love Waves (L-waves) and Rayleigh waves.

How can tsunami waves be generated?

Tsunamis can be generated by various causes such as earthquakes, landslides, undersea volcanoes, and meteor or asteroid impacts. The displacement of a large volume of water in the ocean due to these events can lead to the formation of tsunami waves.

What is a tsunami early warning system, and how does it work?

A tsunami early warning system (TWS) is a network of sensors and communication infrastructure designed to detect tsunamis and issue timely alarms for the evacuation of coastal areas. It consists of two equally important components: a network of sensors to detect

tsunamis and a communications infrastructure to issue timely alarms to permit evacuation of the coastal areas.

Which of the following factors contribute to the formation of a tsunami? **A.** Earthquake **B.** Landslides **C.** Undersea Volcanoes **D.** All of the above

What is the main purpose of a tsunami early warning system? **A.** To predict when earthquakes will occur **B.** To issue timely alarms for evacuation of coastal areas **C.** To prevent tsunamis from forming **D.** To measure the height of tsunamis

1. What is the classification of earthquakes?



Ans. Earthquakes can be classified into different categories based on their magnitude. The most commonly used classification system is the Richter scale, which measures the energy released by an earthquake. Other classification systems include the moment magnitude scale (Mw) and the seismic intensity scale (MMI).

2. What are the consequences of an earthquake?



Ans. Earthquakes can have various consequences, including: - Structural damage to buildings and infrastructure - Loss of human lives and injuries - Displacement of people from their homes - Disruption of essential services such as electricity, water, and communication - Triggering of landslides, avalanches, and tsunamis - Economic losses due to destruction of property and interruption of business activities

3. How is earthquake management carried out?



Ans. Earthquake management involves various measures to mitigate the impact of earthquakes and ensure the safety of affected populations. These measures include: - Developing and implementing building codes and regulations that ensure structural resilience - Conducting seismic hazard assessments to identify high-risk areas - Establishing early warning systems to provide advance notice of an impending earthquake - Educating communities about earthquake preparedness and response measures - Conducting drills and exercises to test emergency response plans - Improving infrastructure resilience and retrofitting vulnerable buildings - Conducting research and monitoring seismic activity to better understand earthquakes

4. What is a tsunami?



Ans. A tsunami is a series of ocean waves that are triggered by large-scale disturbances, such as earthquakes, volcanic eruptions, or underwater landslides. These waves can travel across the ocean at high speeds and cause significant damage when they reach coastal areas. Tsunamis can result in the loss of lives, destruction of infrastructure, and long-term environmental impacts.

5. How do earthquakes and tsunamis affect the environment?



Ans. Earthquakes and tsunamis can have various environmental impacts, including: - Damage to natural habitats and ecosystems - Contamination of water sources due to the disruption of underground aquifers or the release of pollutants - Displacement or extinction of wildlife species - Changes in coastal landforms and erosion - Induced landslides and soil liquefaction - Potential release of hazardous materials from damaged infrastructure It is important to note that the specific environmental consequences can vary depending on the magnitude and location of the earthquake or tsunami.

Concepts of Geomorphic Cycles and Landscape Development

Concepts of Geomorphic Cycles and Landscape Development

The geomorphic cycle, also known as the geographic cycle or cycle of erosion, is a theory that explains the evolution of landforms over time. This concept was first proposed by American geographer William Morris Davis in 1899, who developed a model that showed the sequential changes in landforms through various stages of development, similar to the concept of biological evolution.

- According to Davis, the initial stage of landform development, known as the youthful stage, begins with an uplift that produces fold or block mountains. As streams dissect the area, it reaches maturity and is eventually reduced to a near-sea-level surface called a peneplain during the old-age stage. The cycle can be interrupted by uplift at any point, causing the landform to return to the youthful stage, a process known as rejuvenation.
- The geomorphic cycle can be applied to various landforms, including hills, valleys, mountains, and river drainage systems, and follows a predetermined framework based on the stage of development. The earth's surface is affected by two types of forces: endogenetic and exogenetic. Endogenetic forces create vertical irregularities on the earth's surface, forming various relief features, while exogenetic processes originating from the atmosphere (such as rivers, wind, glaciers, and sea waves) work to remove these irregularities, ultimately reducing the relief to a low, featureless plain called a peneplain.

- The entire period of relief feature creation by endogenetic processes and their destruction by exogenetic processes is called the cycle of erosion. Although initially proposed for humid temperate conditions, the Davisian model has been extended to other landscapes, such as arid, glacial, coastal, karst, and periglacial environments.
- Some modifications have been suggested to Davis's model, such as C.H. Crickmay's proposal of a more powerful process called panplanation, resulting in a panplain rather than a peneplain. L.C. King proposed a new cycle of erosion called the cycle of pediplanation, specifically for arid and savanna regions of Africa.
- In more recent years, some researchers have rejected the evolutionary concept of landform development and advanced alternative theories, such as the dynamic equilibrium model of landform development and the tectonogeomorphic model. Despite these alternative theories, many modern models of landform development can be seen as modified forms of Davis's original concept.

Davis Model of Cycle of Erosion

The Davis Model of Cycle of Erosion, also known as the "geographical cycle," was introduced by Davis in 1899 as a way to classify and describe landforms. The model suggests that landforms undergo an evolutionary sequence due to the effects of denudation (weathering and erosion) processes over time. This leads to an ever-changing landscape that moves from an initial to an ultimate form, with the height and slope of hills declining over time.


- The model is based on three main factors: structure, process, and stage. Structure refers to the composition and arrangement of underlying rocks, including their hardness, porosity, and the presence of folds and faults. Process encompasses the various weathering and erosion agents, such as rivers, wind, and glaciers, as well as mass movements. Stage represents the time duration during which these processes act on the landform.
- Davis used the life stages of living beings - youth, maturity, and old age - as an analogy for the evolutionary stages of landforms. He argued that landforms undergo sequential changes throughout their life cycle as they progress through these stages. However, the duration of each stage is not fixed and depends on various factors, such as the resistance of rocks to erosion.

What are the main stages of the Davis Model of Cycle of Erosion?

- A. Youthful, Intermediate, Old
- B. Youthful, Mature, Old
- C. Initial, Mature, Final
- D. Early, Middle, Late

The Davisian model of the geographical cycle is based on several key assumptions:

- Landforms are shaped by the interaction of internal (endogenetic) forces from within the Earth and external (exogenetic) forces from the atmosphere, such as weathering, erosion, and various erosional agents like rivers, wind, groundwater, sea waves, glaciers, and periglacial processes.
- Landforms evolve in an orderly manner, developing a systematic sequence of forms over time in response to environmental changes.
- Streams erode their valleys quickly until a stable, graded condition is achieved.
- The land mass experiences a rapid rate of uplift, though slower rates can also be considered in the model.
- Erosion does not begin until the uplift is complete, meaning that uplift and erosion do not occur simultaneously. This assumption has been a point of contention among critics of the cyclic concept.



Stages of the cycle

The cycle of erosion begins with the upliftment of a landmass, where there is a rapid rate of short-period upliftment of a homogeneous structure. This phase is not included in the cyclic time as it is the preparatory stage of the cycle of erosion.

The model of the geographical cycle consists of an upper curve (UC) and a lower curve (LC), representing the hilltops or crests of water divides and valley floors, respectively. The horizontal line denotes time, while the vertical axis shows altitude from sea level. The cycle passes through three main stages: youthful, mature, and old.

- **Youthful stage:** Erosion begins after the completion of the landmass upliftment. At this stage, the topsurfaces or summits of the water divides are not significantly affected by erosion as the rivers are small and widely spaced. The rivers and their tributaries experience headward erosion, extending their lengths. Due to the steep slope and channel gradient, rivers actively deepen their valleys through vertical erosion, which results in deep and narrow valleys with steep valley-side slopes. The youthful stage is characterized by a rapid rate of vertical erosion and valley deepening due to the steep channel gradient, increased velocity and kinetic energy of river flow, and increased transporting capacity of the rivers.

- **Mature stage:** As the cycle progresses to the early mature stage, lateral erosion becomes more significant, and the drainage network becomes well integrated. Vertical erosion decreases, and the summits of water divides start to erode, resulting in a decrease in absolute and relative relief. Lateral erosion leads to valley widening, transforming V-shaped valleys of the youthful stage into wide valleys with uniform or rectilinear valley sides.

- **Old stage:** During the old stage, valley incision is almost absent, but lateral erosion and valley widening continue. Water divides are reduced in dimension through both down-wasting and back-wasting, causing a rapid decrease in absolute height. Relative relief also decreases due to active lateral erosion and the absence of vertical erosion. Valleys become almost flat with concave valley-side slopes, transforming the landscape into a peneplain. This stage lasts much longer than the previous two stages combined.

Analysis of Davis cycle of erosion

- The Davis cycle of erosion, specifically the concept of peneplain introduced by Davis, has been met with varied opinions from different geomorphologists. Some consider the peneplain to be a theoretical landform because, for the Davis cycle to fully occur, the landform must remain stable for an extended period. This stability is unlikely in reality, where both endogenetic and exogenetic forces are constantly at work, potentially disrupting the smooth progression of the cycle.
- Additionally, critics argue that the concept of peneplain contradicts the principle of isostatic adjustment. As a river erodes its valley, the removal of the overlying load is compensated by the addition of more material to the landform's base. This process pushes the landform upward, continuing the process of upliftment indefinitely. Therefore, the attainment of the peneplain stage is questioned due to its conflict with the idea of isostatic adjustment.
- Many geomorphologists also believe that the sequential change of landforms proposed by Davis is an oversimplification of the actual process of landform evolution. In reality, the development of landforms is a much more intricate and complex process.
- Furthermore, Walthar Penck criticized the Davis Model for its overemphasis on time. While the Davisian model presents a time-dependent series of landform development, Penck argued for a time-independent series. Penck believed that landforms do not undergo sequential changes over time and suggested removing 'time' from Davis' trio of 'structure, process, and time.'
- Several other researchers, including A.N. Strahler, J.T. Hack, and R.J. Chorley, have also rejected the Davisian concept of 'historical evolution' of landforms and instead proposed the dynamic equilibrium theory. However, the non-cyclic concept of 'dynamic equilibrium' and other open system, non-cyclic models have not generated much enthusiasm among modern geomorphologists.
- Despite these criticisms, the Davis cycle continues to be relevant in the study of slope evolution due to its widespread appeal and clear presentation.

Which of the following assumptions is NOT part of the Davisian model of the geographical cycle?

- A. Landforms are shaped by the interaction of internal and external forces.
- B. Landforms evolve in an orderly manner, developing a systematic sequence of forms over time in response to environmental changes.
- C. Streams erode their valleys quickly until a stable, graded condition is achieved.
- D. Erosion begins as soon as the land begins to rise.

Criticism of Davis Model

The Davis Model has faced significant criticism for its concept of upliftment. Davis described a quick upliftment process of short duration; however, evidence from plate tectonics shows that upliftment is actually an extremely slow and continuous process.

- Additionally, the relationship between upliftment and erosion in the Davis Model is considered flawed. Davis claimed that erosion cannot begin until upliftment is complete, which contradicts the natural process where erosion starts as soon as the land begins to rise. He defended this by stating that he excluded erosion from the upliftment phase to simplify the model and because erosion is insignificant during upliftment.
- Moreover, the Davisian model requires a long period of crustal stability to complete the erosion cycle, but plate tectonics suggests that such a stable period is unlikely due to constant plate movements and frequent tectonic events. Davis acknowledged this objection, arguing that if the desired crustal stability is not possible, the erosion cycle would be interrupted and a new cycle might begin.


Normal Cycle of Erosion

The normal cycle of erosion refers to the process of erosion driven primarily by running water, which is the most common type of erosion worldwide. It is particularly significant in glacial and arid regions. While W.M. Davis believed that humid temperate areas were the most typical case for fluvial cycles of erosion, this idea is still debated.

- In the normal cycle of erosion, a landmass is uplifted relative to sea level, leading to the formation of rivers and the beginning of erosional processes. Eventually, the uplift of the landmass ceases, allowing erosion to become more active. During this period of crustal

stability, the landmass neither rises nor sinks, and river valleys develop in a sequence of stages: youthful, mature, and old. Ultimately, these processes result in a low, featureless plain called a peneplain.

- The youthful stage begins with the uplift of the landmass. At this point, streams are short and few in number, and the primary consequent streams are also limited. Gullies and rills dominate the slopes, and through headward erosion, they lengthen their profiles. The development of tributaries from the primary consequent streams leads to a dendritic drainage pattern. Rivers experience rapid downcutting in their valleys due to the high channel gradient, resulting in narrow, steep valleys with convex side slopes, forming V-shaped juvenile valleys.
- During the mature stage, valley deepening decreases compared to the youthful stage, leading to a reduction in channel gradient and flow velocity. This stage is marked by early maturity, which is characterized by decreased transporting capacity, channel gradient, and river velocity. Lateral erosion becomes more active, causing valleys to widen. As rivers lose some of their transporting capacity, they deposit large boulders and sediments at the foothills, forming alluvial fans and cones. These deposits eventually create piedmont plains as they expand. With the reduced channel gradient, rivers meander and form loops.



- In the old stage, the channel gradient decreases further, and valley deepening ceases altogether. The number of tributary streams also declines, and valleys become wide and flat, with concave slopes. Rivers have a lower transporting capacity, leading to increased sedimentation and weathering. The rivers carve their courses extensively, forming large deltas at their mouths. The landscape ultimately transforms into a vast, flat plain known as a peneplain.

Conclusion

The geomorphic cycle, first proposed by William Morris Davis, is a theory that explains the evolution of landforms over time through various stages of development, including youthful, mature, and old stages. The cycle is driven by endogenetic and exogenetic forces and can be applied to various landforms such as hills, valleys, mountains, and river drainage systems.

Despite criticisms and alternative theories, the Davisian model continues to be relevant in the study of landform evolution and has inspired further research and modifications.

What is the geomorphic cycle?

The geomorphic cycle, also known as the geographic cycle or cycle of erosion, is a theory that explains the evolution of landforms over time. It was first proposed by American geographer William Morris Davis in 1899 and shows the sequential changes in landforms through various stages of development, similar to the concept of biological evolution.

What are the stages in the Davis Model of Cycle of Erosion?

The Davis Model of Cycle of Erosion describes three main stages of landform development: youthful, mature, and old. Each stage is characterized by specific features, such as deep and narrow valleys in the youthful stage, wide valleys with uniform valley sides in the mature stage, and flat valleys with concave slopes in the old stage.

What is a peneplain?

A peneplain is a low, nearly flat, featureless plain that is formed during the old stage of the geomorphic cycle. It occurs when the erosional processes have reduced the relief of the landform to a near-sea-level surface.

What are the main criticisms of the Davis Model of Cycle of Erosion?

The Davis Model has been criticized for its concept of upliftment, where it describes a rapid and short-duration upliftment process, while evidence from plate tectonics shows that upliftment is actually slow and continuous. The model also faces criticism for its overemphasis on time, stating that landforms undergo sequential changes over time, while some researchers argue for a time-independent series.

What is the normal cycle of erosion?

The normal cycle of erosion refers to the process of erosion driven primarily by running water, which is the most common type of erosion worldwide. It is particularly significant in glacial and arid regions. In this cycle, a landmass is uplifted, and river valleys develop through a sequence of stages: youthful, mature, and old.

These processes ultimately result in a low, featureless plain called a peneplain.

1. What are geomorphic cycles and how do they contribute to landscape development?

Ans. Geomorphic cycles refer to the repetitive sequence of processes that shape the Earth's surface over time. These cycles include weathering, erosion, transportation, and deposition of materials. They play a crucial role in landscape development by gradually wearing down mountains, carving out valleys, and reshaping the landforms we see today.

2. Could you explain the Davis Model of the Cycle of Erosion?

Ans. The Davis Model of the Cycle of Erosion, proposed by geographer William Morris Davis, suggests that landscapes go through a series of stages in a cyclic pattern. It consists of three main stages: youth, maturity, and old age. In the youth stage, the landscape is characterized by steep slopes, deep valleys, and fast erosion. In the maturity stage, the landscape becomes more gentle, with wider valleys and less erosion. Finally, in the old age stage, the landscape is flat and featureless, with little erosion occurring.

3. What does the analysis of Davis' cycle of erosion reveal about landscape development?

Ans. The analysis of Davis' cycle of erosion reveals that landscapes are not static, but rather undergo continuous change and evolution. It highlights the dynamic nature of landforms and the processes that shape them. By studying the different stages of the cycle, geographers can understand how landscapes develop over time and the factors that influence their formation.

4. What is a normal cycle of erosion?

Ans. A normal cycle of erosion refers to the idealized sequence of landscape development proposed by Davis. It assumes that landscapes evolve in a predictable pattern, starting from a youthful stage with high erosion rates, progressing to a mature stage with moderate erosion, and eventually reaching an old age stage with minimal erosion. However, it is important to note that real landscapes may not always follow this idealized pattern due to various factors such as tectonic activity, climate change, and human interventions.


5. Are there any frequently asked questions related to the concepts of geomorphic cycles and landscape development?

Ans. Yes, some frequently asked questions related to the concepts of geomorphic cycles and landscape development include: - How long does it take for a landscape to go through a complete cycle of erosion? - What are the main factors that influence the rate of erosion in a landscape? - Can human activities accelerate or disrupt the natural cycles of landscape development? - Are there any exceptions or variations to the normal cycle of erosion proposed by Davis? - How do geomorphic cycles contribute to the formation of specific landforms, such as mountains, canyons, or river valleys?

Denudation Chronology

Denudation chronology is the study of the historical development of landscapes through the processes of denudation, which involve the wearing away of Earth's surface by elements such as water, ice, wind, and waves. This approach aims to reconstruct the denudational history of a region by analyzing present-day landscapes, sediments, and relic landforms. It is based on concepts such as sequential landform change, uniformitarianism, cyclic nature of Earth's history, palimpsest topography, and Davisian model of cyclic evolution of landforms.


Which of the following concepts is crucial to the study of denudation chronology? **A.** Palimpsest topography **B.** Isostatic adjustment **C.** Soil formation **D.** Stream piracy



- The primary goal of denudation chronology is to identify, date, and interpret planation surfaces developed in past cycles and sub-cycles of erosion based on evidence from drainage development, river capture, relic surfaces, and past tectonic events. This approach emphasizes the historical development of landscapes and is based on the concept of palimpsest topography, which refers to surfaces that bear imprints of geomorphological processes from past geological periods.
- However, denudation chronology has some limitations, including the difficulty in explaining only small parts of existing landscapes due to the partial destruction of former surfaces by subsequent erosion. Additionally, this approach is highly speculative and deductive, as it relies on the researcher's ability to deduce the original forms and initial heights of old erosion surfaces and remnant

forms that have been greatly modified by subsequent processes. Moreover, dating erosion surfaces can be challenging due to the lack of valid geological evidence.

- In summary, denudation chronology is an approach that attempts to reconstruct the historical development of landscapes through the study of denudation processes. Despite its limitations, this approach provides valuable insights into the complex topographic features and geological history of a region, helping us better understand the Earth's surface and the processes that have shaped it over time.



Which of the following is a limitation of the denudation chronology approach? **A.** Difficulty in distinguishing between different types of rock formations **B.** Inability to explain the present landscape effectively **C.** Lack of understanding of geomorphological processes **D.** Ignoring the role of human activities in landform development

Conclusion

Denudation chronology is an approach in the study of landforms that focuses on reconstructing the historical development of landscapes through the analysis of erosion surfaces, drainage patterns, and geologic structures. This approach is based on the concept of palimpsest topography, which represents complex surfaces that have been shaped, erased, and reshaped by geomorphological processes over time. Despite its usefulness in understanding landscape evolution, denudation

chronology has its limitations, such as difficulties in accurately identifying and dating remnant landforms and the speculative nature of the approach due to the lack of valid geological evidence.

What is denudation and how does it affect Earth's surface?

Denudation is the process of wearing away the Earth's surface through the action of moving water, ice, wind, and waves. It leads to a reduction in elevation and relief of landforms and landscapes.

What is denudation chronology and why is it important in the study of landforms?

Denudation chronology is the reconstruction of the denudational history of a given region. It involves studying the traces left in present-day landscapes and sediments to understand the long-term landscape changes. This approach helps researchers to better understand the geological and geomorphological processes that have shaped a particular region over time.

What is the concept of palimpsest topography in denudation chronology?

Palimpsest topography refers to a landscape that has been shaped, reshaped, and modified by geomorphological processes over time,

similar to a manuscript that has been written, erased, and rewritten multiple times. This concept is essential in denudation chronology, as it helps researchers to recognize and interpret the complex topographic features that have evolved through different stages of geological time.

What are some limitations of the denudation chronology approach?

Some limitations of the denudation chronology approach include the difficulty in explaining only small parts of the existing landscape, the highly speculative nature of the approach due to the modification of old erosion surfaces, and the challenges in dating erosion surfaces due to the lack of valid geological evidence.

How does the denudation chronology approach contribute to our understanding of Earth's history?

The denudation chronology approach allows researchers to reconstruct the geomorphic history of a region based on present and remnant landforms. This helps in understanding the various geological and geomorphological processes that have occurred over time, providing insights into the Earth's history and the development of its landscapes.

1. What is denudation chronology?



Denudation chronology refers to the study of the sequence and timing of erosional processes that have shaped the Earth's surface over time. It involves understanding the rates of erosion, weathering, and sediment transport, as well as the factors influencing these processes.

2. How is denudation chronology determined?



Denudation chronology is determined through various methods, including the analysis of sediment deposition, dating of landforms, and the measurement of erosion rates. Techniques such as cosmogenic nuclide dating, optically stimulated luminescence dating, and stratigraphic analysis are commonly used to establish the chronology of denudation.

3. What factors influence denudation chronology?



Several factors can influence denudation chronology, including climate, tectonic activity, lithology (rock type), topography, and vegetation cover. For example, areas with higher precipitation and steeper slopes tend to exhibit higher erosion rates compared to regions with lower rainfall and gentler slopes.

4. Why is denudation chronology important in understanding landscape evolution?



Denudation chronology plays a crucial role in understanding the processes that shape landscapes and how they evolve over time. By studying denudation rates and patterns, scientists can gain insights into the geological history of an area, identify past climatic changes, and assess the impact of human activities on the landscape.

5. How does denudation chronology relate to the UPSC exam?




Denudation chronology is a topic that falls under the broader subject of Geomorphology, which is often included in the syllabus of the UPSC exam. Aspirants preparing for the exam may come across questions related to denudation chronology and its significance in understanding the Earth's surface processes.

Channel Morphology

Define Channel Morphology

- Channel morphology, also known as river channel morphology or river morphology, is a comprehensive study of river channels from a geographical perspective and fluid dynamics aspect. It involves examining various channel-related aspects such as channel patterns, geometry, and the factors that influence these forms.
- The factors that affect a channel are the processes that modify it, including the network of tributaries that connect to the main river channel within the drainage basin. Channel development is primarily controlled by two factors: water flow or discharge (in terms of volume and velocity) and sediment movement. These factors are influenced by the channel slope or gradient, which can range from steep to gentle slopes depending on the altitude variations the channel traverses.
- The interrelationship between these parameters can be qualitatively described by Lane's Principle (also known as Lane's relationship), which states that the product of the sediment load and bed grain size is proportional to the product of discharge and channel slope. In simpler terms, this principle highlights the balance between the forces of water flow and sediment movement that shape a river channel's morphology.




Channel

- A channel, in the context of physical geography, refers to the narrow pathway of a water body, such as a river or a strait. It is characterized by its bed and banks that outline the path of the river or stream. The entire network of river

channels, including their connecting branches in the form of tributaries, is proportional to the size of the valley they traverse.


- Channels can be occupied by various types of streams, including permanent ones that flow year-round, intermittent ones that flow at certain times, and ephemeral ones that only flow during and after rain events.
- During floods, the volume of water in a river may exceed the channel's capacity, causing the water to overflow its banks and flood the surrounding floodplain. In this sense, the term "channel" is often used interchangeably with "strait," which refers to a relatively narrow body of water connecting two larger bodies of water. In a nautical context, the terms "strait," "channel," and "passage" can be used interchangeably.
- For instance, in an archipelago, the water between islands is commonly referred to as a channel or passage. A well-known example of a channel is the English Channel, which is the strait separating England and France.



What are the two primary factors controlling channel development and morphology? **A.** Climate and vegetation **B.** Water flow and sediment movement **C.** Human activities and geology **D.** Channel slope and sinuosity

Channel Structure


- Channel structure refers to the various components that make up a river channel, including channel banks, channel bed, and the Thalweg.
- The channel bed is the primary pathway through which a river flows. It is the base of the river, on which the water runs and moves downstream.
- Channel banks are the two sides of the river bed that confine the water flow. These banks can be made up of various materials, such as soil, rocks, and vegetation, and help to shape the overall structure of the river channel.
- The Thalweg is a continuous line that connects the lowest points of the stream channel. It is essentially the deepest part of the river bed, where the flow of water is generally the fastest and the most erosive, often resulting in the formation of deeper channels over time.



River Processes

- River processes involve the wearing away, transportation, and deposition of materials by the river. These processes are essential for shaping the landscape and creating various landforms.
- Erosion is the process of gradually wearing away materials, such as rocks and soil, by the river's movement. There are four primary methods of erosion: attrition, corrosion, corrasion, and hydraulic action. Attrition occurs when the river's load, or the materials it carries, collide with one another, causing them to break apart. Corrasion, also known as abrasion, happens when the river's load grinds against the riverbanks and bed, breaking off pieces. Corrosion, or solution, is the process of the river's water dissolving its load, as well as the bed and banks. Hydraulic action refers to the erosion caused by water and air entering cracks in the riverbanks and bed, which increases pressure and breaks apart the materials.
- Transportation is the process of the river moving its load from one place to another, using excess energy. There are four main ways the river transports materials:

traction, saltation, suspension, and solution. Traction involves large pieces of load rolling along the river bed, while saltation is the process of smaller loads bouncing along the river bed. Suspension refers to tiny pieces of load being carried within the river's flow, and solution involves dissolved materials being transported within the water. Flotation is when materials are transported on the river's surface.



- The term "load" refers to the materials transported by the river. If the materials are being transported along the river bed, they are called bedload. If they are transported within the river's flow, they are often referred to as suspended load.
- Deposition occurs when the river no longer has enough energy to transport its load, causing it to put down or deposit the materials in its path. This process leads to the formation of various landforms, such as deltas and alluvial fans.

Factors Controlling Channel


Morphology

Channel morphology, which includes aspects such as channel pattern and movement, is influenced by two groups of factors: independent factors and dependent factors.

- **Independent factors** are those that are externally imposed on the watershed and are related to the landscape's characteristics. These factors include geology, climate, and human activities. The geology of a watershed is influenced by various processes at the landscape level, such as volcanism, tectonics, and to a smaller extent, surface processes like erosion and deposition. These factors affect the distribution, structure, and type of bedrock, surface materials, and topography within a watershed (Montgomery 1999).
- Climate is another key independent factor that influences channel morphology at the landscape scale. It plays a crucial role in determining the amount of rainfall and water flow in the stream channel. Human activities, such

as deforestation, construction, and agriculture, can also lead to significant changes in the watershed conditions.

- Dependent factors, on the other hand, are determined by the independent variables mentioned above. These include sediment supply, stream discharge, and vegetation (Montgomery and Buffington 1993; Buffington et al. 2003). Furthermore, time is also an essential independent variable that influences channel morphology.



Dependent Factors

Dependent factors in channel morphology are variables that change and adapt according to independent conditions. The morphology of a channel is influenced by the combined effects of these dependent landscape variables, which in turn respond to changes in these factors by adjusting one or more of the dependent channel variables.

- Sediment supply is one of the dependent factors and is determined by the frequency, volume, and size of material delivered to the channel. Stream discharge is another dependent factor, which is characterized by the frequency, magnitude, and duration of stream flows. The variation in stream discharge over time and space has a significant impact on channel morphology.
- Riparian vegetation is the third dependent factor that influences channel morphology. It helps control bank erodibility and near-bank hydraulic conditions, and also serves as a source of in-channel large woody debris (LWD). Traditional models of channel morphology primarily considered stream flow and sediment transport rate, assuming that transport rate and sediment supply were equal under equilibrium conditions. However, these models did not account for the role of vegetation or other boundary conditions, which are crucial in determining channel morphology.

In addition to riparian vegetation, important boundary conditions include elements found within the stream channel and those


that may affect the channel's ability to migrate laterally or build vertically. **Key boundary conditions include:**

- Channel gradient, which is determined by the valley slope. The maximum possible gradient of a stream channel depends on the slope of the valley.
- Bank composition and structure, which influence bank erodibility based on the sedimentology and Bedrock and other non-erodible units, such as colluvial material, compact tills, and lag glaciofluvial deposits, which can limit lateral and vertical channel migration and determine stream channel alignment.
- Erodible sediment stored in valley bottoms, including floodplains, fans, or terraces, as well as alluvial sediments, lacustrine, marine, and glacial outwash deposits, and fine-textured colluvium.
- Human channel alterations, such as bridge crossings and flood protection works.

These boundary conditions are primarily influenced by the geomorphic history of a landscape and the history of human intervention. As a result, the current morphology of a stream is a product of both present-day and historical watershed processes.

Channel Classification

- The channels can be classified into different categories based on different criteria. These criteria include the constituent material of the river channel and the shape or pattern of the river channel. The pattern of the channel is described as channel form. Channels exist in a variety of forms. There are a wide variety of stream channel types based on their forms, such as Single Thread Sinuous Rivers, Wandering Rivers or a meandering river, Braided Rivers, etc. enclosed by the materials of its bed and banks.




- Bedrock Channel:** When the river bed has the cover of rocks rather than the sediment cover and the river erodes into the rock. These are the channels that flow through

non-erodible materials (e.g., bedrock, coarse colluvium, and non-erodible glacial deposits) and their boundary conditions tend to dominate the channel morphology. This type of channel usually has a limited sediment supply and a morphology that is largely determined by the structure and composition of the material through which it flows. Bedrock channels, for example, frequently run along faults or other geologic planes of weakness within the rock. Overall, these channels are relatively insensitive to disturbances, including disturbances from changes occurring upstream (i.e., the channel is relatively stable), but bedrock channels are very effective at transferring disturbances from upstream to downstream reaches.

- Alluvial Channel: When the river cuts the river-transported rock debris, or alluvium, these are referred to as the alluvium channel. These channels are more regular.
- Schumm (1985) has proposed a more detailed channel classification that included three categories:
 - bedrock channel,
 - semi-controlled channel, and
 - alluvial channel;
- But this classification was found to be not addressing the variable geotechnical properties associated with the landforms.
- So Kellerhals in 1976, has suggested that the categories should be based on the materials that determine channel bed and bank strength and the channel's threshold of erodibility. Based on this, three categories of materials constituting channel could be identified (1) nonerodible, (2) semi-erodible, and (3) erodible. Although, these terms (as opposed to the conventional "non alluvial" and "alluvial") are more useful, but by definition these are contradictory, as all alluvial material is erodible and many non-alluvial materials are also highly erodible (e.g., marine and glaciofluvial deposits). Similarly, some alluvial materials are far less erodible than others; for instance, armored channel beds developed by fluvial processes are much more resistant to movement than other alluvium such as gravel-bar deposits, which are rearranged on an annual basis.
- The other classification is based on the shape assumed by the river channel, called a planform pattern. In the words of Leopold (1957) "Channel pattern is used to describe the plan view of a reach of river as seen from an airplane and includes meandering, braiding, or relatively

straight channels. Natural channels characteristically exhibit alternating pools or deep reaches and riffles or shallow reaches, regardless of the type of pattern." The shape of the channel is largely decided by the sinuosity of the river. Sinuosity refers to the ratio of the measured channel distance divided by the straight-line distance of the valley from the beginning of the channel reach to the end of the channel reach.

- Mollard (1973) identified 17 planform channel types that were related to both the physiographic environment in which channels flowed, and the materials that made up the channel bed and banks. He based this channel pattern classification on the factors controlling morphology, specifically streamflow, sediment supply, the relative dominance of fluvial transport processes, and the materials within which the channel is formed.
- Church (1992) classified channel patterns on the basis of the caliber and volume of sediment supply that in turn decides the sinuosity of the river channel. He has separated the patterns into phases of river channel flow during its upper course to lower course, related to how the supplied sediment gets transported.
- **Braided River Channel:** In other situations, the channel becomes too active for stable vegetated islands to develop, and the system divides into numerous individual channels that divide and recombine around unstable gravel bars; these are called "braided channels." Anastomosing rivers or streams are similar to braided rivers in that they consist of multiple interweaving channels but they typically consist of a network of low-gradient, narrow, deep channels with a stable bank.



Hydraulic Geometry of Channel

- The analysis of the relationship among stream discharge, channel shape, sediment load and slope, has been called as hydraulic geometry of stream channels.
- Gaging stations around the world maintained by the respective governments make recordings along the rivers all over the world. At these stations, the water surface level, channel shape, stream velocity, amount of dissolved and suspended minerals and other variables

are periodically recorded. These records provide a detailed history of river flow.

- In 1953, L.B. Leopold and Thomas Maddock published an elaborate analysis of thousands of measurements from stream gaging stations all over the world. It was they who called their analysis of the relationship among stream discharge, channel shape, sediment load and slope as hydraulic geometry of stream channels.
- To understand the hydraulic geometry of stream channels there is need to study the changes in following four parameters at different gaging stations under varying conditions spanning from low flow to bank full discharge and flood:

- **Channel width,**
- **Channel depth,**
- **Stream velocity,**
- **Suspended load**

- These four parameters increases as some small, positive power function of discharge, based on the following equations:

$$w=aQ^b \quad d=cQ^f \quad v=kQ^m$$

where Q stands for water discharge,
 w refers to water surface width,
 d stands for mean water depth, and
 v is the mean current velocity.

- The numerical values for arithmetic constants a , c , and k is not very significant for the hydraulic geometry of stream channels. They are constants of proportionality that convert a given value of discharge into the equivalent values of width, depth and velocity. The numerical values of the exponents b , f and m are very important as they describe the changes of width, depth and velocity for a specified change of discharge are very important.
- During flood, the water rises and stream channel width, depth and current velocity all increase at the gaging stations. This is clearly evident during the flood situation, as the regularity of the average changes drastically.
- In a downstream direction, the changes in channel shape and stream velocity are more significant as was emphasised by Leopold and Maddock in 1953. River discharge in humid area increases downstream.
- They proved that as mean discharge of a river increases downstream, channel width, channel depth, and mean current velocity all increase.

- Leopold and Maddock showed that not only the effluent rivers get both wider and deeper as they grow larger downstream but against the popular belief they also proved that average current velocity and increases downstream. This is because of the high-velocity current in a upper coarse of the river flows in circular eddies, with backward as well as forward motion.
- The numerical value of the three exponents b , f , and m that describe the variations in width, depth and velocity with variable discharge at gaging station are not the same as those that describe the downstream increases in width, depth, and velocity with progressively increasing mean annual discharge.
- Leopold and Maddock said that in a downstream direction, at mean annual discharge, the average values for the exponents were found to be $B=0.5$, $f=0.4$, and $m=0.1$.
- These values show that channel width increases most rapidly with mean annual discharge (as the square root of the discharge), depth next most rapidly, and mean velocity increases only slightly, in the downstream direction.
- Leopold and Maddock proposed that the increasing depth downstream permits more efficient flow in a river and compensates for the decreasing slope, thus providing a slight net increase in velocity at mean annual discharge.
- Thus, the channel width, depth, and velocity change with the streamflow, and against the popular belief, the velocity increases though only slightly in the downstream direction.

What are the four main ways a river transports materials?

A. Traction, saltation, suspension, and solution **B.** Attrition, corrosion, corrasion, and hydraulic action **C.** Erosion, deposition, transportation, and solution **D.** Traction, erosion, suspension, and deposition

Channel Stability and Movement

- Channel stability refers to a channel's tendency for vertical or lateral movement, as described by Church (2006). This stability is constantly changing as sediment is deposited or transported away, which can occur either along the banks of the stream or within the channel itself. These deposits can lead to alterations in the channel's

patterns, and any noticeable changes in these patterns can signal significant shifts in both the watershed and the factors that influence its morphology.

- Geomorphologists can provide evidence of these channel changes, which can then be used by researchers as an indicator of the environmental health of the watershed. It is important to note that different channel types respond differently to changes in sediment supply or discharge. These responses can manifest as either vertical shifts (such as aggradations or degradation, which can be observed in the upward or downward movement of the channel bed) or lateral shifts (where the bed and banks move sideways, as seen in old or abandoned channels on a floodplain).
- In summary, channel stability and movement are crucial factors in understanding the health of a watershed. By observing changes in channel patterns and how different channel types respond to sediment supply and discharge, researchers can gain valuable insight into the overall environmental health of the region.

Vertical Channel shifts:

- Vertical channel shifts refer to the changes in the width-to-depth ratio of a river channel. These shifts occur as a response to variations in sediment load and transportation capacity. When the sediment supply exceeds the river's ability to transport it, the excess sediment accumulates and causes aggradation. Conversely, when the sediment supply is lower than the river's transportation capacity, sediments are removed, leading to degradation.
- Aggradation is the process wherein sediments are deposited within the river channel, causing the width-to-depth ratio to increase. This in turn reduces the stability of the channel. An aerial view of a river experiencing aggradation would show a noticeable expansion of sediment deposits within the channel over time.
- On the other hand, degradation refers to the removal of sediments from the river channel, which happens when the river's transportation capacity exceeds the sediment supply. This process can lead to the narrowing and deepening of the channel.

In summary, vertical channel shifts are changes in the width-to-depth ratio of a river channel as a result of variations in sediment load and transportation capacity. Aggradation occurs when sediment supply surpasses the river's capacity to transport it,

leading to sediment deposition and a decrease in channel stability. Degradation, conversely, takes place when sediment supply is lower than the river's transportation capacity, leading to the removal of sediments and potential channel narrowing.

Lateral Channel movement

- Lateral channel movement refers to the shifting of a river or stream across a flat valley surface, which can indicate changes in the conditions upstream. This movement can be caused by progressive bank erosion, which is the gradual wearing away of the riverbank due to sediment buildup in the channel or natural meandering processes. On the other hand, channel avulsion is a more sudden and significant shift in the channel's position to a new or old part of the floodplain or a minor change within a braided channel or other active channels.
- The lateral movement of a channel affects the riparian zone, which is the area where land meets a river or stream. This movement can cause erosion in some areas and buildup in others. The limits of lateral channel movement are determined by the channel's boundary conditions and its relationship with the valley it flows through. If there are no constraints like valley confinement, bridges, or dykes, and the valley is filled with erodible material, the channel can potentially erode across its entire floodplain.
- When a valley is wider than the channel, there is potential for lateral channel movement. In confined systems where the valley is only slightly wider, the extent of lateral movement is limited. In forested valleys, the bank strength provided by riparian vegetation can restrict lateral channel movement, allowing for stable channel morphology in environments where it might not otherwise occur.
- Changes in the patterns of in-channel sediment storage, such as bars and islands, can also serve as early indicators of future channel issues. Braided channels are among the most active types of stream channels, characterized by rapid lateral migration rates and often experiencing net vertical aggradation, or the buildup of sediment.

Conclusion

Channel morphology is an essential aspect of studying river channels from both geographical and fluid dynamics perspectives. It encompasses various factors that influence the shape and structure of channels, such as water flow, sediment

movement, and the slope of the channel. Channels can be classified based on their constituent materials or their planform patterns, such as straight, sinuous, or meandering channels. Understanding channel morphology is vital to comprehending the processes and factors that shape our landscape and contribute to the formation of diverse landforms.

What is channel morphology?

Channel morphology, also known as river channel morphology or river morphology, is the study of river channels from a geographical perspective and fluid dynamics aspect. It involves examining various aspects of the channel, such as patterns, geometry, and the factors that influence these forms.

What are the main factors that control channel morphology?

Channel morphology is primarily controlled by two factors: water flow or discharge (in terms of volume and velocity) and sediment movement. These factors are influenced by the channel slope or gradient, which can range from steep to gentle slopes depending on the altitude variations the channel traverses.

What is the difference between bedrock channels and alluvial channels?

Bedrock channels are those that flow through non-erodible materials, such as bedrock, coarse colluvium, and non-erodible glacial deposits. Their morphology is largely determined by the

structure and composition of the material through which they flow. Alluvial channels, on the other hand, are those that flow through river-transported rock debris or alluvium. These channels are more regular and are more sensitive to changes in sediment supply and flow conditions.

How are river channels classified based on their shape or planform pattern?

River channels can be classified into four types based on their planform pattern: straight river channels, sinuous river channels, meandering river channels, and wandering river channels. These patterns are determined by the sinuosity of the river, which refers to the ratio of the measured channel distance divided by the straight-line distance of the valley from the beginning of the channel reach to the end of the channel reach.

What are some of the processes involved in shaping river channels?

River processes that shape channels include erosion, transportation, and deposition. Erosion is the gradual wearing away of materials by the river's movement, transportation is the process of the river moving its load from one place to another, and deposition occurs when the river no longer has enough energy to transport its load, causing it to put down or deposit the materials in its path. These processes lead to the formation of various landforms, such as deltas and alluvial fans.

1. What is channel morphology?



Ans. Channel morphology refers to the physical characteristics and features of a river channel, including its shape, size, slope, and cross-sectional profile. It describes the overall form and structure of a river channel and how it changes over time due to various natural and human-induced factors.

2. What factors control channel morphology?



Ans. Several factors control channel morphology, including the type and amount of sediment available in the river, the flow velocity and discharge of water, the slope of the channel, the presence of vegetation, and the geological characteristics of the surrounding landscape. Human activities, such as dam construction and land use changes, can also significantly influence channel morphology.

5. What are some frequently asked questions about channel morphology?



Ans. Some frequently asked questions about channel morphology include: - How does human activity affect channel morphology? - What are the major impacts of channel morphology on ecosystems and human communities? - How can channel morphology be managed or restored to prevent flooding and erosion? - What are the different methods used to study and analyze channel morphology? - How does climate change influence channel morphology and its associated hazards?

3. How are channels classified based on morphology?



Ans. Channels can be classified into different types based on their morphology. The classification is often based on the shape and pattern of the channel, such as straight, meandering, or braided channels. Channels can also be classified based on their size, including small streams, rivers, and large-scale river systems. Additionally, channels can be categorized based on their gradient, bed material, and hydraulic characteristics.

4. What is lateral channel movement?



Ans. Lateral channel movement refers to the horizontal shifting or migration of a river channel over time. It occurs when a river erodes its banks on one side and deposits sediment on the other side, leading to the gradual shifting of the channel. Lateral channel movement can be influenced by factors such as changes in flow patterns, sediment supply, and the presence of vegetation.

Erosion Surfaces

Erosional Surfaces


Erosion or planation surfaces are the outcomes of denudation processes, which result in undulating ground surfaces and remnants of high relief cutting across geological formations and structures. These surfaces are essential for understanding the development of landforms as well as the past geomorphic processes and evolutionary history of landscapes. They can provide insight into the denudation chronology.

- Various scholars have used different terms to describe planation surfaces, including "peneplain" (Davis), "pediplain" (L.C. King), "panplain" (Crickmay), "etchplain" (Wayland 1934, Thomas), "panfan" (A.C. Lawson), "primarrumpf," "gipfelflur," "stockwerk," "schietelflur," and "stripped surface" (Budel 1957). Some minor erosion surfaces include valley-side beaches (terraces), river terraces, and marine benches (platforms).
- Although the study of erosion surfaces became less popular during the second half of the 20th century, the current consensus is that they do exist, and interest in them has been revived. Erosional plains are planation surfaces that cut across various bedrock types and geological structures, and they can be found in both low-lying areas and at higher elevations. Elevated plains

often show signs of an erosional origin, followed by subsequent dissection, like a bevelled cuesta.

- A cuesta is a hill or ridge with a gentle slope on one side and a steep slope on the other. The flat top or bevel on a cuesta can be seen as evidence of an upper erosion surface that existed before differential erosion shaped the cuesta. However, it is essential to note that a single bevelled cuesta does not necessarily indicate a planation surface. It could also be a river terrace or another small flat feature. A planation surface is more likely when multiple bevelled cuestas occur at about the same elevation.

In summary, planation surfaces, also known as erosional surfaces or paleoplains, are the result of various erosion agents, such as wind, running water, and glaciers, acting in different climatic regions. These geographically plain surfaces or faint reliefs are produced during the final phase of the erosion cycle and are essential for understanding the evolution of landforms and the history of geomorphic processes.



Peneplains

- The concept of a peneplain, which means "almost a plain," was developed from W.M. Davis' cyclic view of landscape evolution. According to Davis, as rivers and hillslopes progress through the stages of youth, maturity, and old age, they gradually reduce the relief of the landscape, ultimately creating a plain with extremely low relief. The rate of change in this plain is minimal due to the reduced potential energy for fluvial action.
- In the youthful stage, valleys are typically V-shaped. As the landscape matures, valleys become flat-bottomed due to the dominance of lateral erosion. In the old age stage, valleys have very shallow features as they become part of extensive plains, with lateral erosion eroding away all hills.
- Young landscapes are characterized by large areas of flat topography, which are remnants of the original uplifted peneplain. As landscapes mature, the valleys deepen and widen, creating V-shaped valleys that consume much of the interfluvies, or areas between river systems, which still bear traces of the original land surface. Old landscapes are characterized by peneplains, where the interfluvies are reduced to minor undulations, creating a nearly flat landscape.

Which landscape feature is primarily formed by chemical weathering and weakly acidic water lowering the weathering front? **A. Peneplain B. Pediplain C. Etchplain D. Panfan**





Fig. 16.1 : Graphical presentation of geographical cycle presented by W.M. Davis.


DAVICIAN CYCLE OF EROSION



- A. YOUTH
- B. MATURITY
- C. OLD AGE


Pediplains

- The pediplain concept was developed by Lester Charles King, who built upon Penck's model of slope retreat in his own model of landscape evolution. King's idea was that slope retreat creates pediments, and when enough pediments are formed, a pediplain is the result. He envisioned this process occurring in "cycles of pedimentation."
- Each cycle starts with a sudden burst of cymatogenic diastrophism, followed by a period of diastrophic quiescence. During this calm period, subaerial processes reduce the relief to a pediplain. As the continent is eroded, the sediment is deposited offshore. As sediment is removed from the continental margins, they rise, while the weight of the sediment in offshore regions causes depression.
- This simultaneous uplift and depression lead to the formation of a major scarp near the coast, which then cuts back inland. As the scarp retreats, it leaves behind a pediplain, further unloading the continent and placing additional sediment offshore.
- Over time, new cycles of uplift and depression occur, creating new scarps. This cyclical relationship between continental unloading and offshore loading results in landscapes consisting of a large staircase of erosion surfaces or pediplains, with the oldest steps found deep inland.



Endrumpf

- In the landscape evolution model proposed by Walter Penck, the endrumpf refers to the final stage of erosion that occurs after inselbergs (isolated hills or mountains) and their surrounding pediments (gentle slopes) have disappeared. The endrumpf is characterized by a landscape with low-angle, concave slopes that gradually retreat over time.
- Penck's model is a variation on the earlier model proposed by William Morris Davis, who believed that uplift and planation (the process of creating a flat surface) occurred alternately. However, in many landscapes, uplift and denudation (the wearing away of the earth's surface) actually occur simultaneously. This led Penck to develop a different model of landscape evolution, in which the evolution of individual slopes plays a crucial role in determining the overall landscape's development.
- In Penck's model, the continuous and gradual interaction between tectonic processes (such as uplift and earth movement) and denudation results in a unique landscape evolution pattern, where the endrumpf stage is reached after the inselbergs and their surrounding slopes have been eroded away.




Panplains

- Colin Hayter Crickmay (1933, 1975) introduced the concept of unequal activity in slope retreat, which deviates from the models of landscape evolution proposed by Davis, Penck, and King, who assumed that slope processes act evenly on individual slopes. In reality, geomorphic agents do not act uniformly.
- As a result, a slope may only recede in areas where its base is eroded by a stream (or the sea). In this case, slope denudation is primarily achieved through the lateral erosion of rivers (or marine erosion at the base of cliffs), meaning that some parts of the landscape may remain virtually unaffected by slope recession. Crickmay argued that this lateral erosion by rivers leads to the formation of Panplains.

Etchplains and Etched Surfaces


- Landscape evolution was traditionally thought to be primarily influenced by mechanical erosion. However, recent research indicates that chemical weathering plays a significant role in shaping landscapes, especially in tropical and subtropical environments. One such process is etch planation, which involves the formation of a thick regolith through chemical weathering, followed by erosion.
- An etchplain, or etched peneplain, is a landscape feature resulting from etch planation. The concept was first introduced by E.J. Wayland and
- Bailey Willis in the 1930s and later refined by Julius Budel. Etchplains are mainly formed by chemical weathering, with weakly acidic water lowering the weathering front, similar to how an acid-soaked sponge would etch a metal surface.
- Some researchers propose the theory of double planation, which suggests that surface erosion and chemical etching occur at the same rate, maintaining land surfaces of low relief during slow uplift. This theory involves two planation surfaces: the wash surface and the basal weathering surface (etch surface).
- There are various types of etchplains, ranging from stripped etchplains to those covered with thick waste. One example is the Chotanagpur region, where the bedrock is mostly hidden beneath a layer of soil and regolith. Laterite beds can be found at the surface, acting as a cap rock and helping to maintain even topography.
- Thomas identified five types of etchplains: undissected lateritized surface, dissected lateralized etchplain, extensively stripped surface, partially stripped erosion surface, and incised erosion surface. However, there may be other variations, such as the one seen in Chotanagpur.
- Etchplains can also form on non-crystalline sedimentary rocks, though some researchers question the distinction between etchplains and pediments. Despite this, the term "etchplain" is still used by some geomorphologists, as it helps to explain landscape features found in warmer climates, now present in temperate latitudes.



Panfan

- Lawson coined the term "**Panfan**" to describe the final stage of geomorphic development in arid regions, similar to how a "peneplain" represents the end stage of landform evolution in humid regions.

Which of the following terms was proposed by Lester Charles King as a result of slope retreat creating pediments? **A. Peneplain B. Pediplain C. Endrumpf D. Panplain**



Conclusion

Planation surfaces, including peneplains, pediplains, endrumpf, panplains, etchplains, and panfans, are essential for understanding the evolution of landforms and the history of

geomorphic processes. These surfaces result from various erosion agents and climatic conditions, shaping the landscape through mechanical and chemical processes. The study of these features provides valuable insights into the denudation chronology and the interplay between tectonic processes and erosion in shaping the Earth's surface.

What are erosional surfaces, and why are they important in understanding the evolution of landforms?

Erosional surfaces, also known as planation surfaces or paleoplains, are the result of various erosion agents, such as wind, running water, and glaciers, acting in different climatic regions. These geographically plain surfaces or faint reliefs are produced during the final phase of the erosion cycle and are essential for understanding the evolution of landforms and the history of geomorphic processes.

What is the difference between a peneplain and a pediplain?

A peneplain is a landscape with extremely low relief that forms as rivers and hillslopes progress through the stages of youth, maturity, and old age, gradually reducing the relief of the landscape. A pediplain, on the other hand, is formed when enough pediments (gentle slopes) are created as a result of slope retreat, often occurring in "cycles of pedimentation."

What is an etchplain, and how does it differ from other planation surfaces?

An etchplain, or etched peneplain, is a landscape feature resulting from etch planation, which involves the formation of a thick regolith through chemical weathering, followed by erosion. Etchplains are mainly formed by chemical weathering, with weakly acidic water lowering the weathering front, similar to how an acid-soaked sponge would etch a metal surface. Etchplains differ from other planation surfaces as they primarily result from chemical weathering rather than mechanical erosion.

What is the role of chemical weathering in the formation of etchplains and etched surfaces?

Chemical weathering plays a significant role in the formation of etchplains and etched surfaces, as weakly acidic water lowers the weathering front, similar to how an acid-soaked sponge would etch a metal surface. This process forms a thick regolith through chemical weathering, which is then eroded to create etchplains and etched surfaces.

What is a panfan, and how does it relate to the concept of a peneplain?

A panfan is the final stage of geomorphic development in arid regions, similar to how a peneplain represents the end stage of landform evolution in humid regions. Lawson coined the term "panfan" to describe this final stage in arid environments, where the landscape is shaped by different erosional processes compared to those in humid regions.

1. What are erosion surfaces?



Ans. Erosion surfaces refer to flat or gently sloping surfaces that are formed by the process of erosion over a long period of time. They are characterized by their smooth and level appearance, and are often found in areas that have undergone significant erosion.

2. How are erosion surfaces formed?



Ans. Erosion surfaces are formed through the continuous action of weathering and erosion over a long period of time. The process starts with the weathering of rocks and the subsequent transportation of the resulting sediments by various agents such as wind, water, or ice. As these agents erode the surface, they gradually create a flat or gently sloping landform.

3. Where are erosion surfaces commonly found?



Ans. Erosion surfaces can be found in various regions around the world. They are often found in areas with ancient or highly weathered landscapes, such as plateaus, mountains, or deserts. Examples of famous erosion surfaces include the Colorado Plateau in the United States and the Nullarbor Plain in Australia.

4. What are the features of erosion surfaces?



Ans. Erosion surfaces typically exhibit certain common features. They are generally flat or gently sloping, with a smooth and level appearance. They may have remnants of ancient river channels or valleys, indicating the previous presence of water. Erosion surfaces can also have distinct layers of sedimentary rocks, showing the different stages of erosion over time.

5. Why are erosion surfaces important?



Ans. Erosion surfaces are important for understanding the geological history of a region. By studying these surfaces, geologists can gain insights into past environmental conditions, the effects of climate change, and the evolution of landscapes over millions of years. Erosion surfaces also provide valuable information for industries such as mining and oil exploration, as they can indicate the presence of valuable mineral deposits.

Slope Development

Slopes

- The study of slopes in the physical landscape has long fascinated geomorphologists, as they hold a central role in the study of landforms. Despite their importance, the study of slopes has been largely neglected and faces numerous challenges.
- One of the main difficulties in studying slopes is determining their nature, the rate at which processes operate, and the impact of these processes on the slope. Additionally, it is difficult to trace the entire development of a slope and its changes in form over time.
- In landform studies, there are two main aspects that are typically focused on: form and process. "Form" refers to the morphology of a given region at a given time, while "process" refers to the operation of various agents that cause changes in the physical environment, such as soil creep, surface wash, and weathering.
- To better understand slope development, researchers have followed two main approaches: the historical approach and the process-form approach.
- The historical approach emphasizes the evolution of slopes from their origin to their present form. However, this approach suffers from the problem of accurately reconstructing past forms of the slope, as there is no standard method for verifying the accuracy of these reconstructions. It is also challenging to determine the age of a slope, as there is no universally applicable method for doing so.
- One commonly-held assumption among some researchers is that present-day slopes developed from near-vertical cliffs that weathered and modified over time. However, this assumption may not hold true in many cases, as slopes often border river valleys that were never vertical cliffs. Furthermore, river erosion does not always produce steep, vertical walls, as other processes like weathering and creep can also modify slope edges.
- As a result, the concept of an "initial slope" may be unrealistic. Determining the age of a slope based on its

form is not an easy task, and even extensive field surveys cannot provide completely satisfactory results.

The Process-form Approach

- The Process-form Approach is based on the idea that the shape and steepness of a slope are determined by the interplay of weathering, erosion, transportation, and deposition. These denudation processes work in various combinations and at different rates, leading to a wide array of slope forms with varying degrees of steepness. Factors such as rock types, climate, and vegetation directly influence the types of slope forms that are produced.
- For instance, in a limestone region with sufficient rainfall, convex slopes are commonly found because rainwash is less effective due to the porous nature of the rock. In other regions, this might result in concave slopes. However, this approach, like the historical approach, faces several challenges. It is difficult to observe the various processes at work on slopes since weathering, creep, and rainwash occur very slowly and are not easily perceptible.
- To accurately study these processes, advanced tools and precise methods are needed for recording their operation. Moreover, it is hard to definitively establish that slope processes directly determine the form of the denudational slope. Another issue with the process-form approach is the possibility that many current slopes are not a result of present-day processes (Small, 1978). The relationship between slope form and climate has also been a subject of debate.
- If the slope is solely controlled by slope processes, it is assumed that different slope forms would be found in different regions. Some geomorphologists have emphasized this relationship, providing various examples to support their claims. However, the assumption that a specific type of slope will only form in a particular climate is not always accurate. For example, it was once believed that pediments are exclusive to desert regions,

but now many scientists and geomorphologists agree that they can occur anywhere in the world.

Which of the following slope elements is characterized by a steep, wall-like profile and is typically found below cliffs or free faces? **A. Convex Slope B. Rectilinear Slope C. Concave Slope D. Summital Convexity**

Genetic Classification of slopes

Slopes are formed through two primary processes: Endogenetic and Exogenetic. Based on these processes, slopes can be broadly categorized into Endogenetic Slopes and Exogenetic Slopes.

Endogenetic Slopes

- These slopes are formed as a result of processes occurring within the Earth's interior. Various geological movements, such as folding, faulting, and rift valley formation, contribute to the development of these slopes. They are also commonly referred to as tectonic slopes. Fault scarps are often found in association with faults and rift valleys.
- Volcanic eruptions, which are also a consequence of endogenetic processes happening within the Earth, create new landforms. These eruptions lead to the accumulation of lava and pyroclastic materials, forming various volcanic hills, plateaus, and cones.
- The landforms created by volcanic eruptions or tectonic activities are further modified by subaerial processes, resulting in the development of diverse slope forms.


Exogenetic slopes

- Slopes are formed as a result of external processes that take place at or near the Earth's surface. Factors such as weathering, mass wasting, erosion, and deposition play a crucial role in shaping landscapes. These processes constantly act on the surface, creating and modifying slope forms over time.
- We can classify slopes created by exogenetic processes into two main categories: (a) erosional or degradational slopes and (b) depositional or aggradational slopes.

- Erosional slopes are created by various forces, including wind, running water, waves, and glaciers. For example, glaciers and running water generate numerous landforms within their valleys. Escarpments, watersheds, river terraces, and coastal cliffs are all examples of slopes formed through erosion.
- On the other hand, depositional slopes are formed by the same agents but through the accumulation of material. For instance, running water creates alluvial fans and natural levees, glaciers form moraines, and wind results in sand dunes of various shapes and sizes. Additionally, deposition along coastlines leads to the formation of sandbars, barriers, and beaches. These are all examples of depositional slopes created by the actions of water, wind, and glaciers.


Slope elements

- Slope elements refer to the various forms or shapes that make up a slope's profile. These forms typically include convex (crest), rectilinear, and concave slopes. Convex slopes are usually found at the top of a hill, while concave slopes are located at the base.
- L.C. King and A.Wood believed that a standard composite slope profile consists of four main elements. The first, at the top, is the convex slope or crest. Below the crest, there is a scarp or free face, followed by a rectilinear slope. Finally, at the bottom, there is a concave slope. This is the most common composite slope profile; however, in reality, these elements can be found in various combinations, and not all of them may be present in a single slope profile.
- The combination of these elements in a slope profile depends on factors such as the rock's structure, the rock's nature, and the processes that operate on the surface. Below, we will discuss each of the four main types of slope elements (forms) in greater detail.



Convex Slope


- A convex slope is a type of slope that can sometimes cover an entire hillside, but it is most frequently found in the upper portions of a slope. This convex shape is created by the gradual wearing away of the landscape, known as the denudation process, and is typically found in humid, temperate regions.
- In the higher areas of a hill, a convex slope is often called a "crest" or "summit slope." As you move down the slope from the crest, the angle of the slope becomes steeper. The formation of this convex shape at the summit is primarily attributed to weathering and soil creep, which are the most active processes in creating this unique topography.
- This type of slope is also known as a "waxing slope," a term coined and popularized by the renowned German geomorphologist W. Penck. The term "summital convexity" is often used to describe a waxing slope, as it highlights the distinct curved shape of the slope's upper portion.



Cliff or Free Face


- A cliff, also known as a scarp or free face, is a steep, wall-like slope often found in various landscapes. Due to its steepness, cliffs are typically bare, as regolith or debris cannot accumulate on such a slope. Instead, any material that falls or erodes from the cliff settles at the base of the slope, creating a pile of debris.
- The term "free face" is used by many geomorphologists to describe a cliff, as it remains free from any detritus or debris. Cliffs can form along coastlines, where sea waves cause undercutting, in river valleys, glacial regions, faulted landscapes, and in various other locations. When weathered material falls or slides from the cliff, it accumulates at the base, forming a depositional feature known as a talus slope.
- Over time, the talus slope will continue to grow in size as more weathered material accumulates, eventually covering the lower portions of the free face and protecting it from further weathering. As the talus slope

risks, the length of the free face diminishes, and the entire cliff may eventually disappear, replaced by an aggradational slope with a lesser angle than the original cliff.




Rectilinear Slope

- Rectilinear slopes are a type of slope that is primarily found below cliffs or free faces, and are characterized by their straight profile and relatively constant slope angle. These slopes can vary in size and may sometimes dominate an entire slope area. They often extend from the summit of a hill or mountain down to the bottom of a valley.
- In many cases, rectilinear slopes are situated in the middle of a slope profile, with a broader convex section above and a larger concave section below. Some geomorphologists believe that these slopes are formed only through the process of aggradation, which is the accumulation of sediment or other material. However, this is not always the case.
- As noted by Small (1978), rectilinear slopes can also be a result of denudation, which is the wearing away of the Earth's surface by natural processes like weathering and erosion. In these instances, the slope is underlain by solid rock and may be covered by a thin layer of debris that is slowly moving downhill due to factors like frost and other natural forces.
- Rectilinear slopes are sometimes referred to as debris-controlled slopes, as the presence of debris can play a significant role in their formation and appearance. Additionally, Strahler (1950) used the term "repose slope" to describe these slopes, highlighting their unique characteristics and distinguishing them from other types of slope formations.




Concave slope

- A concave slope is found at the lowest part of a hill's slope profile, typically extending from the bottom of the hill down to the river valley. This area is often covered with debris, such as accumulated scree, which is spread across the slope due to rainwash. As a result, finer particles are dispersed further than coarser ones, leading to the formation of a concave shape.
- The term "waning slope" was coined by Penck to describe such slopes, which can display different characteristics depending on the climate. In arid and semi-arid regions, concave slopes often show a sharp break in gradient between the lower concave section and the steeper slopes above. However, in humid conditions, the concave base transitions more smoothly into the higher slopes.




The above-discussed elements are assumed to be present in a standard hill slope but as has been pointed out by many geomorphologists and thinkers that **all four elements may not be noticed in a hill slope. One or more than one element may be missing from the slope profile owing to various reasons.**



A slope profile can consist of various combinations of elements, and while there may be an infinite number of possible combinations, some are more common than others. Three frequently occurring composite slope combinations are discussed below:

- The convexo-rectilinear-concave slope profile features an upper convex portion, a middle rectilinear section, and a

lower concave part. These three slope elements blend seamlessly, resulting in a curved slope profile (Small, 1978). This type of slope profile is often found in regions with weaker rock types, such as lowland England. The lengths of the different slope elements can vary, creating diverse landscapes. However, in areas with a wide variety of rock types, where hard and soft rocks alternate, or where the region has experienced multiple rejuvenations, the slope profile can become more complex.



- In regions where there are alternating layers of massive and weak strata, high relief, deep valleys, and active weathering, a distinct composite profile can be observed. This profile consists of several free faces and rectilinear slopes, with minimal or no summital convexity and basal concavity (Small, 1978). Massive strata contribute to the formation of free faces, while weak and thinly bedded rocks result in rectilinear slopes.
- In arid regions with hard crystalline rocks, a composite slope profile is formed, which includes an upper section with a free face at a slope of 40° or more, a mid-section with a boulder-controlled slope at an angle of 25° or more, and a lower section with a gentle concave (pediment) slope at angles below 7° .
- Various theories have been proposed by geomorphologists to explain the development of specific slope elements in a slope profile. Earlier attempts focused on linking certain processes to specific slope forms, such as rainwash and soil creep in the formation of convex and concave slope forms.
- American physiographer N. M. Fenneman suggested that the most common convexo-concave profiles are due to the action of running water. According to Fenneman, the upper slopes have less runoff during rainfall, with water moving as a thin sheet. As the water travels downslope, it collects particles and gains volume from additional run-off from higher up the slope. This leads to increased erosion

in the sections of the slope further from the summit, resulting in convexity over time.

- Fenneman also argued that as the water reaches the lower part of the slope, the increased volume causes it to become concentrated in small channels, leading to the formation of gullies and ultimately a concave curve. However, many geomorphologists criticized Fenneman's hypothesis for not considering the role of soil creep in slope formation. Despite this, his ideas were supported by Horton (1945), who argued that erosion by sheet wash becomes more significant downslope due to increased run-off.
- Gilbert (1909) attributed soil creep as a major factor in the rounding of hilltop summits and the development of summital convexity, but his ideas were considered simplistic. Lawson, on the other hand, argued that rainwash was an important process on the upper slope, although he disagreed with Fenneman, suggesting that wash was most effective at the sloping summit.
- Many other scientists and geomorphologists have considered soil creep and rainwash as crucial processes in determining slope form, each proposing their own theories based on their understanding. However, it is important to note that, in addition to soil creep and rainwash, there are numerous other factors at play, and their interactions can be quite complex.

In which type of region is a composite slope profile consisting of an upper free face, a mid-section boulder-controlled slope, and a lower gentle concave slope most likely to be found? **A.** Humid, temperate regions **B.** Arid regions with hard crystalline rocks **C.** Lowland England **D.** Coastal areas

Conclusion

The study of slopes and their development is a complex field in geomorphology, with various factors and processes contributing to their formation. Two main approaches, the historical approach and the process-form approach, have been used to understand slope development. Slopes are classified into endogenetic and exogenetic types based on their formation processes, and their profiles consist of various combinations of convex, cliff or free face, rectilinear, and concave slope elements. The interplay of factors such as rock types, climate, and vegetation, as well as processes like weathering, erosion, and deposition, contribute to the diverse forms of slopes

observed in nature. While progress has been made in understanding these processes, there is still much to learn about the intricacies of slope development and the interactions between various factors and processes.

What are the main challenges in studying slopes in the physical landscape?

The main challenges in studying slopes include determining their nature, the rate at which processes operate, the impact of these processes on the slope, and tracing the entire development of a slope and its changes in form over time.

What are the two main approaches to studying slope development?

The two main approaches to studying slope development are the historical approach, which emphasizes the evolution of slopes from their origin to their present form, and the process-form approach, which focuses on the interplay of weathering, erosion, transportation, and deposition in determining the shape and steepness of a slope.

What are the two primary categories of slopes based on their formation processes?

Slopes can be broadly categorized into Endogenetic Slopes, which are formed as a result of processes occurring within the Earth's interior, and Exogenetic Slopes, which are formed as a result of external processes that take place at or near the Earth's surface.

What are the four main types of slope elements (forms)?

The four main types of slope elements are convex slope, cliff or free face, rectilinear slope, and concave slope. These elements can be found in various combinations in a slope profile, and not all of them may be present in a single slope profile.

How do soil creep and rainwash contribute to the formation of slope forms?

Soil creep, which is the slow, continuous movement of soil and debris down a slope, contributes to the rounding of hilltop summits and the development of summital convexity. Rainwash, on the other hand, is the washing away of soil and debris by rainfall, and can contribute to the formation of both convex and concave slope forms depending on the location and intensity of the runoff.

1. What is the genetic classification of slopes?

Ans. The genetic classification of slopes refers to how slopes are formed based on their origin or formation processes. The different genetic classifications of slopes include convex slopes, cliff or free face slopes, rectilinear slopes, and concave slopes.

2. How are convex slopes formed?

Ans. Convex slopes are formed by the accumulation of loose materials, such as debris, rocks, or soil, at the base of a slope. This accumulation gradually builds up and creates a convex shape. Convex slopes are commonly found in areas with high erosion rates or where there is a deposition of materials.

3. What are cliff or free face slopes?

Ans. Cliff or free face slopes are steep slopes that have a vertical or near-vertical face. They are typically formed through processes such as weathering, erosion, or tectonic activities. Cliff slopes are commonly found in mountainous regions, coastal areas, or areas with steep rock formations.

4. How are rectilinear slopes formed?

Ans. Rectilinear slopes are formed by the movement of large blocks or masses of rocks or soil along distinct planes or fractures. These slopes have a series of steps or terraces due to the movement of these blocks. Rectilinear slopes are often found in areas with jointed or fractured rock formations.

5. What are concave slopes?

Ans. Concave slopes are characterized by a rounded or bowl-shaped appearance. They are formed by erosion processes, such as water or wind erosion, that remove materials from the lower parts of the slope. This erosion creates a concave shape over time. Concave slopes are common in hilly or mountainous regions with softer rock or soil materials.

Applied Geomorphology

Introduction - Applied

Geomorphology

- Understanding landforms is crucial for humans, as it directly or indirectly affects their interaction with the earth's surface features. By correctly interpreting landforms, we can gain insights into the

- geological history, structure, and composition of a region. Applied geomorphology, as defined by D.K.C. Jones, is the process of utilizing geomorphic knowledge to address issues related to land use, resource management, and environmental planning.
- According to R.G. Craig and J.L. Craft, all geomorphological knowledge tends to have practical applications. As our understanding of the earth's processes improves,

geomorphologists can use this knowledge to assess resources, plan development projects, identify potential natural hazards, and reduce the impact of natural disasters. In essence, the study of landforms plays a significant role in managing and planning our interactions with the earth's surface.

- **Geomorphic knowledge and techniques may be applied in the following areas:**
 - Examining the effects of geomorphic and environmental processes on human society and activities, as well as addressing the challenges that result from these impacts;
 - Exploring the alterations in geomorphic and environmental processes caused by human actions and managing the issues that arise from such interactions.
 - Overseeing resource management and tracking changes in the geomorphic system to propose appropriate solutions that ensure sustainable development.

Two Main Lines of Application

The application of geomorphology, according to Charley, Schumn, and Sugden, may be considered along two lines:

Resource evaluation, engineering construction and planning

- In the field of applied geomorphology, various applications can be found, such as resource inventories, environmental management, soil and land evaluation, and producing maps for hydrological, erosional, and stability control. This field also encompasses geomorphic mapping, land system mapping, terrain evaluation, and information retrieval on terrain and other relevant subjects for earth scientists, engineers, and planners.
- Applied geomorphology plays a crucial role in urban planning across diverse geomorphic environments. It aids in the creation of natural hazard maps, morpho-agricultural regionalization, land use planning, and the construction and management of roads. By utilizing applied geomorphology, professionals can better understand and address the challenges posed by various landscapes and environments, ultimately leading to more informed decision-making in urban planning and development.

Human beings as geomorphic agents:

- Applied geomorphology is a field that examines the impact of human activities on the Earth's surface, particularly in terms of intentional or unintentional effects on the various processes and forms that shape the landscape. Throughout history, humans have sought to control and manipulate these natural processes to meet their economic needs.
- For example, people have constructed embankments to prevent river flooding, straightened and redirected river courses, built walls to protect coastal areas from wave erosion, attempted to stabilize sandy areas through planting vegetation, and combated soil erosion through reforestation. These are all examples of intentional human actions that have influenced the geomorphic processes and forms.

- Meanwhile, there are also numerous unintentional consequences of human activities on the Earth's surface. These include deforestation and burning grasslands for agricultural or urban development purposes, land subsidence due to mining activities and groundwater extraction, alteration of the landscape as a result of construction and mining, and increased soil erosion and sedimentation rates due to excessive and unplanned deforestation. Pollution, a significant byproduct of human economic activities, also plays a role in altering geomorphic processes and forms.
- Furthermore, the building of dams can lead to changes in river sediment loads and increased erosion, while construction in high-altitude regions can alter permafrost conditions. In summary, applied geomorphology seeks to understand the complex interplay between human activities and the Earth's surface, with a particular focus on both the intentional and unintentional consequences of these actions on the landscape and its processes.

What are some applications of geomorphology in engineering works? **A.** Road construction **B.** Dam site selection **C.** Airstrip construction **D.** All of the above

Specific Applications

1. **Geomorphology and Hydrology:** Geomorphology plays a crucial role in understanding the movement and distribution of water on Earth's surface. By studying the shape and structure of landscapes, hydrologists can predict and manage water flow, flood risks, and water resource availability.
2. **Geomorphology and Mineral Exploration:** The study of landforms and the processes that shape them can provide essential clues for locating valuable mineral deposits. Geomorphologists can help identify areas with high potential for mineral resources, reducing the time and cost of exploration.
3. **Geomorphology and Engineering Works:** Geomorphologic analysis can help engineers design safe and sustainable infrastructure projects by assessing the stability of slopes, the susceptibility of an area to erosion, and the potential impacts of construction on surrounding landscapes.
4. **Geomorphology and Military Geology:** Geomorphology can provide valuable information for military strategists, such as identifying potential obstacles, determining the optimal placement of defensive structures, and understanding how the terrain may affect troop movement and transportation.
5. **Geomorphology and Urbanization:** As cities continue to grow and expand, geomorphology can help planners make informed decisions about where to build and how to minimize the impacts of urbanization on the natural environment, such as preserving vital ecosystems and reducing the risk of natural hazards.
6. **Geomorphology and Hazard Management:** Geomorphologists can help identify areas at risk for natural hazards, such as landslides, earthquakes, and floods, and develop strategies for

mitigating these risks by understanding the underlying geological processes at work.

7. **Geomorphology and Regional Planning:** By analyzing the physical characteristics of a region, geomorphologists can provide valuable insights for regional planners, helping them make informed decisions about land use, resource management, and environmental protection.

In summary, geomorphology has a wide range of applications that can benefit professionals in various fields, from guiding mineral exploration efforts to informing urban planning decisions. By understanding the processes that shape Earth's landscapes, we can make more informed decisions about how to manage and protect our environment.

Geomorphology and Hydrology

- Water sources for human consumption can be derived from various sources such as streams, lakes, rivers on the Earth's surface, or groundwater. Different geological zones offer diverse conditions for both surface and groundwater availability.
- Limestone terrains have varying water-yielding capacities depending on the rock type. Permeability in limestones can be primary or secondary. Primary permeability depends on the existence of initial interconnected voids in the calcareous sediments from which the rock was formed. Secondary permeability arises from earth movements like faulting, folding, and warping, as well as solution or corrosion mechanisms.
- This secondary permeability is greatly influenced by a region's topography, being most significant beneath and adjacent to topographic lows or valleys. In karst terrain, much of the groundwater is confined to solution channels.
- In the early stages of karst evolution, conditions are not very different from those in other landscapes with similar relief. However, as the cycle progresses, a large portion of water is diverted to solutionally opened passageways, reducing surface water availability. The primary source of water in such regions then becomes karst springs. These springs can provide water to meet moderate demands, but the water quality may be affected by pollutants and bacteria.
- To determine the sources of spring water in cases of pollution, the swallow holes and sinkholes that feed water to the underground drainage systems emerging as springs must be located. This can be achieved by introducing a coloring material, like fluorescein, into the water entering nearby swallow holes (or sinkholes) and testing the various spring waters to identify their source. A knowledge of the region's structural geology is helpful in this context, as groundwater tends to move down rather than up the regional dip.
- The ease of obtaining water in a limestone region depends on the area's geomorphology. If the limestones have adequate permeability and are capped by a sandstone layer, large-yield wells may be easily obtained. Additionally, the water would be naturally filtered as it passes through the sandstone beds.
- However, if the limestone is dense and compact with low mass permeability, groundwater movement will primarily occur through secondary openings. In such circumstances, the water yield may be low, or even if sufficient, subject to contamination. Karst plains lack a filtering cover, and sinkholes, swallow holes, or karst valleys within an area of clastic rocks should raise concerns about the purity of nearby spring water.
- Groundwater potential in glaciated regions can be determined based on the area's geomorphic history, characteristics of glacial deposits, and landforms. Outwash plains, valley trains, and intertill gravels are likely to yield large volumes of water. Most tills are poor water sources due to the clay content, but they contain local strata of sand and gravel that may hold and supply enough water for domestic needs.
- Buried preglacial and interglacial valleys could be good sources of groundwater. Their presence (or absence) can be detected by studying the preglacial topography and geomorphic history of the area. Buried valleys are located by constructing bedrock topography maps of glaciated areas.

Geomorphology and Mineral Exploration

Mineral deposits are often associated with specific geological structures, and the landscape characteristics of certain areas can indicate the presence of these structures. There are several types of minerals and ore bodies that can be identified by their surface expressions or through the study of geomorphology.

- **Surface Expression of Ore Bodies:** Some ore bodies, such as lead-zinc lodes or quartz veins, have visible surface expressions in the form of topographic forms, outcrops, gossans, or structural features like faults, fractures, and breccia zones. These can help in identifying the presence of mineral deposits in the area.
- **Weathering Residues:** Minerals like iron ore, clay, caliche, bauxite, and some manganese and nickel ores are formed as weathering residues of present or ancient geomorphic cycles. These minerals are typically found on pleneplain or near-pleneplain surfaces, which are remnants of Tertiary erosional surfaces.
- **Placer Deposits:** Placer deposits are mixtures of heavy metals formed through chemical weathering or erosion of metallic formations. These deposits are formed due to specific geomorphic processes and can be found in particular topographical positions. Examples of minerals found in placer deposits include gold, tin, platinum, and diamonds.
- **Oil Exploration:** Many oil fields have been discovered due to their distinct topographic expression. Oil is usually trapped in rocks under structural or stratigraphic traps, with porous and permeable rocks like sandstone and limestone being ideal locations for oil deposits. Techniques such as drainage analysis using aerial photography can help identify areas with geological structures favorable for oil accumulation.

In summary, the study of geomorphology can be highly beneficial in mineral exploration. Knowledge of the bedrock geology, geophysical surveying, test drilling, and aerial-photograph interpretation are essential components in the search for mineral deposits. A thorough understanding of the geomorphic history of a region can significantly aid in the discovery of valuable mineral resources.

Geomorphology and Engineering Works

Engineering projects typically involve evaluating various geologic factors, with terrain characteristics being among the most common. Geomorphology, the study of landforms and their processes, plays a vital role in several engineering fields, including road construction, dam site selection, airstrip construction, locating sand and gravel pits, military geology, urbanization, hazard management, and regional planning.

- **Road Construction:** Determining the best highway routes depends on the topographic features of the area. Understanding the geologic structure, lithological and stratigraphic characteristics, strength of surface deposits, and geomorphic history of the area is crucial in road engineering. Different terrains, such as karst plains, glacial terrains, and areas with considerable relief, present unique engineering challenges. Modern highway construction also requires knowledge of soil profiles and their relationships to varying topographic conditions and types of parent material.
- **Dam Site Selection:** Selecting appropriate dam sites involves synthesizing knowledge of geomorphology, lithology, and geologic structure of terrains. Good reservoir sites must meet several geologic conditions, including a watertight basin of adequate size, a narrow outlet with a suitable foundation, an adequate spillway, availability of construction materials, and assurance of a long reservoir life. Understanding the subsurface topography is essential for constructing a dam that will not leak or become structurally compromised.
- **Airstrip Construction:** Constructing airstrips requires engineering skill and knowledge of landscape characteristics. Ideal airstrip sites have a flat surface with resistant materials, a nearly level slope, and good visibility. Geomorphological knowledge can be used to create a morphological map of the region where the airstrip will be built.
- **Locating Sand and Gravel Pits:** Sand and gravel have numerous engineering, commercial, and industrial uses. Selecting suitable sites for sand and gravel pits involves evaluating geologic factors such as variation in grade sizes, lithologic composition, degree of weathering, amount of overburden, and continuity of deposits. Geomorphological knowledge helps identify the type of deposit and its potential for development.
- **Military Geology:** Geomorphology plays a crucial role in military operations, as terrain appreciation and analysis are essential for proper planning. Geomorphologists can provide valuable insights into terrain conditions, influencing the interpretation of topographic maps and aerial photographs.
- **Urbanization:** Urban geomorphology, the study of landforms in relation to urban development and management, has become a

distinct branch of geomorphology. An urban geomorphologist can help in the planning and development process by providing detailed information on topography, soils, hydrology, lithology, and terrain characteristics.

- **Hazard Management:** Geomorphology is also crucial in identifying, predicting, and managing natural and man-made hazards. Geomorphic knowledge can be used to monitor seismic events, volcanic eruptions, floods, landslides, avalanches, and other hazards, as well as to assess their potential impacts and develop management strategies.
- **Regional Planning:** Applied geomorphology also plays a role in regional planning, as balanced economic growth requires an understanding of each region's resources, both natural and human. Detailed information on topography, soils, hydrology, lithology, and terrain characteristics can help regional planners devise development projects best suited for the region.

Other Applications

Besides the areas mentioned above, applied geomorphology is also useful in fields such as soil mapping, beach engineering, soil erosion control, and land classification. It can also help control the adverse effects of human activities on geomorphic forms and processes, leading to the development of anthropogeomorphology as a separate branch.

Techniques of Applied geomorphology

Applied geomorphology focuses on the relationship between human activities and geomorphological processes. Some key aspects of applied geomorphology include mapping landforms that affect or are modified by human activity, interpreting aerial photographs and remote sensing images, monitoring environmental changes, assessing the causes of unsustainable changes, and proposing solutions to hazards resulting from these changes.

- Aerial photographs and satellite images have made the preparation and interpretation of specialized maps more accurate and efficient. These images are useful for evaluating landforms and land use in relation to urban development plans, major construction projects, and more. Satellite images also play a vital role in studying global and country-level climatic phenomena, mineral prospecting, preparing land use inventories, and forecasting agricultural outputs.
- Remote sensing involves collecting information about objects from a distance without direct contact, using sensors and cameras to measure the spectral behavior of the objects under study. The most common remote sensing techniques involve sensing electromagnetic radiation emitted by terrestrial objects. Each object has a unique scattering property, or signature, based on its molecular composition. Knowledge of these signatures is crucial for interpreting satellite images.
- Remote sensing is essential for the sustainable management of natural resources such as soil, forests, crops, and oceans, as well as urban and town planning. Resource planners rely on remote sensing for up-to-date information on resource conditions and extent. Since these resources are dynamic and replenishable,

ground-based monitoring systems often cannot keep up with their changes. Satellite-based surveys, on the other hand, can provide more frequent and consistent data.

- Geographical Information Systems (GIS) technology is often used in conjunction with remote sensing techniques. GIS are spatial data-handling programs used to collect, store, and retrieve spatial data from the real world, containing only the data that researchers consider relevant.

Remote sensing surveys offer several advantages over ground surveys, including:

- A broader view, covering large areas at once.
- A permanent record of ground conditions that can be verified later.
- Faster data interpretation compared to ground surveys.
- Access to thermal and microwave regions not visible to the naked eye.
- Lower costs and less infrastructure required compared to ground surveys.
- More cost-effective for repeated surveys.
- Versatility, as the same data can be used for various purposes, such as soil, groundwater, or crop surveys.
- Unaffected by weather conditions or other obstacles that hinder ground surveys.

In summary, applied geomorphology is a vital field, utilizing aerial photographs, satellite images, and remote sensing techniques to study the relationship between human activities and geomorphological processes. This information is essential for sustainable resource management and urban planning.

How do remote sensing techniques benefit applied geomorphology?

A. They provide a broader view, covering large areas at once **B.** They offer a permanent record of ground conditions for later verification **C.** They allow access to thermal and microwave regions not visible to the naked eye **D.** All of the above

Conclusion

Applied geomorphology plays a critical role in various aspects of human interaction with the Earth's surface, from resource management and urban planning to hazard mitigation and infrastructure development. By understanding landforms and the processes that shape them, professionals in diverse fields can make more informed decisions, leading to sustainable development and better management of natural resources. The use of aerial photographs, satellite images, and remote sensing techniques has greatly enhanced the efficiency and accuracy of applied geomorphology, making it an indispensable tool for addressing the challenges posed by our ever-changing planet.

What is applied geomorphology, and why is it important?

Applied geomorphology is the process of using geomorphic knowledge to address issues related to land use, resource management, and environmental planning. It is essential because it helps assess resources, plan development projects, identify potential natural hazards, and reduce the impact of natural disasters, ultimately leading to more informed decision-making in various fields, such as urban planning and mineral exploration.

What are some specific applications of applied geomorphology?

Applied geomorphology can be applied in various fields, including hydrology, mineral exploration, engineering works, military geology, urbanization, hazard management, and regional planning. It can help professionals in these fields better understand and address challenges posed by different landscapes and environments.

How does applied geomorphology contribute to urban planning?

Applied geomorphology can help urban planners make informed decisions about where to build and how to minimize the impacts of urbanization on the natural environment. This includes preserving vital ecosystems, reducing the risk of natural hazards, and guiding land use planning and development in diverse geomorphic environments.

What role does remote sensing play in applied geomorphology?

Remote sensing involves using sensors and cameras to collect information about objects from a distance without direct contact. It is crucial for interpreting satellite images, monitoring environmental changes, and providing accurate, up-to-date information about resource conditions and extent. Remote sensing is essential for sustainable management of natural resources and urban planning, as it offers a broader view, faster data interpretation, and more cost-effective monitoring compared to ground-based surveys.

What is the relationship between GIS and applied geomorphology?

Geographical Information Systems (GIS) technology is often used in conjunction with remote sensing techniques in applied geomorphology. GIS are spatial data-handling programs that collect, store, and retrieve spatial data from the real world. They help researchers analyze and interpret data collected through remote sensing, providing valuable insights for managing and planning our interactions with the earth's surface.

5. What are some techniques used in applied geomorphology?



Ans. Some techniques used in applied geomorphology include aerial photography, satellite imagery, LiDAR (Light Detection and Ranging), topographic surveys, and geomorphological mapping. These techniques help in analyzing landforms, surface processes, and their spatial distribution, which are crucial for understanding and solving practical problems in various fields, such as geology, hydrology, environmental management, and engineering.

1. What is applied geomorphology?



Ans. Applied geomorphology is the study of landforms and the processes that shape them, with the purpose of understanding and solving practical problems. It involves the application of geomorphological knowledge and techniques to real-world issues and challenges.

2. What are the two main lines of application of geomorphology?



Ans. The two main lines of application of geomorphology are: - Geomorphology and Mineral Exploration: Geomorphological studies are used to identify potential mineral deposits by analyzing the landforms and surface processes that are indicative of mineralization. - Geomorphology and Engineering Works: Geomorphological knowledge is applied in engineering projects to assess and mitigate natural hazards, such as landslides, river erosion, and coastal erosion.

3. How is geomorphology applied in mineral exploration?



Ans. Geomorphology is applied in mineral exploration by examining the landforms and surface processes associated with mineralization. Certain landforms, such as alluvial fans or glacial moraines, can indicate the presence of mineral deposits. By studying the geomorphological characteristics of an area, geologists can identify potential locations for mineral exploration.

4. In what ways is geomorphology used in engineering works?



Ans. Geomorphology is used in engineering works to assess and mitigate natural hazards. By studying the landforms and surface processes, engineers can identify areas prone to landslides, river erosion, or coastal erosion. This knowledge helps in designing and implementing appropriate measures to prevent or reduce the impact of these hazards on engineering structures and human settlements.

Development of Modern Landforms

Fluvial Landforms

- Fluvial landforms are geological features created by the action of running water, primarily rivers. The term "fluvial"

originates from the Latin word "fluvius," which means river. These landforms vary greatly in size, ranging from small features like rills to massive continental-scale units such as large rivers and their drainage basins.

- Approximately 68% of the Earth's land surface is drained by rivers flowing into the oceans. Rivers typically originate in elevated areas with a downward slope, allowing water to run off. These highland regions serve as the catchment areas for rivers, with mountain crests acting as divides or watersheds from which the streams flow.
- The initial stream, called the consequent stream, forms due to the downward slope. As this stream erodes the surface, it is joined by tributaries from either side. The drainage basin, or watershed, is a crucial component of fluvial geomorphology, encompassing the primary river and its tributaries.

Fluvial erosion occurs in various ways, including:

- **Hydration:** The force of running water wears down rocks.
- **Corrosion/Solution:** Chemical reactions cause weathering.
- **Attrition:** The transported material experiences wear and tear as it rolls and collides with other materials.
- **Corrasion or abrasion:** The solid materials carried by the river strike against rocks, wearing them down.
- **Downcutting (vertical erosion):** The base of a stream is eroded, leading to valley deepening.
- **Lateral erosion:** The walls of a stream are eroded, leading to valley widening.
- **Headward erosion:** Erosion at a stream's origin causes it to move backward, lengthening the stream channel.
- **Hydraulic Action:** River water mechanically loosens and sweeps away materials, primarily by surging into crevices and cracks in rocks and breaking them apart.

Braiding is another fluvial process where the main water channel divides into multiple, narrower channels. Braided rivers consist of a network of channels separated by small, temporary islands called braid bars. This phenomenon occurs in rivers with a low slope and/or a large sediment load.

Which of the following erosional processes involves water mechanically loosening and sweeping away materials by surging into crevices and cracks in rocks and breaking them apart? **A.** Hydration **B.** Corrasion **C.** Hydraulic Action **D.** Attrition

- Streams are few during this stage with poor integration and flow over original slopes
- The valley developed is thus deep, narrow and distinctly V-shaped with no floodplains or with very narrow floodplains.
- Downcutting predominates over lateral corrasion
- Streams divides are broad and flat with marshes, swamp and lakes.
- Some of the outstanding features which are developed in this stage are gorges, canyons waterfalls, rapids and river capture etc.

Mature



- During this stage, streams are plenty with good integration.
- Lateral corrasion tends to replace vertical corrasion
- The valleys are still V-shaped but wide and deep due to an active erosion of the banks;
- Trunk streams are broad enough to have wider floodplains within which streams may flow in meanders confined within the valley.
- Swamps and marshes of youth stage, as well as flat and broad inter-stream areas, disappear. The stream divides turn sharp.
- Waterfalls and rapids disappear.
- Meander and slip off slopes are the characteristic features of this stage

Old

- The river moving downstream across a broad level plain is heavy with sediments.
- Vertical corrasion almost ceases in this stage though lateral corrasion still goes on to erode its banks further
- Smaller tributaries during old age are few with gentle gradients.
- Streams meander freely over vast floodplains. Divides are broad and flat with lakes, swamps and marshes.
- Depositional features predominate in this stage
- Most of the landscape is at or slightly above sea level
- Characteristic features of this stage are floodplains, oxbow lakes, natural levees and Delta etc.

River Course

Youth



Splash Erosion

- Splash erosion or rain drop impact represents the first stage in the erosion process. Splash erosion **results from the bombardment of the soil surface by rain drops**.
- Rain drops behave as little bombs when falling on exposed or bare soil, displacing soil particles and destroying soil structure.

Sheet Erosion

- Sheet erosion occurs as a **shallow 'sheet' of water flowing over the ground surface**, resulting in the removal of a uniform layer of soil from the soil surface.
- Water moving fairly uniformly with a similar thickness over a surface** is called sheet flow, and is the cause of sheet erosion.

Rills, and Gully

- A rill is a **shallow channel in some soil**, created by the erosion of flowing water. Rills can generally be easily removed by tilling the soil. When rills get large enough that they cannot easily be removed, they're known as gullies.



Nala or Rivulet

- A rivulet is a small stream.

Ravine

- A ravine is a landform that is narrower than a canyon and is often the product of streambank erosion.
- Ravines are typically classified as larger in scale than gullies, although smaller than valleys.

River Valleys

- The extended depression on the ground through which a stream flows throughout its course is called a river valley.
- At different stages of the erosional cycle, the valley acquires different profiles
- Valleys start as small and narrow rills
- The rills will gradually develop into long and wide gullies
- The gullies will further deepen, widen and lengthen to give rise to valleys.
- Depending upon dimensions, shape, types and structure of rocks in which they are formed, many types of valleys like the V-shaped valley, gorge, canyon, etc. can be recognised.

I-shaped valley/Gorge

- When the sides of the valley are almost parallel to each other, they form an 'I' shape and hence, these valleys are known as I-shaped valley.
- A gorge is a deep and narrow valley with very steep to straight sides
- A gorge is almost equal in width at its top as well as its bottom.
- Gorges are formed in hard rocks.
- Example- Indus Gorge in Kashmir



Canyon


- A canyon is a variant of the gorge.
- Unlike Gorge, a canyon is wider at its top than at its bottom.
- A canyon is characterised by steep step-like side slopes

- Canyons commonly form in horizontal bedded sedimentary rocks
- Example Grand Canyon carved by Colorado River, USA




V-shape valley

- The river is very swift as it descends the steep slope, and the predominant action of the river is vertical corrosion
- The valley developed is thus deep, narrow and distinctly V-shaped



Interlocking Spurs


- Interlocking spurs are **projections of high land that alternate from either side of a V-shaped valley.**



Waterfalls & Rapid

- When rivers plunge down in a sudden fall of some height, they are called waterfalls
- Their great force usually wears out a plunge pool beneath

- Waterfalls are formed because of several factors like the relative resistance of rocks lying across the river, the relative difference in topographic reliefs e.g. in Plateau etc.
- A rapid is similarly formed due to an abrupt change in gradient of a river due to variation in resistance of hard and soft rocks traversed by a river
- Waterfalls are also transitory like any other landform and will recede gradually and bring the floor of the valley above waterfalls to the level below.



Potholes & Plunge Pool

- Potholes are more or less circular depressions formed over the rocky beds of hill-streams, because of stream erosion aided by the abrasion of rock fragments.
- Once a small and shallow depression forms, pebbles and boulders get collected in those depressions and get rotated by flowing water and consequently the depressions grow in dimensions.
- Eventually, such depressions are joined leading to deepening of the stream valley.
- At the foot of waterfalls also, large potholes, quite deep and wide, form because of the sheer impact of water and rotation of boulders. These deep and large holes at the base of waterfalls are referred to as plunge pools.
- These pools also help in the deepening of valleys



Cataract

- Cataracts are **waterfalls on very large rivers.**
- The term cataract is usually applied to that section of a rapidly flowing river where the running water falls

suddenly in a sheer drop. When the drop is less steep, the fall is known as a **cascade**.

- A cataract is a powerful, even dangerous, waterfall.




Ait/Eyot

- An ait or eyot is a small island.
- It is especially used to refer to river islands found on the River Thames and its tributaries in England.
- Aits are typically formed by the deposit of sediment in the water, which accumulates over a period of time.
- An ait is characteristically long and narrow, and may become a permanent island should it become secured and protected by growing vegetation.
- However, aits may also be eroded: the resulting sediment is deposited further downstream and could result in another ait. A channel with numerous aits is called a braided channel.

Incised or Entrenched Meanders

- Incised or entrenched meanders are found cut in hard rocks. They are very deep and wide.
- In streams that flow rapidly over steep gradients, normally erosion is concentrated on the bottom of the stream channel.
- Entrenched meander normally occurs where there is a rapid cutting of the river bed such that the river does not get to erode the lateral sides.
- Meander loops are developed over original gentle surfaces in the initial stages of development of streams and the same loops get entrenched into the rocks normally due to erosion or gradual uplift of the land over which they started.
- They are widened and deepened over a long period of time and can be found as deep gorges and canyons in the areas where hard rocks are found.
- They give an indication of the status of original land surfaces over which streams have developed.
- Incised meanders are said to be an impact of river rejuvenation.




Structural Benches

- Step like sequence of geomorphic surfaces
- Differential erosion of alternately arranged hard and soft rocks forming step-like valleys known as structural benches
- The benches formed due to differential erosion of alternate bands of hard and soft rock beds are called structural benches or terraces because of lithological control in the rate of erosion and consequent development of benches.

River Terraces

- River terraces refer to surfaces relating to old valley floor or floodplain levels.
- They may be bedrock surfaces without any alluvial cover or alluvial terraces consisting of stream deposits.
- River terraces are basically products of erosion as they result due to vertical erosion by the stream into its own depositional floodplain.
- There can be a number of such terraces. They are found at different heights indicating former river bed levels.
- The river terraces may occur at the same elevation on either side of the rivers in which case they are called paired terraces




Peneplain

- A peneplain (an almost plain) is a low-relief plain which is formed as a result of stream erosion

- The peneplain is meant to imply the representation of a near-final (or penultimate) stage of fluvial erosion during times of extended tectonic stability.

Drainage Basin

- Other terms that are used to describe drainage basins are **catchment, catchment area, catchment basin, drainage area, river basin, and water basin.**
- The drainage basin includes both the streams and rivers and the land surface.
- The drainage basin acts as a funnel by collecting all the water within the area covered by the basin and channelling it to a single point.
- In **closed (endorheic) drainage** basins the water converges to a single point inside the basin, known as a sink, which may be a permanent lake (e.g. Lake Aral, also known Aral Sea, Dead Sea), dry lake (some desert lakes like Lake Chad, Africa), or a point where surface water is lost underground (sinkholes in Karst landforms).



Drainage Divide

- Adjacent drainage basins are separated from one another by a drainage divide.
- Drainage divide is usually a ridge or a high platform.
- Drainage divide is conspicuous in case of youthful topography (Himalayas), and it is not well marked in plains and senile topography.

Drainage Patterns

- The drainage pattern of a stream refers to the typical shape of a river course as it completes its erosional cycle
- They are governed by the topography of the land, resistance and strength of base rocks and the gradient of the land

There are various types of drainage patterns which are described briefly as below:

Dendritic Drainage Pattern

- It is the most common form of drainage system.
- The drainage pattern resembling the branches of a tree is known as dendritic. In a dendritic system, there are many contributing streams, which are then joined together into the tributaries of the main river
- The examples of Dendritic Pattern include the rivers of northern plain such Indus.

Trellis Drainage Pattern

- In the trellis drainage pattern, the primary tributaries of rivers flow parallel to each other and they are joined by secondary tributaries at the right angle.
- The geometry of a trellis drainage system is similar to that of a common garden trellis used to grow vines.
- Trellis drainage is characteristic of folded mountains,
- Examples of trellis pattern include the drainage system of the Appalachian Mountains in North America and Seine and its tributaries in Paris basin (France) etc.

Parallel Drainage Pattern

- A parallel drainage system is a pattern of rivers caused by steep slopes with some relief.
- The parallel drainage pattern is observed in a uniformly sloping region where the tributaries seem to be running parallel to each other.
- A parallel pattern sometimes indicates the presence of a major fault that cuts across an area of steeply folded bedrock.
- Examples of this system include the rivers of Lesser Himalaya

Rectangular Drainage Pattern

- Rectangular drainage develops on rocks that are of approximately uniform resistance to erosion, but which have two directions of joining at approximately right angles.
- In the rectangular drainage pattern, the mainstream curve at right angles and the tributaries join the mainstream at right angles.
- Example Colorado river the USA

Angular Drainage Pattern

- Angular drainage pattern is commonly observed in foothill regions.
- Angular drainage patterns form where bedrock joints and faults intersect at more acute angles than rectangular

drainage patterns. Angles are both more and less than 90 degrees

- the mainstream is joined by the tributaries at acute angles.

Radial Drainage Pattern

- When the rivers originate from a hill and flow in all directions, the drainage pattern is known as radial .
- Volcanoes usually display excellent radial drainage. Other geological features on which radial drainage commonly develops are domes and laccoliths.
- The rivers originating from the Amarkantak range present a good example of it.


Centripetal Drainage Pattern

- When the rivers discharge their waters from all directions in a lake or depression, the pattern is known as centripetal .
- Examples – streams of Ladakh, Tibet and Loktak Lake in Manipur (India)

Annular Drainage Pattern

- In an annular drainage pattern streams follow a roughly circular or concentric path along a belt of weak rock, resembling in plan a ring-like pattern.
- Example of such system include Black Hill streams of South Dakota, USA

Drainage patterns




Fluvial Landforms – Depositional

- Fluvial Depositional landforms are made by river sediments brought down by extensive erosion in the upper course of the rivers.
- Rocks and cliffs are continually weathered and eroded in the youth stage or upper course of the river.
- The river moving downstream on a level plain brings down a heavy load of sediments from the upper course.

- The decrease in stream velocity in the lower course of the river reduces the transporting power of the streams which leads to deposition of this sediment load.
- Coarser materials are dropped first and finer silt is carried down towards the mouth of the river
- This depositional process leads to the formation of various depositional landforms through fluvial action such as Delta, Levees and Flood Plain etc.

Alluvial Fans and Cones


- An alluvial fan is a cone-shaped depositional landform built up by streams, heavy with sediment load.
- Alluvial fans are formed when streams flowing from mountains break into foot slope plains of low gradient.
- Normally very coarse load is carried by streams flowing over mountain slopes. This load gets dumped as it becomes too heavy to be carried over gentler gradients by the streams
- Furthermore, this load spreads as a broad low to a high cone-shaped deposit called an alluvial fan that appears as a series of continuous fans.
- Alluvial fans in humid areas show normally low cones with a gentle slope from head to toe and they appear as high cones with a steep slope in arid and semi-arid climates.



Floodplains

- Floodplain is a major landform of river deposition.
- Deposition develops a floodplain just as erosion makes valleys.
- Rivers in the lower course carry large quantities of sediments
- Large sized materials are deposited first when stream channel breaks into a gentle slope.
- Sand, silt and clay and other fine sized sediments are carried over gentler channels by relatively slow-moving waters
- During annual or sporadic floods, these materials are spread over the low lying adjacent areas. A layer of sediments is thus deposited during each flood, gradually building up a floodplain

- In plains, channels shift laterally and change their courses occasionally leaving cut-off courses which get filled up gradually by relatively coarse deposits.
- The flood deposits of spilt waters carry relatively finer materials like silt and clay.
- **Active Floodplain** – A riverbed made of river deposits is the active floodplain.
- **Inactive Floodplain** – The floodplain above the bank is an inactive floodplain. Inactive floodplain above the banks basically contains two types of deposits flood deposits and channel deposits.
- **Delta plains** – The floodplains in a delta are called delta plains.




Doab

- Doab is the tract of land between two converging rivers.
- Doab is a term used in South Asia particularly in India and Pakistan to refer to "tongue" or tract of land lying between two converging rivers.


Natural Levees

- This is an important landform associated with floodplains.
- They are found along the banks of large rivers.
- They are low, linear and parallel ridges of coarse deposits along the banks of rivers on both sides due to deposition action of the stream, appearing as natural embankments.
- At the time of flooding, the water is spilt over the bank. As the speed of flow of the water comes down, large sized sediments with high specific gravity are dumped along the bank as ridges.
- They are high nearer the banks and slope gently away from the river.
- Generally, the levee deposits are coarser
- When rivers shift laterally, a series of natural levees can form.
- Artificial embankments are formed on the levees to minimize the risk of the floods.
- But sudden bursts in the banks due to the pressure of water can cause disastrous floods.
- An example of such flood can be seen in Hwang Ho river which is also called China's sorrow.



Point Bars & Cut Banks

- Point Bar is also associated with floodplain
- Point bars are also known as meander bars.
- A point bar is a depositional feature
- It is formed by alluvium that accumulates in a linear fashion on the inside bends of streams and rivers below the slip-off slope.
- They are found on the convex side of meanders of large rivers.
- They are almost uniform in profile and in width and contain mixed sizes of sediments.
- Long and narrow depressions can be found in between the point bars where there is more than one ridge
- Rivers build a series of them depending upon the water flow and supply of sediment.
- As the point bars are built by the rivers on the convex side, erosion takes place on the concave side of the bank.
- Cut banks are found on the outside of a bend in a river. Cut banks are caused by the moving water of the river wearing away the earth.




Meanders

- In large flood and delta plains, rivers rarely flow in straight courses. Loop-like channel patterns called meanders develop over flood and delta plains
- Normally, in meanders of large rivers, there is active deposition along the convex bank and undercutting along the concave bank.
- If there is no deposition and no erosion or undercutting, the tendency to meander is reduced.

- The concave bank is known as a **cut-off bank** which shows up as a steep scarp and the convex bank presents a long, gentle profile and is known as the **slip-off bank**.


Oxbow Lake

- In the lower course of a river, meanders become very much more pronounced
- As meanders grow into deep loops, the same may get cut-off due to erosion at the inflexion points and are left as independent water bodies, known as ox-bow lakes.
- Through subsequent floods that may silt up the lake, oxbow lakes are converted into swamps in due course of time. It becomes marshy and eventually dries up



Riffle and Pool

- Pools:** An area of the stream characterized by deep depths and slow current. Pools are typically created by the vertical force of water falling down over logs or boulders. The movement of the water carves a deeper indentation in the stream bed. Pools are important because they can provide depth and still water.
- Riffles:** An area of stream characterized by shallow depths with fast, turbulent water. The riffles are short segments of the stream where water flow is agitated by rocks. The rocky bottom provides protection from predators, food deposition, and shelter. Riffle depths vary depending upon stream size but can be as shallow as 1 inch or deep as 1 meter. The turbulence and streamflow result in high dissolved oxygen concentration.



Bluff

- A bluff is a small, rounded cliff that usually overlooks a body of water, or where a body of water once stood.
- Bluff is a ridge of land that extends into the air.

Braided Channels

- A braided channel consists of a network of river channels divided into multiple threads and separated by small and often temporary islands called eyots.
- Braided channels are commonly found where water velocity is low and the river is heavy with sediment load
- Deposition and lateral erosion of banks are essential for the formation of the braided pattern.
- There is the formation of central bars due to selective deposition of coarser material which diverts the flow towards the banks causing extensive lateral erosion
- As the valley widens due to continuous lateral erosion, the water column is reduced and more and more materials get deposited as islands and lateral bars developing a number of separate channels of water flow.



Delta

- Deltas are fan-shaped alluvial areas, resembling an alluvial fan
- This alluvial tract is, in fact, a seaward extension of the floodplain
- The load carried by the rivers is dumped and spread into the mouth of the river at sea. Further, this load spreads and piles up as a low cone
- Unlike in alluvial fans, the deposits making up deltas are very well sorted with clear stratification. The coarsest sediments are deposited first and the finer sediments are carried out further, into the sea.
- Deltas extend sideways and seaward at an amazing rate
- As the delta grows, the river distributaries continue to increase in length and Delta continues to build up into the sea.
- Some deltas are extremely large. For example, the Ganges delta is as big as the whole west of Malaysia



Types of Deltas: There are great variations in size, shape, growth, and importance of Deltas. A great number of factors influence the eventual formation of deltas such as depth of the river, sedimentation, sea-bed, character of tides, waves, and

currents, etc. owing to these factors several types of deltas can be found.

- **Bird's foot delta** It is a kind of delta featuring long, stretching distributary channels, which branch outwards resembling the foot of a bird. Deltas that are less subjected to wave or tidal action culminate to a bird's foot delta. Example the Mississippi River has a bird's foot delta extending into the Gulf of Mexico
- **Arcuate delta** Arcuate is the most common type of delta. This is a fan-shaped delta. It is a curved or bowed delta with a convex margin facing the sea. Arcuate deltas have a smooth coastline due to the action of the waves and the way they are formed. Examples – The Nile, Ganges, and Mekong river deltas
- **Cuspate delta** A few rivers have tooth-like projections at their mouth, known as the cuspate delta. Cuspate deltas are formed where the river flows into a stable water body (sea or ocean). The sediments brought down by the rivers collide with the waves. As a result, sediments are spread evenly on either side of its channel. Example Ebro river delta in Spain
- **Estuarine delta** some rivers have their deltas partly submerged in the coastal waters to form an estuarine delta. This may be due to a drowned valley because of a rise in sea level. Example Amazon river delta

Conditions Favorable for the Formation of Delta


- Active vertical and lateral erosion in the upper course of the river to provide extensive sediments to be eventually deposited as deltas.
- The coast should be sheltered preferably tideless.
- The sea adjoining the delta should be shallow or else the load will disappear in the deep waters.
- There should be no large lakes in the river to filter off the sediments.
- There should be no strong current running at right angles to the river mouth, washing away the sediments.



Types of Delta

Marine Landforms


- Waves are the most potent agents of marine erosion, primarily caused by winds sweeping over the water surface, generating a series of rolling swells that move forward. As these waves approach the shallow waters near the shoreline, their speed decreases, and they become curved and refracted according to the coast's alignment.
- When the shallow water is less than the height of the waves, their forward movement is impeded, causing the wave crest to curl over and break on the shore. The water that rushes up the beach and throws rock debris against the land is known as swash, while the water that retreats or is pulled back is referred to as backwash.
- Another component of offshore drift is an undertow, a current that flows near the bottom, away from the shore. This current exerts a pulling effect that can be hazardous for those swimming in the sea.



Types of Erosion


Erosion is the process of breaking down and carrying away rocks and sediments by natural forces such as wind, water, and ice. In coastal areas, several types of erosion are carried out by sea waves, winds, currents, tides, and storms. There are four main types of erosion:

1. **Corrasion or Abrasion:** This type of erosion occurs when waves carry rock debris of various sizes and shapes, which crash against the base of cliffs and wear them down. This process, known as corrasion, helps break down the cliffs and make them more susceptible to further erosion. Incoming tides and waves then complete the process by carrying away the eroded material into the sea.
2. **Attrition:** Attrition is the process by which waves transport beach materials such as boulders, pebbles, shingles, and fine sand, causing these materials to collide with each other and break into smaller pieces. This grinding and polishing of materials against the cliff faces and one another result in the formation of the fine sand that makes up beaches.
3. **Hydraulic Action:** Hydraulic action refers to the erosion that takes place when the movement of water against a rock surface results in mechanical weathering. It involves the ability of moving water, such as flowing currents or waves, to dislodge and transport rock particles. Hydraulic action is different from other water-related erosion processes, like static erosion, where water dissolves salts and carries away organic material from unconsolidated sediments, or chemical erosion, which is more commonly known as chemical weathering. Hydraulic action is a mechanical process in which the water current flows against the banks and bed of a river, removing rock particles as it goes.
4. **Solvent Action:** Solvent action is another form of erosion, though it mainly affects soluble rocks like limestone and chalk. In this process, the solvent or chemical action of seawater on calcium carbonate causes chemical changes in the rocks, leading to their disintegration. This type of erosion is most pronounced along limestone coasts, where the interaction between seawater and the calcium carbonate in the rocks leads to the breakdown and eventual erosion of the rock formations.



Marine Transportation


- **Solution** – minerals are dissolved in the water and carried along in solution.
- **Suspension** – fine light material is carried along in the water.
- **Saltation** – small pebbles and stones are bounced along the sea bed.
- **Traction** – large boulders and rocks are rolled along the sea bed.



Marine Landforms – Erosional

Headlands and Bays

- Coastal cliffs do not erode at a uniform rate, leading to the formation of headlands and bays when the coastline consists of varying rock types. Softer rocks like clay and sand are less resistant to erosion and tend to wear away more quickly, resulting in the creation of bays. A bay is a curved indentation in the coastline where the land slopes inward, typically featuring a beach.
- On the other hand, harder rocks such as chalk are more resistant to erosion. As the softer rocks erode, these harder rocks protrude into the sea, forming headlands. Erosional features like wave-cut platforms and cliffs are commonly found on headlands, as they are more exposed to the force of the waves. Bays tend to be more protected, with constructive waves depositing sediment to create a beach.





Cliffs and Wave-Cut platforms

- **Cliffs are shaped through erosion and weathering. Soft rock erodes quickly and forms gentle sloping cliffs, whereas hard rock is more resistant and forms steep cliffs.** A wavecut platform is a wide gently sloping surface found at the foot of a cliff.


A Wave-Cut Platform is Formed When the Following Occurs

- The ocean constantly assaults the cliff's foundation, specifically in the area between the high and low tide marks.
- A wave-cut notch, which is a small indentation in the cliff, forms as a result of erosion processes like abrasion and hydraulic action, typically occurring at high tide level.
- As this notch grows larger, the cliff's stability is compromised, leading to its eventual collapse and the retreat of the cliff face.
- The water then washes away the debris from the erosion, leaving behind a flat, wave-cut platform. This cycle continues, causing the cliff to progressively recede over time.



Caves, Arches, Stacks, and Stumps

- Caves, arches, stacks, and stumps are all examples of erosional features that can typically be found on a headland. These unique formations are created through a series of natural processes, mainly hydraulic action and abrasion, which help to widen existing cracks in the headland.
- Over time, waves persistently erode and grind against these cracks, causing them to expand and eventually develop into a cave. As the erosion continues and the cave grows larger, it may eventually break through the entire headland, resulting in the formation of an arch.
- The base of this arch is continually subjected to further erosion, causing it to widen until the weight of the arch's roof becomes too heavy to support itself. At this point, the roof collapses into the sea, leaving behind a stack, which is a tall, isolated column of rock.
- When a section of the sea arch collapses, the remaining vertical structure is referred to as a stack, skarry, or chimney rock. As the erosion process continues, the base of the stack is undercut, eventually causing it to collapse and form a stump.



Ria, and Cove

- A ria is a type of coastal inlet created when a river valley that has not been glaciated becomes partially submerged. This results in a drowned river valley that remains connected to the sea.
- On the other hand, a cove is a small, sheltered bay or inlet that typically has a narrow or restricted entrance. Coves can also be found at the opening of a creek or another small body of water. They are typically formed through the erosion of soft rock formations, leaving behind more durable rock that shapes a circular or oval bay with a limited entrance.
- Coves are generally smaller in size, with a width of less than 1000 feet, and can be even smaller, with some measuring less than 100 feet in diameter.

Geos and Gloops (blowholes or marine geyser)

- The intermittent splashing of waves on a cave's roof can cause the joints to expand as air is repeatedly compressed and released within them. This process can eventually create a natural shaft that might break through the surface. When waves crash into the cave, they can force water or air out of this hole, which is referred to as a Gloop or blowhole.
- As blowholes continue to enlarge and waves persistently impact the cave, the roof becomes weaker. If the roof eventually collapses, a long, narrow creek called a Geos may form.

Chasms


- These are **narrow, deep indentations (a deep recess or notch on the edge or surface of something) carved due to headward erosion (downcutting) through vertical planes of weakness in the rocks by wave action.**
- With time, further headward erosion is hindered by lateral erosion of chasm mouth, which itself keeps widening till a bay is formed.

Creek, and Inlet

- **Creek: natural stream of water normally smaller than and often tributary to a river.** creek is a narrow, sheltered waterway, especially an inlet in a shoreline or channel in a marsh.
- Inlet: An inlet is an **indentation of a shoreline, usually long and narrow, such as a small bay or arm**, that often leads to an enclosed body of salt water, such as a sound, bay, lagoon, or marsh.

Marine Landforms – Depositional

When water loses its energy, any sediment it is carrying is deposited. The build-up of deposited sediment can form different features along the coast.




Beaches

- Beaches consist of eroded materials that have been carried from other locations and deposited by the sea. This process requires waves with limited energy, which is why beaches frequently form in sheltered areas like bays. Constructive waves, characterized by a strong swash and a weak backwash, contribute to the formation of beaches.
- Sandy beaches are commonly found in bays with shallow water and less energetic waves. In contrast, pebble beaches tend to develop in areas with cliff erosion and higher energy waves. The cross-section of a beach is referred to as the beach profile, which consists of multiple ridges called berms. These berms indicate the lines of high tide and storm tides.
- Sandy beaches usually have a gentle slope in their profile, while shingle beaches can be much steeper. At the top of the beach, the material is larger due to the high-energy storm waves carrying larger sediment. The smallest material is found closest to the water, where the waves break and gradually wear down the rock through a process called attrition.


Spits and Hook

- A **spit is an extended stretch of sand or shingle jutting out into the sea from the land.** Spits occur when there is a change in the shape of the landscape or there is a river mouth.
- **This is how spits are formed:**
 - Sediment is carried by longshore drift.
 - When there is a change in the shape of the coastline, deposition occurs. A long thin ridge of material is deposited. This is the spit.
 - A hooked end can form if there is a change in wind direction.
 - Waves cannot get past a spit, therefore the water behind a spit is very sheltered. Silts are deposited here to form salt marshes or mudflats.




Bars, Lagoons and Barrier

- Sometimes a **spit can grow across a bay and join two headlands together. This landform is known as a bar.** They can trap shallow lakes behind the bar, these are known as **lagoons**. Lagoons do not last forever and may be filled up with sediment.
- **Barrier:** It is the **overwater counterpart of a bar.**



Tombolos & Dumb Ball

- A tombolo is a **sandy isthmus**. Sometimes, islands are connected **with mainland** by a bar called **tombolo**.
- If **two islands** are connected **to each other** by a bar is called **Dumb Ball**.



Marine Dunes & Dune Belts


- With the force of on-shore winds, a **large amount of coastal sand is driven landwards forming extensive marine dunes that stretch into dune belts.**
- Their advance inland may engulf farms, roads & even the entire villages;
- Hence to arrest the migration of dunes, sand binding species of grass & shrubs, such as marram grass & pines are planted.

Karst Landforms

- Karst topography is a landscape formed by the dissolution of soluble rocks, such as limestone, dolomite, and gypsum. This type of terrain is characterized by underground drainage systems, sinkholes, and caves. It is most prominently developed in dense carbonate rock, like limestone, which is thin-layered and highly fractured.
- Karst topography is not commonly found in chalk, as chalk is highly porous rather than dense. This means that groundwater flow is not concentrated along fractures. Karst landscapes are also more likely to develop in areas with low water tables, such as uplands with entrenched

valleys, and where there is moderate to heavy rainfall. This encourages the rapid downward movement of groundwater, promoting the dissolution of bedrock. In contrast, stagnant groundwater becomes saturated with carbonate minerals, preventing further dissolution of the bedrock.

- Karst landscapes can be found in various parts of the world, including the Causses in France; the Kwangsi region of China; the Yucatán Peninsula; and the Middle West, Kentucky, and Florida in the United States. In India, karst topography is present in the Vindhya region (primarily southwestern Bihar), the Himalayas (parts of Jammu & Kashmir, Robert Cave, Sahasradhara, the eastern Himalayas, and areas near Dehradun), Pachmarhi in Madhya Pradesh, Gupt Godavari Cave in Chitrakoot (U.P.), the surrounding coast near Vishakhapatnam (Borra Caves), and Bastar in Chhattisgarh.
 - Borra Caves, also known as Borra Guhalu, are situated on the East Coast of India, within the Ananthagiri hills of the Araku Valley.




Characteristics of Karst Landforms

- Karst regions are characterized by a stark landscape, with occasional steep slopes. These areas typically lack surface drainage, as most of the water seeps underground, leaving surface valleys dry. Streams tend to carve their paths along the joints and fissures in the rock, creating an intricate system of underground channels.
- When water reaches the base of the limestone and encounters non-porous rocks, it resurfaces as a spring or resurgence. Limestone is heavily jointed, and rainwater seeps into the rock through these joints and cracks. Over time, the process of dissolution widens these cracks, eventually forming unique features known as limestone pavements. The widened joints are referred to as grikes, while the isolated, rectangular blocks are called clints.

- Swallow holes, also known as sinkholes, are small depressions on the limestone surface, created by the dissolution process as rainwater sinks into the limestone at weak points. Water that seeps into limestone carves out caverns and passages along the joints. When multiple swallow holes merge, they form a larger hollow known as a doline. The gradual caving or subsidence of several dolines can result in an even larger depression called an uvala. In some instances, such as in Yugoslavia, large depressions called poljes may form, covering up to 100 square miles and being partially caused by faulting.
- Underground streams descending through swallow holes lead to the formation of caves and caverns, which may contain ponds or lakes. The most striking features of

limestone caves are stalactites, stalagmites, and calcite pillars. Water carries dissolved calcium, and when this water evaporates, it leaves behind solidified crystalline calcium carbonate.

- Stalactites are sharp, slender, downward-growing pinnacles that hang from cave ceilings. As moisture drips from the ceiling, it flows down the stalactites and falls to the floor, depositing calcium and forming stalagmites, which are shorter, wider, and more rounded. Over an extended period, the stalactite hanging from the ceiling eventually connects with the stalagmite growing from the floor, creating a pillar.



What is the primary difference between a sandy beach and a shingle beach? **A.** Sandy beaches have a steeper slope in their profile. **B.** Shingle beaches are found in areas with cliff erosion and higher energy waves. **C.** Sandy beaches usually form in bays with deep water and more energetic waves. **D.** Shingle beaches consist of eroded materials carried from other locations and deposited by the sea.

Erosional landforms of Karst topography

Blind Valley

- A **steephead valley, steephead or blind valley** is a deep, narrow, flat bottomed valley with an abrupt ending.
- Karst valley **abruptly terminated by the passage underground of the watercourse** which has hitherto resisted the karst processes and remained at the surface.

Swallow Hole/Sinkholes/Doline

- A sinkhole, also known as a **cenote, sink, sink-hole, swallet, swallow hole, or doline** (the different terms for sinkholes are often used interchangeably), is a **depression or hole in the ground caused by some form of collapse of the surface layer.**
- Most are caused by karst processes – for example, the chemical dissolution of carbonate rocks or suffosion processes
- The surface streams which sink disappear underground through swallow holes.
- **Local names:**
 - Black hole – Sea water
 - Blue hole- deep under water
 - Cenotes- Belize (British Honduras)
 - Sotanos- Mexico
 - Tomo- New Zealand

Clift

- When the Solution hole is dippen over a period of time then the dipped part is called Clift.

Pinnacles

- Vertical rock blades fretted sharpened by dissolution.

Lapies/Karren

- It is formed due to differential solution activity along parallel to sub-parallel joints. They are also called grooved, fluted and ridge-like features in an open limestone field.
- The most widespread surface karst landforms are small solution pits, grooves and runnels, collectively called Karren.

Limestone Pavements

- It is a smoother form of lapies.

Sinking Creeks/Bogas

- In a valley, the water often gets lost through cracks and fissures in the bed. These are called sinking creeks, and if their tops are open, they are called bogas.

Karst Window/Fenster


- Karst fenster is a geomorphic feature formed from the dissolution of carbonate bedrock.
- In this feature, a spring emerges, then the discharge abruptly disappears into a sinkhole.
- A karst fenster is caused by a caving in of portions of the roof of a subterranean stream, thus making some of the underground stream visible from the surface.
- When a number of adjoining sink holes collapse, they form an open, broad area called a karst window.

Uvalas

- Karst depressions that are much larger than sinkholes and that display gentler slopes and more complex three-dimensional shapes are known as uvalas.
- Uvalas is collection of multiple smaller individual sinkholes that coalesce into a compound sinkhole.
- A single uvala typically contains numerous sinkholes within it.

Polje

- A polje, also karst polje or karst field, is a large flat plain found in karstic geological regions of the world, with areas usually 5 to 400 km².



Pools

- An opening at the top with water collected in the void of the surface with varying depth


Caves/Cavern

- This is an underground cave formed by water action by various methods in a limestone or chalk area.
- Cave formation is prominent in areas where there are alternating beds of rocks (shales, sandstones, quartzites) with limestones or dolomites in between or in areas where limestones are dense, massive and occurring as thick beds



Karst Lake

- Karst lakes are formed as the result of a collapse of subterranean caves, especially in water soluble rocks.
- such as limestone, gypsum, and dolomite.
- This process is known as karstification. They can cover areas of several 100 square kilometers.
- Their shallow lakebed is usually an insoluble layer of sediment so that water is impounded, leading to the formation of lakes. Many karst lakes only exist periodically but return regularly after heavy rainfall.



Depositional landforms of Karst

topography

Stalactites & Helictite

- The water containing limestone in solution, seeps through the roof in the form of a continuous chain of drops.
- A portion of the roof hangs on the roof and on evaporation of water, a small deposit of limestone is left behind contributing to the formation of a stalactite, growing downwards from the roof.
- Usually, the base is broader than the free end of the hanging stalactites.
- The ones that descend vertically are known as **stalactites**, whereas the ones that extend horizontally or diagonally are known as **helictites**.

Stalagmites & Halagmite

- A stalagmite is a type of rock formation that rises from the floor of a cave due to the accumulation of material deposited on the floor from the ceiling drippings.
- It is an **upward-growing mound** of mineral deposits that have precipitated from water dripping onto the floor of a cave.
- Ones that extend horizontally or diagonally from stalagmites are known as Halagmite.

Cave Pillars

- The combination or fusion of stalactites and stalagmites form the pillars
- The diameters of pillars vary

Drapes/Curtain

- Numerous needle-shaped dripstones hanging from the cave ceiling are called Drapes or Curtains.

Tufa

- Tufa is a **variety of limestone formed when carbonate minerals precipitate out of ambient temperature water**.
- Geothermally heated hot springs sometimes produce similar (but less porous) carbonate deposits, which are known as travertine.

Travertine

- Travertine is a sedimentary rock formed by the chemical precipitation of calcium carbonate minerals from freshwater, typically in springs, rivers, and lakes that is, from surface and ground waters.
- Developed form of Tufa is Travertine.

Drip Stone


- **Calcium carbonate rock deposited in caves by the precipitation of calcite from water as excess dissolved carbon dioxide** is diffused into the atmosphere.
- Dripstone takes various forms, including stalactites, helictites (having spiral form), curtains, ribbons, and stalagmites.

Terra Rossa

- Terra Rossa is a well-drained, reddish, clayey to silty clayey soil with neutral pH conditions.

Human Activities of Karst Region


- Karst regions are **often barren & at best carry a thin layer of soil**.
- The **porosity of the rocks & the absence of surface drainage make vegetative growth difficult**, hence limestone can usually **support only poor grass**.
- Limestone vegetation in tropical regions is luxuriant because of heavy rainfall all the year around.
- The only mineral found in association of limestones is **lead**.
- **Good quality limestone is used as building materials & quarried for cement industry**.



Glacial Landforms

Glaciers

- A glacier is a **huge mass of ice that moves slowly over land**. The term “glacier” comes from the French word glace (glah-SAY), which means ice. **Glaciers are often called “rivers of ice.”**
- Glaciers normally assume the shape of a tongue, broadest at the source & becoming narrower downhill.
- Though glacier is not liquid, but **it moves gradually under the continual pressure from the snow accumulated above**.
- The **rate of movement is greatest in the middle** where there is little obstruction.
- The sides & bottom are held back by the frictions due to valley sides & valley floors.
- If a row of stakes is planted across a glacier in a straight line, they will eventually take a curved shape down the valley, showing that the glacier moves faster at the center than at the sides.
- **Glacial Landforms can be found in locations that currently have no active glaciers or glaciation processes.**
- Glaciers fall into two groups: **alpine glaciers and ice sheets**
- **Alpine glaciers**
 - Alpine glaciers **form on mountainsides and move downward through valleys**.
 - Sometimes, **alpine glaciers create or deepen valleys** by pushing dirt, soil, and other materials out of their way.
 - Alpine glaciers are **found in high mountains** of every continent except Australia (although there are many in New Zealand).
 - The Gorner Glacier in Switzerland and the Furtwangler Glacier in Tanzania are both typical alpine glaciers.
 - **Alpine glaciers are also called valley glaciers or mountain glaciers.**
- **Ice sheets**
 - Ice sheets, unlike alpine glaciers, **are not limited to mountainous areas**. They form **broad domes and spread out from their centers in all directions**.
 - As ice sheets spread, **they cover everything around them with a thick blanket of ice, including valleys, plains, and even entire mountains**.
 - The **largest ice sheets, called continental glaciers**, spread over vast areas. Today, continental glaciers cover most of Antarctica and the island of Greenland.
 - **Massive ice sheets covered much of North America and Europe during the Pleistocene time period**. This was the last glacial period, also known as the Ice Age. Ice sheets reached their greatest size about 18,000 years ago. As the ancient glaciers spread, they carved and changed the Earth's surface, creating many of the landscapes that exist today.
 - **During the Pleistocene Ice Age, nearly one-third of the Earth's land was covered by glaciers. Today, about one-tenth of the Earth's land is covered by glacial ice.**



The Ice-Age and Ice Masses

- An ice age is a **long period of reduction in the temperature of Earth's surface and atmosphere**, resulting in the presence or expansion of continental and polar ice sheets and alpine glaciers.
- Earth's climate alternates between ice ages and greenhouse periods, during which there are no glaciers on the planet.
- Earth is currently in the Quaternary glaciation.

- Today, **only 2 major ice caps are present in this world – Antarctica & Greenland**, along with many highlands above the snowline surviving in the world.
- Ice from the ice cap creeps out in all directions to escape as glaciers.
- When the ice sheets reaches down to the sea they float as ice shelves in polar waters.
- When ice sheets break into individual blocks, these are called icebergs.
- **While afloat in the sea, only 1/9th of the iceberg's mass is visible above the surface.**
- They diminish in size when reaching warm waters & eventually melted, dropping the rock debris that was frozen inside them on the sea bed.
- Permanent snowfield is sustained by heavy snowfall in winters & ineffective snow melting & evaporation in summers as part of snow that melt during the day is refrozen during the night.
- This refreezing process repeats until it forms a hard, granular substance known as neve or firn.
- Owing to the gravitational forces, neve of the upland snowfield is drawn towards the valley below, which marks the **beginning of the flow of glacier (river of ice)**.

Glacial Cycle of Erosion


- **Youth:** The stage is marked by the inward cutting activity of ice in a cirque. Aretes and horns are emerging. The hanging valleys are not prominent at this stage.
- **Maturity:** The valley glacier gets transformed into a trunk glacier and hanging valleys start emerging. The opposite cirques come closer and the glacial trough acquires a stepped profile that is regular and graded.
- **Old Age:** The emergence of a 'U'-shaped valley marks the beginning of old age. An outwash plain with features such as eskers, kame terraces, drumlins, kettle holes, etc. is a prominent development. The opposite cirques coalesce and the summit heights are greatly reduced. Mountain tops become rounded.

Glacial Landforms

- **Glaciations generally gives rise to erosional features in the highlands & depositional features on lowlands.**
- It erodes its valley by two processes viz. **plucking & abrasion**.
- **Plucking:** Glacier freezes the joints & beds of underlying rocks, tears out individual blocks & drags them away.

- **Abrasion:** Glacier scratches, scrapes, polishes & scours the valley floor with the debris frozen into it.

Glacial Landforms – Erosional




Snout or Glacier Terminus

- A **glacier terminus, toe, or snout**, is the **end of a glacier at any given point in time**.
- The terminus is the **usually the lowest end** of glacier.

Corrie, Cirque or cwm

- The downslope movement of a glacier from its snow-covered valley head & the the intensive shattering of the upland slopes, tend to produce a depression where neve or firn accumulates.
- Plucking & abrasion further deepen the depression into a steep horse shoe-shaped basin called Cirque (in French), cwm (in wales) & Corrie (in Scotland)
- There is a rocky ridge at the exit of the corrie & when the ice eventually melts, water collect behind this barrier known as Corrie Lake or tarn.



Cols

- Cols form when two cirque basins on opposite sides of the mountain erode the arête dividing them.
- Cols create saddles or passes over the mountain.

Horns


- Horns are a single pyramidal peak formed when the summit is eroded by cirque basins on all sides.

Aretes and Pyramidal Peaks

- When two corries cut back on opposite sides of the mountain, knife edged ridges are formed called arêtes
- When three or more cirques cut back together, recession will form an angular horn or pyramidal peak.


Bergschrund

- At the head of a glacier, where it begins to leave the snowfield of a corrie, a deep vertical crack opens up called a Bergschrund or Rimaye
- This happens in summer when although the ice continues to move out of the corrie, there is no new snow to replace it
- In some cases, not one but several such cracks occur which present a major obstacle to climbers
- Further down, where the glacier negotiates a bend or a precipitous slope, more crevasses or cracks are formed.




Roche Moutonnée

- Basically, a resistant residual rock hummock or mound, striated by the ice movement.
- Its upstream or stoss side is smoothed by abrasion & its downward or leeward side is roughened by plucking & is much steeper.
- It is believed that plucking may have occurred on the leeward side due to a reduction in pressure of the glacier moving over the stoss slope
- Therefore providing the opportunity for water to refreeze on the lee side and pluck the rock away.



Crag and Tail

- A crag and tail is a larger rock mass than a Roche moutonnée
- Like a Roche moutonnée, it is formed from a section of rock that was more resistant than its surroundings.
- Crag is a mass of hard rock with a steep slope on the upward side, which protects the softer leeward slope from being completely worn down by the oncoming ice.
- It therefore has a gentle tail strewn with the eroded rock debris.



Nunatak

- A nunatak is the summit or ridge of a mountain that protrudes from an ice field or glacier that otherwise covers most of the mountain or ridge. They are also called glacial islands.
- **Nunatak is a mountain peak or other rock formation that is exposed above a glacier or ice sheet.**

Paternoster Lakes


- A paternoster lake is one of a series of glacial lakes connected by a single stream or a braided stream system.
- Paternoster lakes are formed in the low depression of a U-shaped valley.
- Paternoster lakes are created by recessional moraines, or rock dams, that are formed by the advance and subsequent upstream retreat and melting of the ice.

U shaped glacial Troughs & Ribbon lakes

- Glaciers on their downward journey are fed by several corries & grind the bedrock with straightening out any protruding spurs.
- The interlocking spurs are thus blunted to form truncated spurs with the floor of the valley deepened.
- Hence, the valley which has been glaciated takes the characteristic of U shape, with a wide flat floor & very steep sides.
- After the disappearance of the ice, the deep sections of these long, narrow glacial troughs may be filled with water forming Ribbon lakes also known as Trough lakes or Finger Lakes.

Hanging Valleys

- The main valley is eroded much more than the tributary valley as it contains a much larger glacier.
- After the ice has been melted, a tributary valley hangs above the main valley & plunges down as a waterfall.
- Such Tributary valleys are termed as hanging valleys.
- Hanging valleys may form a natural head of water for generating hydroelectric power.




Rock Basins and Rock Steps


- A glacier erodes & excavates the bed rock in an irregular manner.
- The unequal excavation gives rise to many rock basins later filled by lakes in valley trough.
- Where a tributary valley joins a main valley, the additional weight of ice in the main valley cuts deeper into the valley floor & deepest at the point of convergence forming rock steps.
- A series of such rock steps may also be formed due to different degrees of resistance to glacial erosion of the bedrocks.

Fjord

- If the glacier flows right down to the sea, it drops its load of moraine in the sea.
- If sections break off as icebergs, moraine material will only be dropped when they melt
- Where the lower end of the trough is drowned by the sea, it forms a deep, steep side inlet called a Fjord, a typical of Norway & Chilean coast.
- Fjords are common in Norway, Greenland and New Zealand.



Glacial Landforms – Depositional



Boulder clay or Glacial till

- This is an unsorted glacial deposit comprising a range of eroded materials such as boulders, sticky clays & fine rock flour.
- It is spread out in sheets, not mounds, & forms gently undulating till or drift plains with monotonous landform.
- The degree of fertility of such glacial plains depends very much on the composition of the depositional materials.

Erratics

- Boulders of varying sizes are transported by ice & left stranded in the regions of deposition when the ice melted, called erratics because they are composed of materials entirely different from those of the regions in which they have been transported.
- Useful in tracing the source & direction of ice movements but their presence in large numbers causes hindrance in farming.

- Also known as perched blocks as sometimes they are found perched in precarious positions as the ice dropped them.

Moraines

- Moraines are made up of pieces of rock that are shattered by frost action, embedded in the glaciers & brought down the valley.
- Those that fall on the sides of the glacier form lateral moraines.
- When two glaciers converge, their inside lateral moraines unite to form a medial moraine.
- The rock fragments which are dragged along, beneath the frozen ice, are dropped when the glacier melts & spread across the floor of the valley as ground moraine.
- The glacier eventually melts on reaching the foot of the valley & the pile of transported materials left behind at the snout is terminal moraine or end moraine.
- The deposition of end moraines may be in several succeeding waves, as the ice may melt back by stages so that a series of recessional moraines are formed.

Drumlins

- **Drumlins are oval-shaped hills, largely composed of glacial drift, formed beneath a glacier or ice sheet and aligned in the direction of ice flow.**
- They are **widespread in formerly glaciated areas** and are especially **numerous in Canada, Ireland, Sweden, and Finland.**
- They are low hills up to 1.5 km long and 60 mm tall & appear steeper on the onset side & taper off at the leeward side.
- They are arranged diagonally & commonly referred as a **basket of eggs topography.**
- 'Basket of egg topography' is the **topography of drumlins, which are generally found in clusters.**
- **Drumlin fields are areas with numerous drumlins.**

Eskers

- Eskers are the sinuous ridges composed of glacial material mainly sands & gravel deposited by meltwater currents in glacial tunnels
- Glacial tunnels mark the former sites of sub glacial melt water streams
- Their orientation is generally parallel to the direction of glacial flow, and they sometimes exceed 100 kilometers in length.

Outwash Plains


- Made up of fluvio glacial deposits washed out from the terminal moraines by the streams of stagnant ice mass.
- The melt waters sort & redeposit the material mainly consisted of layers of sand and other fine sediments.
- Such plains with their sandy soils are often used for specialized kinds of agriculture, such as the potato.

Kettle lake

- Depressions are formed when the deposition takes place in the form of alternating ridges.
- Shallow, sediment-filled bodies of water formed by retreating glaciers.

Kames

- Small rounded hillocks of sand & gravel which cover part of the plain.
- Kames are often associated with kettles, and this is referred to as kame & kettle topography.
- Kame is also called Hummocks.




Periglacial Landforms

- The term **periglacial** (near-glacial) literally means **around the ice or peripheral to the margins of the glaciers** but now this term is used for both '**periglacial landscape**' and '**periglacial climate**'.
- Periglacial areas are those which are in **permanently (perennially) frozen condition but without permanent ice cover on the ground surface.**
- The periglacial climate is characterized by **mean annual temperature ranging between 1°C and – 15°C** and **mean annual precipitation of 120 mm to 1400 mm (mostly in solid form).**
- Periglacial landform is a feature resulting from the **action of intense frost, often combined with the presence of permafrost.** Periglacial landforms are restricted to areas

that **experience cold but essentially nonglacial climates**.

- Periglacial regions are **very active geomorphological areas**. The Mechanical splitting of rocks by ice (gelifraction), frost heaving of the ground (geliturbation), solifluction and nivation are all important processes. In addition, each spring, large quantities of water from melting snow and ice rapidly erode the debris scattered and moved down the slopes. Wind action is also a significant force.
- **In 1950 L.C Peltier put forward the concept of a periglacial cycle of erosion.** This is similar to the Davisian concept of the normal cycle of erosion, and the periglacial landscape attains old age after passing through the stages of youth and maturity.
- During this entire period there is erosion of the higher parts and deposition in low-lying parts, resulting in overall reduction in relief, and the slope profile gradually becomes smooth and flat. **The process by which there is gradual flattening of the surface in the periglacial regions is called 'cryoplanation'.**
- In this process there is parallel retreat of the scarp face by frost shattering and the development and gradual extension of gentle slope at its base by deposition and transport of debris.
- **Cryoplanation is thus achieved primarily by the processes of intense frost action or congelifraction and solifluction or congeliturbation.** The penultimate landform is a surface of low local relief, not controlled by any base level, and has been called a 'cryoplain' or an 'altiplain'.
- The **present-day periglacial zones are found in the Arctic regions of Alaska, Canada, Greenland, and Siberia and also in Antarctica, and the fossil zones of Pleistocene, and other past Ice Ages.**
- **Permafrost** and the **active layer** are the two most striking features of periglacial areas.
 - **Permafrost is a condition where a layer of soil, sediment, or rock below the ground surface remains frozen for a period greater than a year (for at least 2 years).** Permafrost is not a necessary condition for creating periglacial landforms. However, many periglacial regions are underlain by permafrost and it influences geomorphic processes acting in this region of the world.
 - Permafrost is found in about 25 percent of the Earth's non-glaciated land surface. It occurs

wherever the mean annual ground temperature is **less than 0° Celsius**. Common depths of permafrost are several hundred meters. Most deposits of permafrost have an upper **active layer that is between 1 to 3 meters thick**. This active layer is subject to a cyclic thaw during the summer season.



Evolution of the Periglacial Erosion Cycle

The evolution of the periglacial erosion cycles its different stages have been as:

Stage 1 (Initial surface)

- The cycle begins with the **onset of periglacial conditions**. In this initial stage **congelifraction is especially active** as a result of which there is **shattering of the bare rocks on the upper slopes, and block fields are formed** by the accumulation of angular rock fragments.

Stage 2 (Early or young stage)


- The **rock fragments move downhill** along the slope by gravity, and **talus or congeliturbate mantled slope is formed at the base of the slope**.
- During this period **nivation hollows**, as well as **cryoplanation terraces**, also begin to be formed on the higher slopes.
- Till the end of the young stage, the scarp of the terraces starts retreating by frost action, and the terrace benches are extended by scarp retreat.

Stage 3 (Mature Stage)

- With further passage of time, the **scarps are destroyed by continual recession** and in their place only residual or for features are left on the summits or slopes.
- **This is the beginning of maturity** when the original landscape is lost and the valleys and lower slopes are covered with a mantle of frost shattered and soliflucted materials.

Stage 4 (Old Stage of cryoplanation)

- Towards the end of maturity and in old age, the higher land is further reduced and levelled on account of solifluction, and the accumulation of debris goes on increasing in the neighbouring valleys and low-lying lands, and the periglacial surface is converted into an almost level plain.
- The debris becomes fine on account of continual frost weathering and solifluction (gelifluction) by further disintegration and abrasion.
- The finer debris is transported by the wind, and loess and sand dunes are formed at places, and areas of ventifacts and lag deposits develop by the blowing away of finer materials by the wind.




Taking the different slope forms found in periglacial regions as the basis, Peltier has tried to synthesize them and arrange them in a sequence. His hypothesis thus provides an overall framework for the understanding of the slope evolution in periglacial regions, but it does not provide an analysis of how exactly slope form is influenced by frost shattering and solifluction. Further, adequate attention has not been paid to other processes, especially the influence of running water. But the concept of scarp recession by frost shattering and the resultant extension of the terraces is by itself an important and useful concept that helps us to understand the formation of cryoplanation terraces, summit tors, nivation hollows & cryopediments.

Periglacial Landforms – Erosional

Freeze-Thaw Cycle

- Freeze-thaw weathering is a process of erosion that happens in cold areas where ice forms. A crack in a rock can fill with water which then freezes as the temperature

drops. As the ice expands, it pushes the crack apart, making it larger. When the temperature rises again, the ice melts, and the water fills the newer parts of the crack. The water freezes again as the temperature falls, and the expansion of the ice causes further expansion to the crack. This process continues until the rock breaks.



Nivation Hollow

- Hollows produced by snow-patch erosion or nivation are called nivation hollows which are generally found along the hillsides in various forms. They extend from a few metres to 1.5 kilometres.
- In time, these hollows may trap more snow and may deepen further with more nivation so that cirques or thermocirques are formed.
- They are classified on the basis of shape into: (i) Transverse hollows (ii) Longitudinal hollows.

Asymmetrical Valley

- An asymmetric valley is a valley that has steeper slopes on one side. Valley has one side steeper than the other, the opposing slopes having significantly contrasting angles.
- This contrast may be caused by geologic structure or variation in the nature and intensity of erosional (e.g. periglacial) processes and there may be contrasts in the vegetation on the opposing slopes.
- Such valleys are common in past and present periglacial environments, where aspect has a significant effect on the nature of frost-based processes and on the depth of the active layer.

Cryoplanation Terraces or Goletz Terraces

- Cryoplanation terraces (also known as altoplanation or goletz terraces and by several other terms) are periglacial landforms consisting of nearly horizontal bedrock surfaces or benches backed by frost-weathered bedrock cliffs.
- Cryoplanation Terraces (CTs) are erosional landforms reminiscent of giant staircases, with alternating shallow sloping treads and steep scarps leading to extensive flat summits.

Patterned Ground

- When Ice from below makes upward pressure the overlying sedimentary flat rocks are being weathered and cracks are developed, these cracks it seems are having some pattern on ground.
- A process called frost heaving is responsible for these features.

Stone Strips

- A stone stripe is an elongated concentration of mostly talus-like basalt rock found along a hillside or the base of a cliff. Many stone stripes occur without cliffs.
- **Stone stripes are thought to have been originally created by periglacial conditions of the Quaternary period during an Ice age.** It is likely their formation originates from multiple processes including **frost action, surface erosion, eluviation, and mass wasting.** However, it is likely that intense freeze and thaw cycles account for the natural sorting of the rock debris within a stone stripe, and also accounts for the shallow depth of the stripes, since frost penetration is thought to not penetrate deeper than 1 meter in the region.
- A stone stripe is identified by its lack of vegetative cover.

Pingo

- Pingo is an Eskimo word that means **isolated dome-like low mounds or hills found in permafrost areas.** This word was first used by A.E. Porsild in the year 1938. They are found in continuous as well as discontinuous permafrost areas.
- **They are abundantly found in the arctic areas (65° N) of Canada, Alaska, Greenland and Siberia.**
- They range in height from a few metres to 60 metres (sometimes they are as high as 100 metres) and from a few metres to 300 metres in diameter. Small pingos have closed tops whereas big pingos have open tops.

Hummocks

- Small upstanding wrinkles on the surface of permafrost are called hummocks. These are formed due to squeezing of the ground surface because of lateral pressure exerted by freezing of active layer.

Palsa

- A special category of hummock found in swampy areas and **composed of peats having thin ice layers inside**

is called palsa which is about 10m in height and 10 to 20 m in diameter and is mostly found in the periglacial environment of arctic and subarctic areas.

- Palsa may be destroyed by rise in water table in the nearby swamps or by fracture in its upper surface.
- It is **formed by frost heaving under the influence of ice segregation.**

Periglacial Landforms – Depositional

BlockFields or Felsenmeer

- Blockfields are areas covered by large angular blocks, traditionally believed to have been created by freeze-thaw action.
- Block field refers to the natural collection of large stone blocks at the flat surface of the hill tops in the periglacial areas. **These block fields are also called blockmeer and falsenmeer.** The stone blocks are formed due to frost weathering (conglifraction).

Patterned Ground

- **Blockfields sometimes takes some geometrical shapes** (like circles, polygons, nets, stripes and garlands) and are so systematically arranged are **called Patterned Ground.**
- Patterned Ground is **erosional** as well as **depositional** landforms in periglacial.



Tors

- Tors, one of the most controversial landforms, are **piles of broken and exposed masses of hard rocks having a crown of rock blocks of different sizes on the top and clitters (trains of blocks) on the sides.** The rock blocks, main components of tors, may be cuboidal, rounded, angular, elongated etc. in shape.
- They may be seated at the top of the hills, on the flanks of the hills or on flat basal platforms ranging from 6 m to 30 m in height. They are found in different climates varying from cold to hot and dry to humid. Though tors

have developed over almost all types of rocks but they are frequently found in the regions of granites.



Boulder Fields

- Boulder Field is a periglacial feature that **formed as a direct result of its proximity to the end moraine**. **Accumulations of rock debris in the valley floors are called stone streams or boulder fields**.
- The stone is well sorting of rock debris in the stone streams. The upper layer consists of large and coarse debris while the lower layer is dominated by fine materials. Water channel is developed between the valley walls and stone stream. Sorting of rock debris occurs through the process of frost Heaving. Stone streams move downslope due to the force of gravity, frost heaving and solifluction.




Desert Landforms/Arid

Landforms

Desert

- A desert is a **barren area of landscape** where little precipitation occurs and, consequently, living conditions are hostile for plant and animal life. **The lack of vegetation** exposes the unprotected surface of the ground to the processes of denudation. About one-third of the land surface of the world is arid or semi-arid.
- This includes much of the Polar Regions, where little precipitation occurs, and which are sometimes called polar deserts or "**cold deserts**". Deserts can be classified

by the amount of precipitation that falls, by the temperature that prevails, by the causes of desertification, or by their geographical location.




- **About 1/5 of the world's land is made up of deserts.**
- Deserts that are **absolutely barren, where nothing grows are known as true deserts**.
- Insufficient & irregular rainfall, high temperature & rapid rate of evaporation are the main causes of the desert's aridity.
- Almost all the deserts are confined within **15 degree – 30 degree parallels to N-S of the equator** known as **trade wind deserts or tropical deserts**.
- They **lie in the trade wind belt on the western parts of the continents**.
- Offshore trade winds are often bathed in cold currents which produces a desiccating (dehydrating) effect, hence moisture is not easily condensed into precipitation.

Types of Desert


- **Hamada/Rocky Desert**
 - Consist of large stretches of bare rocks, swept clear of sand & dust by wind.
 - Exposed rocks are thoroughly smoothened, polished & highly sterile
- **Reg/Stony Desert**
 - Composed of extensive sheets of angular pebbles & gravels which the wind is not able to blow off.
 - Stony deserts are more accessible than sandy deserts & large herds of camels kept there.
- **Erg/Sandy Desert**
 - Also known as the sea of sand
 - Winds deposit vast stretches of undulating sand dunes in the direction of winds
- **Badlands**
 - Consists of gully & ravines formed on hill slopes & rock surfaces by the extent of water action
 - Not fit for agriculture & survival

- Finally leads to the abandonment of the entire region by its inhabitant
- **Mountain Deserts**
 - Deserts which are found on the highlands such as on plateaus & mountain ranges, where erosion has dissected the desert highland into rough chaotic peaks & uneven ranges.
 - Their steep slopes consist of Wadis (dry valleys) with sharp & irregular edges carved due to the action of frost.



Mechanism of Desert/Arid Erosion

- **Weathering**
 - Most potent factor in reducing rocks to sand in arid regions.
 - Even though the amount of rain that falls in a desert is small, but manages to penetrate into rocks & sets up chemical reactions in various minerals it contains.
 - Intense heating during the day & rapid cooling during the night by radiations, set up stresses in already weakened rocks, hence they eventually crack.
 - When water gets into cracks of a rock, it freezes at night as the temperature drops below the freezing point & expands by 10 % of its volume.
 - Successive freezing will prise off fragments of rocks which get accumulated as screes.
 - As heat penetrates rock, its outer surface gets heated & expands, leaving its inner surface comparatively cool.
 - Hence, outer surface prise itself from the inner surface & peels




Action of Wind

- Efficient in arid regions as little vegetation or moisture to bind the loose surface materials.


It is carried out in the following ways:

- **Deflation**
 - Involves lifting & blowing away of loose materials from the ground
 - Blowing capacity depending largely on the size of the material lifted from the surface
 - Finer dust & sands may be removed miles away from their place of origin & may get deposited even outside the desert margins.
 - Deflation results in the lowering of the land surface to form large depressions called Deflation hollows.
- **Abrasion**
 - Sandblasting of rock surfaces by the wind when they hurl sand particles against them
 - This results in rock surfaces being scratched, polished & worn away
 - Abrasion is most effective near the base of the rocks, where the amount of material the wind is able to carry is greatest.
 - This explains why telegraphic poles in the deserts are protected by covering of metal for a foot or two above the ground.
- **Attrition**
 - When wind-borne particles roll against one another in the collision, they wear each other away
 - Hence their sizes are greatly reduced & grains are rounded into millet seed sand.



Desert Landforms by Wind Erosion

- **Rock pedestals/Mushroom Rocks**
 - Formed by the sandblasting effect of winds against any projecting rock masses
 - It wears down the softer layer leading to the formation of irregular edges on alternate bands of softer & harder rocks.
 - Grooves & hollows cut in the rock surfaces, carve them into grotesque-looking pillars known as rock pedestals.
 - Such rock pillars will be further eroded near their bases where friction is greatest.
 - This process of undercutting produces rocks of mushroom shape called mushroom rocks.



- **Messa and Butte**

- Mesa is a Spanish word meaning 'table'. It is a flat, table-like landmass with a very resistant horizontal top layer and very steep sides. The hard stratum on the surface resists denudation by both wind and water, and thus protects the underlying layers of rocks from being eroded away.
- Mesas may be formed in canyon regions e.g. Arizona, or on fault blocks e.g. the Table Mountain

of Cape Town, South Africa. Continued denudation through the ages may reduce mesas in the area so that they become isolated flat-topped hills called buttes.



- **Zeugen**

- Tabular masses which have a layer of soft rocks lying beneath a surface layer of more resistant rocks
- Difference in the erosional effect of the wind on soft & resistant rock surfaces, carve them into weird looking ridge & furrow landscape
- Mechanical weathering initiates their formation by opening up joints of the surface rocks
- Wind abrasion further eats into the underlying softer layer so that deep furrows are developed
- The hard rock then stand above the furrows as ridges or Zeugen
- Zeugen may stand 10 to 100 feet above the sunken furrows
- Continuous abrasion by winds gradually lowers the Zeugen & widens the furrow.



- **Yardangs**


- Yardangs look quite similar to Zeugen but instead of lying in horizontal strata upon one another, the hard & soft rocks of Yardangs are vertical bands
- Rocks are aligned in the direction of prevailing winds.
- Wind abrasion excavates the bands of softer rocks into long, narrow corridors, separating the

steep-sided over hanging ridges of hard rock called Yardangs.




- **Isenberg (Island Mountain)**

- They are basically isolated residual hills rising abruptly from the ground level
- Characterized by very steep slopes & rather rounded tops
- They are often composed of granite or gneiss
- Are probably relics of an original plateau, which has been almost entirely eroded away.



- **Ventifacts & Dreikanter**

- These are pebbles faceted by sand-blasting. They are shaped and thoroughly polished by wind abrasion to shapes resembling Brazil nuts. Rock fragments, mechanically weathered from mountains and upstanding rocks, are moved by wind and smoothed on the windward side.
- If the wind direction changes another facet is developed. Such rocks have characteristic flat facets with sharp edges.
- Amongst the ventifacts those with three wind-faceted surfaces are called dreikanter. These wind-faceted pebbles form the desert pavement a smooth, mosaic-like region, closely covered by the numerous rock fragments and pebbles.



- **Deflation Hollows**


- Winds lower the ground by blowing away the unconsolidated materials, and small depressions may form. Similarly, minor faulting can also initiate depressions and the eddying action of on-coming winds will wear off the weaker rocks until the water table is reached.
- Water then seeps out forming oasis or swamps, in the deflation hollows or depressions.
- Large areas in the western U.S.A., stripped of their natural vegetation for farming, were completely deflated when strong winds, moved materials as dust storms, laying waste crops and creating what is now known as the Great Dust Bowl.



Deserts Landforms by wind deposition


- Materials eroded & transported by winds must come to rest somewhere.
- The **finest dust travels enormous distances in the air sometimes as long as 2300 miles** before they settle down.
- The dust from Sahara desert is sometimes blown across the **Mediterranean to fall as blood rains in Italy or on the glaciers of Switzerland.**
- **Dust that settles in Hwang Ho basin (also known as Hwangtu – the yellow earth) from the Gobi desert has been accumulated over past centuries** to a depth of several hundred feet
- As wind borne **materials are shifted according to their coarseness**, it can be expected that the coarser sands will be too heavy to be blown out of desert limits.

- They remain as dunes or other depositional landforms within desert themselves.




Dunes

- Dunes are, in fact, hills of sand formed by the accumulation of sand and shaped by the movement of winds.
- They may be active or live dunes, constantly on the move, or inactive fixed dunes, rooted with vegetation.
- Dunes are most well represented in the erg desert where a sea of sand is being continuously moved, reshaped, and re-deposited into a variety of features.
- Two most common types of dunes are **Barchan & Seifs**.
- **Barchan:**
 - These are crescentic or moon-shaped dunes that occur individually or in groups. They are live dunes that advance steadily before winds that come from a particular prevailing direction. They are most prevalent in the deserts of Turkestan and in the Sahara.
 - Barchans are initiated probably by a chance accumulation of sand at an obstacle, such as a patch of grass or a heap of rocks. They occur transversely to the wind, so that their horns thin out and become lower in the direction of the wind due to the reduced frictional retardation of the winds around the edges.
 - The windward side is convex and gently sloping while the leeward side, being sheltered, is concave and steep (the slip-face).
 - The crest of the sand dune moves forward as more sand is accumulated by the prevailing wind.
 - The sand is driven up the windward side and, on reaching the crest, slips down the leeward side so that the dune advances.
 - The migration of the barchans may be a threat to desert life for they may encroach on an oasis burying palm trees or houses.
- Long-rooted sand-holding trees and grasses are therefore planted to halt the advance of the dunes thus preventing areas of fertile land from being devastated.
- Under the action of winds, barchans take a chaotic changing pattern. Several barchans may coalesce into a line of irregular ridges, ever-changing with the direction of the winds. Ergs or sandy deserts are thus most difficult to cross.



Seifs or Longitudinal Dunes:

- They are long, narrow ridges of sand, often over a hundred miles long lying parallel to the direction of the prevailing winds. The Crestline of the seif rises and falls in alternate peaks and saddles in regular successions like the teeth of a monstrous saw.
- The dominant winds blow straight along the corridor between the lines of dunes so that they are swept clear of sand and remain smooth. The eddies that are set up blow towards the sides of the corridor, and, having less power, drop the sand to form the dunes.
- In this manner, the prevailing winds increase the length of the dunes into tapering linear ridges while the occasional crosswinds tend to increase their height and width. Extensive seif dunes are found in the Sahara Desert, south of the Qattara Depression; e.g. the Thar Desert and the West Australian Desert.




D Longitudinal dunes (seifs)

Loess

- The fine dust blown beyond the desert limits is deposited on neighboring lands as loess. It is a yellow, friable

material and is usually very fertile. Loess is in fact, fine loam, rich in lime, very coherent, and extremely porous. Water sinks in readily so that the surface is always dry.

- Streams have cut deep valleys through the thick mantle of soft loess and badland topography may develop. It is so soft that roads constructed through a loess region soon sink and their walls rise steeply. The most extensive deposit of loess is found in northwest China in the loess plateau of the Hwang- Ho basin.
- It is estimated to cover an area of 250,000 square miles, and the deposits have accumulated to a depth of 200 to 500 feet! In China, such yellowish wind-borne dust from the Gobi Desert is called 'Hwangtu' — the yellow earth! But the original term loess actually comes from a village in Alsace, France bearing that name, where such deposits occurred.
- Similar deposits also occur in some parts of Germany France and Belgium and are locally called Limon They are also wind-borne but were blown from material deposited at the edge of ice sheets during the Ice Ages. In parts of the Mid-West, U.S.A. loess was derived from the ice sheets which covered northern North America and is termed adobe.




Landform of Water Actions in Desert

- Few deserts in the world are entire without rain or water. The annual precipitation may be small and comes in irregular showers. But thunderstorms do occur and the rain falls in torrential downpours, producing devastating effects.
- A single rainstorm may bring several inches of rain within a few hours, drowning people who camp in dry desert streams and flooding mud-baked houses in the oasis.
- As deserts have little vegetation to protect the surface soil, large quantities of rock wastes are transported in the sudden raging torrents or flash-floods. Loose gravels, sand, and fine dusts are swept down the hillsides.
- They cut deep gullies and ravines forming bad-land topography. Subsequent downpours widen and deepen the gullies when they wash down more soft rocks from

the surface. There is so much material in the flash floods that the flow becomes liquid mud.


- When the masses of debris are deposited at the foot of the hill or the mouth of the valley, an alluvial cone or fan or 'dry delta' is formed, over which the temporary stream discharges through several channels, depositing more material.
- The pasty alluvial deposits are subjected to rapid evaporation by the hot sun and downward percolation of water into the porous ground, and soon dry up leaving mounds of debris. Apart from gullies there are many larger dry channels or valleys.
- Temporary lakes(Playas):**
 - Also known as **Playas, Salina or Salars.**
 - Formed in arid or semi-arid areas by intermittent streams flowing into depressions
 - Contain high percentage of salts due to high evaporation & lower precipitation.
 - The playa plain covered up by salts is called alkali flats.




Bajada & Pediment: The floor of the desert depression is made up of two features viz. Bajada & Pediment.

- Bajada**
 - Depositional feature made up of alluvial material lay down by intermittent streams.
 - Bajada is formed by the coalescence of alluvial fans
 - These fan-shaped deposits form from the deposition of sediment by a stream from upland region onto flat land at the base of a mountain
 - Bajadas are common in arid areas where a large quantity of sediment is deposited by flash floods
 - Bajadas frequently contain playa lakes
- Pediment**
 - An erosional plain formed at the base of the surrounding mountain scarps -steep slope.

- They are gently inclined rocky floors close to the mountains at their foot with or without a thin cover of debris.
- They form through the erosion of mountain front through a combination of lateral erosion by streams and sheet flooding.
- Through parallel retreat of slopes, the pediments extend backwards at the expense of mountain front
- Gradually, the mountain gets reduced leaving an inselberg which is a remnant of the mountain.
- That's how the high relief in desert areas is reduced to low featureless plains called pediplains.



- **Alluvium fans**
 - Alluvium fans are cone-shaped accumulations of sand that form at the end of a wadi or valley. A wadi is a narrow, dry valley characterized by its intermittent water flow, which only occurs during heavy rainfall. For the majority of the time, the valley remains dry and parched. However, during heavy downpours, the wadi can fill up with water, carrying alluvium from the higher ground downstream as sheet wash.
 - As the wadi comes to an end and opens up into a larger space, the flow of water loses energy, causing the transported alluvium to spread out and form a fan-like shape. This process results in the creation of alluvium fans at the exit of the wadi or valley.
- **Canyons/Gorges**
 - Gorges, also known as canyons in America, are deep and narrow valleys that have been carved out and eroded by rivers flowing through desert landscapes.
 - The Grand Canyon in Arizona, USA, is an example of such a formation, created by the vertical erosion of sedimentary rock layers by the Colorado River over millions of years.



Conclusion

Fluvial, marine, and karst landforms are created by various erosional and depositional processes involving water, wind, and other geological factors. These landforms, such as headlands, bays, caves, arches, stacks, stumps, spits, and alluvium fans, are found all over the world and contribute to the diverse and fascinating landscape of our planet. Understanding the processes that shape these landforms is essential for studying geomorphology, as well as for sustaining and preserving these unique features for future generations.

What are the main differences between fluvial, marine, and karst landforms?

Fluvial landforms are created by the action of rivers and running water, while marine landforms are shaped by the processes occurring along coastlines and in the ocean. Karst landforms are formed by the dissolution of soluble rocks, such as limestone, resulting in features like caves and sinkholes.

How do headlands and bays form on coastlines?

Headlands and bays form when there is a variation in the resistance of rocks along a coastline. Softer rocks, such as clay and sand,

erode more quickly and form bays, while harder rocks, such as chalk, erode more slowly and form headlands.

What factors contribute to the formation of karst topography?

Karst topography develops in areas with dense, soluble rocks (e.g., limestone), low water tables, and moderate to heavy rainfall. These factors encourage the rapid downward movement of groundwater, promoting the dissolution of bedrock and the formation of features like caves and sinkholes.

What are some examples of depositional landforms in marine environments?

Some examples of depositional marine landforms include beaches, spits, and hooks. These features are created by the accumulation of sediments deposited by the sea, often in sheltered areas or where there are changes in the shape of the coastline.

What is the difference between a gorge and a canyon?

Gorges and canyons both refer to deep, narrow valleys that have been carved out and eroded by rivers. The main difference between them is regional terminology, with "gorge" being more commonly used in Europe and "canyon" in America. The Grand Canyon in Arizona, USA, is an example of such a formation.

1. What are fluvial landforms?

Ans. Fluvial landforms are landforms that are created by the action of running water, such as rivers and streams. Examples of fluvial landforms include valleys, canyons, floodplains, and meanders.

2. What are some examples of marine landforms?

Ans. Marine landforms are landforms that are created by the action of the ocean or sea. Examples of marine landforms include beaches, cliffs, sea stacks, and coral reefs.

3. What are the types of erosion?

Ans. There are several types of erosion, including: 1. Water erosion: This occurs when running water, such as rivers or rainfall, wears away the land and carries away sediment. 2. Wind erosion: This occurs when wind blows over the land and picks up and carries away loose particles of soil and sand. 3. Glacier erosion: This occurs when a glacier moves across the land, scraping and plucking away rocks and sediment. 4. Coastal erosion: This occurs when the action of waves and currents wears away the land along a coastline.

4. What are karst landforms?

Ans. Karst landforms are created by the dissolution of soluble rocks, such as limestone or dolomite, by water. They often feature unique formations such as sinkholes, caves, and underground drainage systems.

5. What are some examples of desert landforms?

Ans. Desert landforms, also known as arid landforms, include: 1. Sand dunes: These are mounds or hills of sand that are formed by wind or water erosion and deposition. 2. Desert pavement: This is a flat surface made up of closely packed pebbles and stones, formed by wind erosion removing finer particles. 3. Playas: These are flat, dry lake beds that form in desert basins. 4. Arroyos: These are dry riverbeds that only fill with water during heavy rainfall or flash floods.

Factors Controlling Landform Development

What are Landforms?

- Landforms are three-dimensional structures found on the Earth's surface, ranging from small to medium-sized areas. These natural formations can take various shapes, such as mountains, plains, plateaus, valleys, and hills, among others.

Before going to the factors that control the development of landforms, we need to know the answers to some of the main questions like -

Why do landforms develop? Why is the Earth's surface uneven?

Landforms develop and the Earth's surface is uneven due to several factors, primarily involving the dynamic nature of the Earth's crust and the varying intensities of internal and external forces.

- The Earth's crust is constantly moving, both vertically and horizontally. This movement contributes to the formation and reshaping of various landforms over time.
- The geothermal gradient, or the variation in the Earth's internal heat, causes uneven distribution of energy within the planet. This energy generates internal forces that create different types of high and low land surfaces, which are collectively known as landforms.
- External forces, primarily driven by sunlight, also influence the Earth's surface. These forces include weathering, erosion, and mass movement, which alter the structure of landforms over time. For instance, large mountains may gradually erode into smaller hills and eventually become flat plains due to the continuous action of these external forces.
- The intensity of both internal and external forces varies across the Earth's surface. This variation leads to the development of diverse landforms in different regions.

Which factors primarily contribute to the development of landforms and the unevenness of the Earth's surface? **A.**

The Earth's crust movement and internal heat **B.** Rock composition and relief **C.** Climate and external forces **D.** Biological activities and human activities

In summary, the development of landforms and the unevenness of the Earth's surface are primarily due to the dynamic nature of the Earth's crust and the varying intensities of internal and external forces. Other factors also contribute to the formation and evolution of landforms, but these primary forces play a significant role in shaping the Earth's diverse landscapes.

Following are the controlling factors of landform development:

- **Rock Composition or Parent Rocks:** The rate at which landforms develop or form is influenced by their rock composition or parent rocks. This is due to the variation in the size and composition of rocks, which determines their hardness or softness. For example, quartz is a hard rock that erodes and weathers slowly, while limestone is a soft rock that undergoes weathering and erosion more quickly. This results in the rapid degradation of landforms composed of softer rocks. Each rock type has unique characteristics that make it more or less resistant to different weathering agents, leading to the formation of diverse landforms.
- **Relief:** Relief refers to the elevation difference between the highest and lowest points in an area. Landforms develop more quickly in areas with greater relief due to increased erosion. For instance, landform development in the Himalayan region is faster than in the Aravalli region because the relief is higher. Greater relief also results in stronger gravitational forces, leading to mass movements and large-scale landslides.
- **Geological Formation:** The geological structure of rocks, such as their layers, faults, orientations, inclinations, joint presence or absence, hardness or softness, and chemical sensitivity, also affects landform development.
- **Exogenous Agents:** External factors like water, wind, ice, currents, waves, and tides also influence the development process of landforms.
- **Climate:** Temperature and humidity are crucial climate factors that impact landform development. High humidity enhances chemical and biological weathering processes, while high rainfall and wind speeds increase erosion

activities. Large temperature variations cause rocks to expand and contract, leading to their fragmentation. Landform development processes and intensities vary across different climatic zones and even within the same climate zone due to varying rainfall and temperature intensities.

- **Energy:** Energy sources, both internal and external, affect landform development. Internal energy refers to differences between convergent and divergent convection cells within the Earth, while external energy includes temperature variations, wind, and pressure changes.
- **Biological Activities:** The presence of vegetation can accelerate chemical weathering processes as plants release acids and moisture onto rocks. Conversely, a lack of vegetation can increase erosion, leading to the rapid degradation of elevated landforms.
- **Human Activities:** Human actions, such as agriculture, dam construction, infrastructure development, and brick-making, can also influence landform development.

How does the relief of an area influence landform development? **A.** It determines the rock composition of the area **B.** It affects the climate of the area **C.** It impacts the rate of erosion and mass movements **D.** It influences biological activities in the area

Conclusion

Landforms are natural structures found on the Earth's surface and are formed and reshaped by various factors. These factors include the rock composition, relief, geological formations, exogenous agents, climate, energy, biological activities, and human actions. The dynamic nature of the Earth's crust and the

varying intensities of internal and external forces significantly influence the development of diverse landforms across the planet. Understanding these factors and their impact on landform development is crucial for predicting and managing the Earth's changing landscapes.

What are the main types of landforms?

Some of the main types of landforms include mountains, plains, plateaus, valleys, and hills. These natural formations vary in size and shape and are found on the Earth's surface.

What are the internal and external forces that contribute to landform development?

Internal forces include the movement of the Earth's crust and the geothermal gradient, which generates energy that shapes the Earth's surface. External forces, driven by sunlight, include weathering, erosion, and mass movement, which alter the structure of landforms over time.

How does rock composition affect landform development?

The rock composition or parent rocks influence the rate of landform development due to the variation in the size and composition of rocks, which determines their hardness or softness. Harder rocks like quartz erode and weather slowly, while softer rocks like limestone undergo weathering and erosion more quickly.

How does climate influence landform development?

Temperature and humidity are crucial climate factors that impact landform development. High humidity enhances chemical and biological weathering processes, while high rainfall and wind speeds increase erosion activities. Large temperature variations cause rocks to expand and contract, leading to their fragmentation.

How do human activities affect landform development?

Human actions, such as agriculture, dam construction, infrastructure development, and brick-making, can influence landform development by altering natural processes like erosion and weathering. These activities can lead to the rapid degradation of elevated landforms and the alteration of the Earth's surface.

5. How do human activities impact landform development?



Ans. Human activities can significantly impact landform development. Mining operations can alter the topography by removing large amounts of soil and rocks, leading to the formation of mining pits or spoil heaps. Deforestation can result in increased soil erosion and landslides, changing the landscape and affecting landform development. Construction activities like dam building can modify river courses and create reservoirs, altering the natural flow of water and impacting landforms downstream. Human-induced climate change can also have long-term effects on landforms through sea-level rise, increased erosion, and altered precipitation patterns.

1. What are the factors controlling landform development?

Ans. Factors controlling landform development include geology, climate, tectonic activity, erosion, and human activities. Geology determines the type of rocks and their properties, which influence landform formation. Climate affects landforms through processes like weathering, erosion, and deposition. Tectonic activity, such as plate movement and volcanic activity, can create mountain ranges and other landforms. Erosion by rivers, glaciers, wind, and waves also plays a significant role. Human activities like mining, deforestation, and construction can alter landforms.

2. How does geology influence landform development?

Ans. Geology influences landform development by determining the type of rocks present and their properties. Different types of rocks have varying resistance to weathering and erosion. For example, hard rocks like granite are more resistant and can form mountains, while softer rocks like sandstone may erode into valleys or cliffs. Geology also affects the formation of specific landforms like caves, canyons, and coastal features. The composition and structure of rocks influence the processes of weathering and erosion, ultimately shaping the landforms we see.

3. What role does climate play in landform development?

Ans. Climate plays a crucial role in landform development through processes like weathering, erosion, and deposition. Temperature, precipitation, and wind patterns determine the rate and intensity of these processes. In humid regions with high rainfall, weathering and erosion are more pronounced, leading to the formation of valleys, gorges, and waterfalls. In arid regions, wind erosion and deposition can create desert landscapes like sand dunes. Cold climates with glaciers can carve out U-shaped valleys and create glacial landforms like moraines and cirques.

4. How does tectonic activity influence landform development?

Ans. Tectonic activity, such as plate movements and volcanic activity, can greatly influence landform development. When tectonic plates collide, they can create mountain ranges like the Himalayas or the Andes. This process, known as orogenesis, involves folding, faulting, and uplift of rocks, resulting in the formation of large-scale landforms. Volcanic activity can also shape the landscape by creating volcanic mountains, lava plateaus, and volcanic landforms like calderas and lava flows. Tectonic activity can be responsible for both gradual and rapid landform changes.

Geo-Hydrology and Hydrological Cycle

Overview - Geo-Hydrology and Hydrological Cycle

Geo-hydrological Investigation is a broad term referring to the study of the availability of water resources in specific areas. In India, this type of investigation encompasses various aspects, including:

- Assessing groundwater levels before and after the monsoon season
- Determining soil moisture content

- Evaluating the quality and quantity of water in various sources, such as rivers, lakes, groundwater, and reservoirs
- Analyzing the impact of urbanization on water demand
- Evaluating the influence of industrial and other economic activities on water demand
- Developing strategies for flood and drought control

Mapping plays a crucial role in geo-hydrological investigations, as it offers several benefits:

- Maps provide a clear and concise representation of complex water resource information, simplifying the investigation process.
- Topographic maps help identify flood- and drought-prone areas, allowing for better management and mitigation strategies. For instance, low elevation areas are more susceptible to flooding, while high elevations are more prone to drought.
- Mapping assists in strategic planning for projects such as river linking, by outlining routes and identifying potentially affected areas like national parks and wildlife sanctuaries.
- The use of maps aids in pollution mitigation and environmental conservation by providing visual data on water resources and their surrounding areas.

In summary, geo-hydrological investigation is a comprehensive study of water resource availability in specific regions. In India, this includes assessing groundwater levels, soil moisture, water quality and quantity, and the impact of human activities on water demand.

Hydrological Cycle Explained

- The hydrological cycle, also known as the water cycle, is a continuous process in which water is exchanged between the atmosphere, oceans, and continents through evaporation, transpiration, condensation, and precipitation. This cycle is essential for maintaining a constant amount of water vapor in the atmosphere, which in turn supports the Earth's climate and ecosystems.
- Evaporation is the process where water from bodies of water, such as oceans and lakes, is transformed into vapor and released into the atmosphere. Transpiration occurs when plants release water vapor through their leaves. Together, these processes are called evapotranspiration, which contributes to the moisture content in the atmosphere.
- In the next stage, this evaporated moisture condenses to form clouds. The process of condensation occurs when water vapor in the atmosphere cools down and turns back into liquid droplets, forming clouds. Following condensation, precipitation occurs when these water droplets, now in the form of rain, snow, or other forms, fall back to the Earth's surface.
- The hydrological cycle is a continuous process with no specific beginning or end, as water is constantly circulating within the Earth's system. This cycle plays a crucial role in shaping the Earth's climate and supporting life on our planet. Factors such as the sun's energy and air currents drive the hydrological cycle, enabling the

various stages to occur and maintain balance within the Earth's water system.


What are the four main stages of the hydrological cycle? **A.** Evaporation, condensation, precipitation, and runoff **B.** Evaporation, transpiration, precipitation, and infiltration **C.** Transpiration, evaporation, condensation, and runoff **D.** Evaporation, transpiration, condensation, and precipitation

Process of the Water Cycle

The water cycle is a continuous process that involves several stages, including evaporation and transpiration, condensation, precipitation, and runoff and infiltration.

1. **Stage I: Evaporation and Transpiration:** The first stage of the water cycle involves the sun's energy heating up bodies of water such as lakes, rivers, and oceans, causing the temperature of the water to rise. This increase in temperature leads to the evaporation of some water into the air. Rising air currents then carry the water vapor up into the atmosphere. Simultaneously, plants and trees lose water to the atmosphere through a process called transpiration, in which they release water vapor that also rises into the sky.
2. **Stage II: Condensation:** In the second stage, the water vapor that has risen into the atmosphere cools down due to the lower surrounding temperatures. As a result, the water vapor transforms back into a liquid through a process called condensation.
3. **Stage III: Precipitation:** During the third stage, wind movements cause the cloud particles to collide with each other. This causes the clouds to become heavy with water, eventually leading to the formation of rain-bearing clouds. The water then falls back onto the Earth's surface through a process known as precipitation, which can take the form of rain, hail, snow, or sleet, depending on temperature conditions.
4. **Stage IV: Runoff and Infiltration:** After precipitation, the water either runs off into oceans and rivers, flows along the ground surface, or is absorbed into the soil through a process called infiltration. Once the water has gone through these steps, it is ready to evaporate once more and repeat the earlier stages of the cycle.

By undergoing all these processes, the hydrological or water cycle continually replenishes the Earth's supply of fresh water.



Water Vapour in the Atmosphere

The atmosphere contains varying amounts of water vapor, ranging from zero to four percent by volume. On average, water vapor makes up approximately 2% of the atmospheric composition. A hygrometer is an instrument used to measure humidity, or the amount of water vapor present in the atmosphere.

- Significance of Atmospheric Moisture:** Water vapor plays a critical role in regulating the Earth's temperature by absorbing both incoming and terrestrial radiation. This absorption process contributes to the heating of the Earth's surface. The amount of water vapor in the atmosphere also determines the energy available for the formation of storms and cyclones. Additionally, atmospheric moisture influences the rate at which the human body cools down in response to temperature fluctuations.
- Humidity:** Humidity refers to the amount of water vapor present in the air or atmosphere. It is often expressed as relative humidity, which is a percentage of the maximum amount of water vapor that the air can hold at a specific temperature.

Why is atmospheric moisture important? **A.** It helps regulate Earth's temperature and influences storm formation **B.** It is essential for plant growth and soil fertility **C.** It assists in the breakdown of pollutants in the atmosphere **D.** It serves as a source of drinking water for humans and animals

Conclusion

Geo-hydrological investigation is essential for understanding water resource availability and management in specific regions, such as India. The hydrological cycle, a continuous process involving evaporation and transpiration, condensation,

precipitation, and runoff and infiltration, plays a crucial role in maintaining the Earth's climate and ecosystems. Water vapor in the atmosphere, which makes up approximately 2% of its composition, is significant in regulating temperature and influencing weather patterns. Overall, studying these processes and their impacts is vital for sustainable water resource management and environmental conservation.

What is the primary objective of a geo-hydrological investigation?

The primary objective of a geo-hydrological investigation is to study the availability of water resources in specific areas. This includes assessing groundwater levels, soil moisture content, the quality and quantity of water in various sources, and evaluating the impact of human activities on water demand.

How does mapping benefit

geo-hydrological investigations?

Mapping plays a crucial role in geo-hydrological investigations by providing a clear and concise representation of complex water resource information, helping to identify flood- and drought-prone areas, assisting in strategic planning for projects, and aiding in pollution mitigation and environmental conservation.

What are the main stages of the hydrological cycle?

The main stages of the hydrological cycle include evaporation and transpiration, condensation, precipitation, and runoff and infiltration. These processes work together to maintain the Earth's water system and support the planet's climate and ecosystems.

How does the water cycle help maintain the Earth's water supply?

The water cycle is a continuous process that helps maintain the Earth's water supply by constantly circulating water within the Earth's system through evaporation, transpiration, condensation, precipitation, and runoff and infiltration. This process replenishes the Earth's supply of fresh water and supports life on our planet.

What is the significance of atmospheric moisture in the hydrological cycle?

Atmospheric moisture, or water vapor, plays a critical role in regulating the Earth's temperature by absorbing both incoming and terrestrial radiation. The amount of water vapor in the atmosphere also determines the energy available for the formation of storms and cyclones, and influences the rate at which the human body cools down in response to temperature fluctuations.

1. What is geo-hydrology?

Ans. Geo-hydrology is a branch of hydrology that focuses on the study of groundwater, its occurrence, movement, and quality in the Earth's subsurface. It involves understanding the geological formations and their impact on groundwater resources.

2. What is the hydrological cycle?

Ans. The hydrological cycle, also known as the water cycle, is a continuous process through which water circulates between the Earth's surface and the atmosphere. It involves various processes such as evaporation, condensation, precipitation, and runoff, which collectively contribute to the distribution and availability of water on Earth.

3. How does the water cycle work?

Ans. The water cycle starts with the evaporation of water from oceans, lakes, and rivers, which then rises into the atmosphere as water vapor. The water vapor condenses to form clouds, and eventually, precipitation occurs in the form of rain, snow, or hail. This precipitation is then stored in various reservoirs such as lakes, rivers, and groundwater. The stored water eventually returns to the atmosphere through evaporation and transpiration, completing the cycle.

4. Why is understanding water vapor in the atmosphere important?

Ans. Understanding water vapor in the atmosphere is crucial because it plays a significant role in the Earth's climate system. Water vapor is the most abundant greenhouse gas and acts as a natural insulator, trapping heat in the atmosphere. It also influences the formation of clouds and precipitation, which are essential for the water cycle and overall weather patterns.

5. How does geo-hydrology contribute to water resource management?

Ans. Geo-hydrology provides valuable insights into the distribution, movement, and quality of groundwater resources. This information is essential for effective water resource management, as it helps in assessing the availability and sustainability of groundwater sources. Geo-hydrological studies also aid in identifying potential contamination risks and developing strategies for groundwater protection and conservation.

Economic Geology and Environment

Introduction to Economic Geology

- Economic geology is a scientific field that concentrates on investigating and exploring ore deposits for financial gain.
- Ore deposits are geological formations containing valuable minerals that can be extracted for profit.

For example, while iron oxide may be present in large quantities in a particular area, it might not be lucrative on its own.

- Therefore, it is crucial to comprehend the properties and locations of ore deposits to maximize their economic benefits.
- In essence, ore bodies and deposits are mineral-rich rocks that can be mined for profit.
- The Earth's crust contains numerous resources that humans utilize to power their daily lives.
- The combination of geological and economic exploration forms the foundation of a thriving resource industry, which constitutes a significant portion of worldwide exports and drives economic growth in nations such as Canada, Australia, Norway, and South Africa.

benefits not only the present generation but also future generations.

- By continually discovering and developing new methods and technologies, they contribute to the long-term sustainability of our planet and its resources.
- In summary, Economic Geology is a significant field of study that plays a vital role in securing the resources necessary for human progress while also promoting responsible and sustainable extraction practices for the benefit of future generations.

Why is Economic Geology considered significant in shaping a sustainable and prosperous future for humanity?

Significance of Economic

Geology

- Economic Geology plays a crucial role in shaping a sustainable and prosperous future for humanity, as it focuses on the discovery and responsible extraction of natural resources. These resources are essential for providing people with a higher quality of life and contribute to various industries and economic growth.
- As an applied science, Economic Geology often leads to the commercial exploitation of resources, making it an integral part of industrial development.
- This field of study relies heavily on public funding from government agencies such as the National Science Foundation, National Aeronautics and Space Administration, Department of Energy, Geological Survey Canada, and universities.
- As a result, researchers and professionals in this field are well-versed in navigating both commercial and scientific environments.
- Economic Geologists play a vital role in finding new and innovative ways to extract natural resources responsibly, ensuring that their work

- A.** It focuses on discovering and extracting natural resources that are essential for providing a higher quality of life.
- B.** It contributes to the fashion industry.
- C.** It helps in the development of new sports equipment.
- D.** It is essential for the growth of the entertainment industry.

Advantages of Economic Geology

- Economic geology offers numerous benefits, primarily due to the collaboration between economists, chemists, and physicists. These professionals from different fields can work together as a cohesive team, utilizing their unique research areas that intersect with economic geology.
- By pooling their expertise, these scientists can efficiently conduct necessary calculations to locate ore deposits, saving both time and money.

- This interdisciplinary approach also eliminates the need for repeated testing, as each team member can contribute their specialized knowledge to the project.
- Furthermore, the synergy between geology, chemistry, and physics leads to a more comprehensive understanding of fundamental processes and intricate chemical details.
- Overall, economic geology promotes efficiency, cost-effectiveness, and a deeper comprehension of the subject matter through the collaboration of experts from diverse scientific backgrounds.

Economic and Environmental

Geology

- Economic and environmental geology may focus on different problems, but there are areas where they can collaborate and complement each other. Environmental geologists need to understand the economic implications of their work and the decisions they make throughout their careers.
- Environmental geologists play a crucial role in making well-informed decisions based on their understanding of Earth's processes and how humans respond to this information. It is essential for them to adopt an economic perspective as well, as their expertise can significantly influence decision-making related to economic activities and their locations.
- As highlighted in an economic geology journal, the work of environmental geologists can greatly benefit the economy. The integration of environmental science and geology helps ensure clean water availability, economic growth, environmental preservation, sustainable development, and social equity for all.

What are some advantages of Economic Geology?

- A.** It promotes efficiency and cost-effectiveness in locating ore deposits.
- B.** It helps in developing new flavors of ice cream.
- C.** It encourages the growth of the tourism industry.
- D.** It enhances the performance of electronic devices.

Conclusion

Economic Geology is a crucial field of study within the realm of geology, as it plays a significant role in determining the success and prosperity of countries. This branch of geology focuses on the investigation and analysis of geological processes that lead to the creation, transformation, and distribution of mineral resource deposits. The ultimate objective is to locate ore deposits and generate profit from them.

The study of Economic Geology encompasses various aspects, including surface mapping, geological mapping (which may involve rock units), mineralogy, geochemistry, statistical analysis, and numerical modeling. It is considered an intriguing field of science by many people due to its direct impact on our daily lives. For instance, the exploration and extraction of precious metals like gold and silver significantly contribute to the economy of nations worldwide.

The significance of mineral production has grown exponentially in contemporary society, as it serves as a critical component in various industries and applications. Therefore, the study of Economic Geology remains essential for the continued growth and development of economies across the globe.

What is the main goal of economic geology?

The primary objective of economic geology is to investigate and analyze geological processes that lead to the creation, transformation, and distribution of mineral resource deposits. The ultimate goal is to locate ore deposits and generate profit from their extraction.

How do economic geologists contribute to sustainable development?

Economic geologists play a vital role in finding new and innovative ways to extract natural resources responsibly, ensuring that their work benefits not only the present generation but also future generations. They continually discover and develop new methods and technologies that contribute to the long-term sustainability of our planet and its resources.

How does the interdisciplinary approach of economic geology benefit the field?

The interdisciplinary approach of economic geology promotes efficiency, cost-effectiveness, and a deeper comprehension of the subject matter through the collaboration of experts from diverse scientific backgrounds, such as economists, chemists, and physicists. This collaborative effort allows for efficient calculations to locate ore deposits, saving both time and money.

How do environmental geologists and economic geologists collaborate?

Environmental geologists need to understand the economic implications of their work and the decisions they make throughout their careers. By adopting an economic perspective, their expertise can significantly influence decision-making related to economic activities and their locations. The integration of environmental science and geology helps ensure clean water availability, economic growth, environmental preservation, sustainable development, and social equity for all.

How does the study of economic geology impact our daily lives?

The exploration and extraction of precious metals like gold and silver significantly contribute to the economy of nations worldwide. The significance of mineral production has grown exponentially in contemporary society, as it serves as a critical component in various industries and applications. Therefore, the study of economic geology remains essential for the continued growth and development of economies across the globe.

1. What is economic geology and why is it significant?



Ans. Economic geology is a branch of geology that focuses on the study of Earth materials that can be economically extracted for their valuable resources. This field plays a crucial role in identifying and evaluating mineral deposits, petroleum reserves, and other natural resources. It helps in understanding the distribution, origin, and characteristics of these resources, which is essential for their sustainable exploitation. Economic geology is significant as it provides the foundation for resource exploration, mine planning, and resource management, contributing to the development of industries and economies worldwide.

2. What are the advantages of economic geology?



Ans. Economic geology offers several advantages. Firstly, it aids in the discovery and assessment of mineral deposits, oil and gas reserves, and other valuable resources, which are vital for meeting the increasing demand for raw materials. Secondly, it provides valuable information for resource planning and management, ensuring the sustainable utilization of Earth's resources. Moreover, economic geology supports the development of mining and petroleum industries, creating employment opportunities and contributing to economic growth. Lastly, it helps in understanding geological processes, which can have implications for hazard assessment and environmental management.

3. How does economic geology relate to the environment?

Ans. Economic geology and environmental geology are closely linked. Economic geology involves the study of Earth materials and resources, while environmental geology focuses on the impact of human activities on the environment. The extraction and utilization of natural resources, such as mining and drilling, can have significant environmental consequences. Therefore, economic geologists play a crucial role in ensuring responsible resource extraction practices that minimize environmental degradation. They assess the potential environmental impacts of resource extraction, develop strategies for environmental protection and restoration, and contribute to sustainable development.

4. What role does economic geology play in sustainable development?

Ans. Economic geology plays a vital role in sustainable development. It helps in identifying and evaluating mineral deposits and other valuable resources, ensuring their sustainable extraction. By understanding the distribution and characteristics of resources, economic geologists contribute to efficient resource management, preventing resource depletion and promoting long-term sustainability. They also contribute to environmental protection by assessing the potential impacts of resource extraction and suggesting mitigation measures. Furthermore, economic geology supports the development of industries and economies, providing opportunities for economic growth and improving living standards.

5. How can economic geology contribute to energy transition and climate change mitigation?

Ans. Economic geology can contribute to energy transition and climate change mitigation in several ways. Firstly, it helps in the exploration and assessment of renewable energy resources such as geothermal, wind, and solar energy. By identifying suitable locations for renewable energy projects, economic geologists support the transition from fossil fuels to clean energy sources. Secondly, economic geology plays a role in the exploration and development of critical minerals required for clean technologies, including lithium for batteries and rare earth elements for wind turbines and electric vehicles. By ensuring a sustainable supply of these minerals, economic geology supports the development of low-carbon technologies. Lastly, economic geologists contribute to carbon capture and storage projects by identifying and characterizing suitable geological formations for carbon sequestration, aiding in climate change mitigation efforts.
